



Psychometric properties and measurement invariance of a European Portuguese version of the fear of COVID-19 scale

Margarida Jarego¹, Alexandra Ferreira-Valente^{1,2,3}, Rui Miguel Costa¹, Mafalda Tavares⁴, and José Pais-Ribeiro^{1,5}

1 William James Center for Research, Ispa – University Institute, Lisbon, 1149-041 (Portugal)

2 Department of Rehabilitation Medicine, University of Washington, Seattle, WA, 98104 (U.S.A)

3 Research Center for Human Development, Faculty of Education and Psychology, Universidade Católica Portuguesa, Porto, 4169-005 (Portugal)

4 Ispa – University Institute, Lisbon, 1149-041 (Portugal)

5 School of Psychology and Education Sciences, University of Porto, Porto, 4200-135 (Portugal)

Título: Propiedades psicométricas e invariancia de medida de una versión en portugués europeo de la escala de miedo de la COVID-19.

Resumen: El miedo generalizado de la COVID-19 parece haber exacerbado el impacto negativo de la pandemia. Por eso, es relevante monitorear el miedo de la COVID-19 y su asociación con la salud mental, el bienestar y los comportamientos de las personas, siendo necesarias medidas válidas y fiables de miedo de la COVID-19. Este estudio tuvo como objetivo evaluar las propiedades psicométricas de una versión en portugués europeo de la escala Fear of COVID-19 (FCV-19S-P). Un objetivo secundario fue evaluar la invariancia de medición transversal multigrupo de FCV-19S-P (mujeres vs. hombres). Una muestra de 572 adultos portugueses (72 % mujeres) completaron el FCV-19S-P y medidas de depresión, ansiedad y estrés. Los resultados del estudio respaldaron la validez de esta versión y fiabilidad (alfa de Cronbach = .84; confiabilidad compuesta = .83) y una estructura factorial similar a la versión original. El miedo de la COVID-19 se asoció positivamente ($.23 < r < .31$) con depresión, ansiedad y estrés. Los resultados del análisis transversal de invariancia multigrupo respaldaron la invariancia escalar total de la FCV-19S-P y su invariancia residual parcial, lo que sugiere que esta medida puede usarse para llegar a conclusiones válidas con respecto a las comparaciones de género en muestras de adultos portugueses.

Palabras clave: Escala de Miedo a la COVID-19. Validez. Fiabilidad. Invariancia de medida. Portugal.

Abstract: The novelty and uncertainty of the pandemic nourished a generalized fear of the COVID-19, which seems to have exacerbated the pandemic's negative impact. It is thus relevant to monitor fear of COVID-19 and its association with individuals' mental health, well-being, and behaviors. Valid and reliable measures of fear of COVID-19 are necessary for that purpose. This study aimed at assessing the psychometric properties of a European Portuguese version of the Fear of COVID-19 scale (FCV-19S-P). A secondary aim was to assess FCV-19S-P's multigroup measurement invariance (female vs. male). A sample of 572 Portuguese adults (72 % female) completed the FCV-19S-P and measures of depression, anxiety, and stress. The study results supported this version validity and reliability (Cronbach's alpha = .84; Composite Reliability = .83), and a factorial structure similar to the original version. Fear of COVID-19 was positively associated ($.23 < r < .31$) with depression, anxiety, and stress. Results of the multigroup invariance analysis supported the FCV-19S-P total scalar invariance and its partial residual invariance, suggesting that this measure may be used to reach valid conclusions in respect to gender comparisons in samples of Portuguese adults in regard to group observed composite means.

Keywords: Fear of COVID-19 Scale. Validity. Reliability. Measurement invariance. Portugal.

Introduction

COVID-19 has spread worldwide and it has been one of the greatest challenges for societies and countries universally (Bitan et al., 2020). SARS-COV-2 virus has a high infection rate, and COVID-19 contributes to the overload of health care systems and governments (Ferrara & Albano, 2020; Worldometer, 2020). In addition, the COVID-19 pandemic was also associated with a relatively low level of knowledge about the disease, especially in its beginning, and with high uncertainty of its development over time and of its negative consequences on different fronts (e.g., a threat to health, job loss; Bambra et al., 2020; Carlucci et al., 2020; Nicola et al., 2020). Consequently, a sense of fear of the COVID-19 grew among the general population (Fofana et al., 2020; Lee et al., 2020). This sense of fear is associated with the virus and the disease in itself (e.g., fear of being near people infected with COVID-19, of getting sick, and of dying), and of not having

access to adequate healthcare in case of need due to the overload of healthcare services (Cawcutt et al., 2020; Ferrara & Albano, 2020; Lin, 2020; Okereke et al., 2021).

These concerns resulted in fear of the COVID-19 and are associated with a deterioration of pre-existing mental health conditions (Colizzi et al., 2020). Even in the absence of such conditions, negative effects of the COVID-19 pandemic and of social isolation on individuals' mental health are to be expected (Bao et al., 2020; Ornell et al., 2020; Usher, Durkin, et al., 2020; Usher, Jackson, et al., 2020). Although fear may predict behavior changes in the short-term in the direction desired by governmental and health agencies – e.g., physical distancing (Harper et al., 2020) – previous literature has shown that long-term fear has limitative effects and hampers behavioral changes (Adolphs, 2013; Leventhal, 1970; Ropeik, 2004). In the long run, such fear might decrease individuals' compliance with health behaviors and with (inter)national health agencies' recommendations (Centers for Disease Prevention and Control., 2019; Fofana et al., 2020). Additionally, fear seems also to be associated with the exacerbation of physical and psychological issues. For example, fear weakens the immune system, and is associated with cardiovascular diseases, fatigue, long-term memory impairment, mood swings, among others (Robinson et al., 2013;

* Correspondence address [Dirección para correspondencia]:

Alexandra Ferreira-Valente. Research Center for Human Development, Faculty of Education and Psychology, Universidade Católica Portuguesa, Rua de Diogo Botelho, Porto, 1327 4169-005 (Portugal).

E-mail: mafvalente@gmail.com

(Article received: 20-01-2022; revised: 22-07-2022; accepted: 22-07-2022)

Roest et al., 2017; Rosenberg, 2017; Segerstrom & Miller, 2004). In recent studies about the effects of COVID-19-related fear, fear of COVID-19 was found to be associated with an increase in reported depression, anxiety, stress, social isolation, loneliness, sleep problems, and cognitive functioning impairment (Baker et al., 2016; Bitan et al., 2020; Bjursell, 2020; Fofana et al., 2020; Koçak et al., 2021; Pyszczyński et al., 2020). It is therefore paramount to monitor fear of COVID-19 over time, in order to study its association with health-related outcomes, health behaviors, and behavioral changes.

Ahorsu and colleagues (2020) have recently developed a measure to assess individuals' fear of COVID-19: the Fear of COVID-19 Scale (FCV-19S). First, a panel of experts identified and analyzed the content of the items of 28 different measures of fear associated with different stimuli in different study populations. The panel of experts retained 17 items which were sent to a second panel of experts for a second round of content analysis. The 10 items retained after this second round of analysis were piloted on a sample of 46 individuals (Ahorsu et al., 2020), and the three items with low inter-item correlations ($r < .06$) were eliminated from the final seven-item scale. Exploratory factor analysis suggested a unidimensional reliable (Cronbach's alpha of .82) seven-item scale. The validity of the FCV-19S was further inspected through the analysis of the correlations (positive and moderate) of the total score of this measure with measures of anxiety, depression, and perceived vulnerability to the disease (Ahorsu et al., 2020).

The FCV-19S has been translated, adapted, and validated in 32 countries (see Appendix 1), sometimes more than once, originating a total of 38 different translated versions of this measure (Ahorsu et al., 2020; Alyami et al., 2020; Barbosa-Camacho et al., 2021; Barrios et al., 2020; Bitan et al., 2020; Cavalheiro & Sticca, 2020; Caycho-Rodríguez et al., 2020; Chang et al., 2020; Chi et al., 2021; El-Bardan & Lathabhavan, 2021; Elemo et al., 2020; Faro et al., 2020; Giordani et al., 2020, 2021; Huarcaya-Victoria et al., 2020; Iversen et al., 2021; Lathabhavan, 2021; Magano et al., 2021; Mahmood et al., 2020; Mailliez et al., 2021; Martínez-Lorca et al., 2020; Masuyama et al., 2020; Medeiros et al., 2021; Midorikawa et al., 2021; Nguyen et al., 2020; Pang et al., 2020; Perz et al., 2020; Pilch et al., 2021; Piqueras et al., 2021; Reznik et al., 2020; Sakib et al., 2020; Satici et al., 2020; Soraci et al., 2020; Tsipropoulou et al., 2020; Winter et al., 2020). The FCV-19S was found to sustain good reliability across several languages and cultures, but debate remains relative to this measure's factorial structure. The validation studies of the American (Perz et al., 2020), Bangladeshi (Sakib et al., 2020), Brazilian (Cavalheiro & Sticca, 2020; Faro et al., 2020; Giordani et al., 2020; Medeiros et al., 2021), Dominican (Piqueras et al., 2021), Egyptian (El-Bardan & Lathabhavan, 2021), Ethiopian (Elemo et al., 2020), French (Mailliez et al., 2021), Greek (Tsipropoulou et al., 2020), Italian (Soraci et al., 2020), Malaya (Pang et al., 2020), Mozambican (Giordani et al., 2021), New Zealand (Winter et al., 2020), Polish (Pilch

et al., 2021), Saudi Arabian (Alyami et al., 2020), Spanish (Martínez-Lorca et al., 2020; Piqueras et al., 2021), Taiwanese (Chang et al., 2020), Turkish (Satici et al., 2020), and Vietnamese (Nguyen et al., 2020) versions of the FCV-19S yielded a one-factor solution, similar to the one found for the original version of this measure. For the Argentinian (Caycho-Rodríguez et al., 2020), Belarusian (Reznik et al., 2020), Chinese (Chi et al., 2021), Israeli (Bitan et al., 2020), Norwegian (Iversen et al., 2021), Paraguayan (Barrios et al., 2020), Peruvian (Huarcaya-Victoria et al., 2020), and Russian versions (Reznik et al., 2020), however, a two-factor structure was found. Some validation studies of the FCV-19S within the same country originated mixed results regarding the factorial structure of the scale. For instance, Lathabhavan (2021) evaluated the one- and two-factor structures of the FCV-19S in a sample of 992 participants from India. This author concluded that both the one- and two-factor models showed good psychometric properties. The same scenario happened in a validation study with 1216 Mexican respondents, where both one- and two-factor models had adequate psychometric properties (Barbosa-Camacho et al., 2021). In Japan, three independent studies evaluated the psychometric properties of a Japanese version of the FCV-19S (Masuyama et al., 2020; Midorikawa et al., 2021; Wakashima et al., 2020). Wakashima and colleagues (2020) were able to replicate the one-factor solution of the original scale, while Masuyama and colleagues (2020) and Midorikawa and colleagues (2021) found a two-factor solution to be the better fit for their data. Finally, in Pakistan, one study found a one-factor structure for the FCV-19S (Mahmood et al., 2020), while another study found a three-factor structure (Basit et al., 2021).

A European Portuguese (i.e., Portuguese as spoken in Portugal) version of the FCV-19S has been studied by Magano and colleagues (2021). Though this version has shown good internal consistency in a convenience and snowball sample of adults from the Portuguese general population recruited between September and November of 2020 through the author's study social network website, the authors were not able to replicate the one-factor structure found by Ahorsu and colleagues (2020). The authors performed exploratory factorial analysis ($n = 561$), followed by confirmatory factorial analysis ($n = 561$) testing the measurement model found in the exploratory factor analysis. Their findings yielded and confirmed a two-factor solution for this version of the FCV-19S.

While translated and validated measures in different languages and cultures are necessary to enable the comparison of data from several subjects from different backgrounds – facilitating cross-cultural research and cross-country comparisons – valid cross-country/cultural and cross-group comparisons require at least a minimum form of measurement invariance, i.e. the extent to which the psychometric properties of a given questionnaire generalize across groups (Brown, 2006; Gregorich, 2006; Vandenberg & Lance, 2000). The starting point for measurement invariance is configural invariance, i.e. the factor structure of a measure is the same

in the different groups (Gregorich, 2006; Vandenberg & Lance, 2000). Although a measure's factor structure alone in different groups does not provide evidence of configural invariance, equal factor structures are a prerequisite for configural invariance and for testing configural invariance. Additionally, loadings' pattern must be similar (Byrne, 2016). Attempting to replicate the factorial structure of the original version of the FCV-19S is a first step in verifying this prerequisite to further test stronger forms of invariance (i.e., metric, scalar, and residual/uniqueness invariance), which are necessary for cross-country comparisons. The factor structure variability observed in the different translated versions of the FCV-19S may be associated with the self-report nature of the measure, its inherent subjectivity, the influence of eventual minor psycholexical (as well as of contextual and cultural) inequivalences between the translated versions regarding items' interpretation (Beaton et al., 2000; Kim & Zabelina, 2015; Solano-Flores & Nelson-Barber, 2001; Suzuki & Ponterotto, 2008).

The purpose of this study is to assess the psychometric properties of a European Portuguese version of the FCV-19S (FCV-19S-P; see Appendix 2) in a sample of Portuguese adults, testing if a one-factor solution – similar to the factorial structure found for the original version of the FCV-19S – would be supported in Portuguese adults from the general population. A secondary aim of the study was to test the multigroup (female vs. male participants) measurement invariance of the FCV-19S-P. We anticipate that: (a) confirmatory factor analysis will support a one-factor solution; (b) internal consistency and composite reliability will range between .73 and .93; (c) the FCV-19S total score will be positively moderately associated with depression, anxiety, and stress; and that (d) female participants will report significant-

ly higher levels of self-reported fear of COVID-19 than their male counterparts. We also anticipate that the FCV-19S-P will show multigroup measurement invariance, allowing for meaningful comparisons between Portuguese female vs. male adult individuals.

Methods

Participants

Participants were adults from the general population living in Portugal between November 2020 and January 2021. Inclusion criteria were: (a) being at least 18 years old; (b) living in Portugal at the time of assessment; (c) being able to read and understand Portuguese; and (d) being willing to participate.

The minimum sample size required to perform the confirmatory factor analysis using structural equation modeling was determined using a priori power calculation, as described below (Cohen, 1988; Soper, 2018). This analysis resulted in a minimum sample size recommended of 400 participants. A total of 580 individuals agreed to participate. Participants with missing values ($n = 6.1\%$) were excluded from the study sample, resulting in a sample of 574 participants. Another two participants, who were found to be multivariate outliers for the FCV-19S-P items, were also excluded from the study sample, resulting in a final sample of 572 individuals (response rate of 99%). Table 1 summarizes the demographic characteristics of the study sample. Most participants were women ($n = 410, 72\%$), aged between 18 and 88 years old ($M = 33.46; SD = 13.44$). Most participants had a college degree ($n = 456, 80\%$) and were single ($n = 396, 69\%$).

Table 1

Characteristics of the Study Sample (n = 572).

		<i>n</i>	<i>%</i>
Gender	Female	410	71.7
	Male	162	28.3
Educational level	9 th grade or lower	5	.9
	10 th to 12 th grade	107	18.7
	(At least some) College	456	79.7
Marital status	Single	396	69.2
	Married/Marital relationship	137	24
	Divorced	34	5.9
	Widow	5	0.9

Measures

Participants provided basic sociodemographic information (e.g., age, self-identified gender, marital status, level of education). Additionally, participants completed the Portuguese version of the Depression, Anxiety, and Stress Scale – Short Form (DASS-21; Lovibond & Lovibond, 1995; Pais-Ribeiro et al., 2004) and a Portuguese version of the Fear of COVID-19 Scale (FCV-19S; Ahorsu et al., 2020).

Fear of COVID-19 Scale.– The FCV-19S (Ahorsu et al., 2020) is a self-report measure aimed at assessing the fear of COVID-19. It is a seven-item scale and participants are requested to respond on a 5-point Likert scale (1 = “strongly disagree”; 5 = “strongly agree”). The total score ranges between 7 and 35, with a higher sum score indicating higher fear of COVID-19.

Depression, Anxiety, and Stress Scale – Short Form.– The DASS-21 (Lovibond & Lovibond, 1995) is a three-dimensional self-report measure of anxiety, depression, and

stress. It is composed by 21 items grouped in three subscales of seven items each: (a) Depression; (b) Anxiety; and (c) Stress. Participants are asked to respond using a four-item type of Likert scale ranging from 1 (“did not apply to me at all”) to 4 (“applied to me very much”). Both the original (Lovibond & Lovibond, 1995) and the Portuguese version (Pais-Ribeiro et al., 2004) of the DASS-21 showed acceptable to excellent internal consistency ($.84 < a < .91$ and $.74 < a < .85$, respectively). This measure has shown good to excellent internal consistency ($.86 < a < .91$) in the study’s sample.

Procedure

The study was reviewed and approved by Ispa’s Ethical Committee for Research (reference I/033/04/2020). Permission was obtained from the authors of the original version of the FCV-19S to translate and study the psychometric properties of the measure in a sample of Portuguese adults. The initial phase of the study consisted of translating and back-translating the instructions and items of the FCV-19S. A consensus version was achieved via expert discussion who analyzed the content of the items to certify that the consensus version measured the same construct as the original version. The study data were collected via an online survey platform between November 2020 and January 2021. Prospective participants were invited to participate through multiple channels: (a) website and newsletter of the Order of Portuguese Psychologists – OPP; (b) a circular email sent to organizations (e.g., educational and health institutions and their professionals) and individuals, who were asked to disclose the study to their collaborators, and professional and personal contacts; and (c) social media. Potentially interested individuals were redirected to the study website and online survey, including an informed consent form with a full description of the study aims and procedures. Participants were assured that participation was anonymous and voluntary, and that they could drop out at any time. Upon providing informed consent, participants were invited to complete the study measures.

Data Analysis

The analysis of the psychometric properties of the FCV-19S was based on classical test theory (CTT) analysis, which included confirmatory factor analysis (CFA), average variance extracted, internal consistency, composite reliability, corrected item-total correlation, and assessment of this measure’s correlation pattern with criterion measures (DeVellis, 2006). We determined the minimum sample size required to perform the confirmatory factor analysis using an online a priori power calculator (Soper, 2018), considering an effect size (d de Cohen) of 0.30, a significance level of .05, and a power of 0.80 (Cohen, 1988).

We first computed frequencies, means, standard deviations of the study variables with descriptive purposes. Second, skewness (S_k) and kurtosis (K_u) analysis was performed

to evaluate items’ sensitivity, with absolute values lower than three and 10, respectively, standing for absence of severe violation of normality assumption and item sensitivity (Byrne, 2016; Kline, 2000). Third, we performed confirmatory factor analysis (CFA) to test the one-factor solution of the FCV-19S-P. Model fit was assessed considering χ^2 and its subsequent ratio with degrees of freedom (χ^2/df), comparative fit index (CFI), root mean square error of approximation (RMSEA), and standardized root mean square residual (SRMR). Model fit was considered acceptable or good, respectively, in the event that χ^2/df was lower than 5 or 2 (Wheaton, 1987), CFI was higher than .80 and .90 (Bentler, 1990), RMSEA was lower than .08 or .05 (Schermelel-Engel et al., 2003), and SRMR was lower than .10 or .05 (Hoyle, 1995; Schermelleh-Engel et al., 2003). According to Barbara Byrne’s recommendations (Byrne, 2016), the model’s adjustment was performed step-by-step, through the analysis of correlation among errors, analyzing Modification Indexes (MI) higher than 4 ($p < .001$; Arbuckle, 2008). The χ^2 difference test [$\Delta\chi^2(df)$, p -value], expected cross-validation index (MECVI) and Akaike information criterion (AIC) were computed to compare fit of the initial and final models after adjustments. Statistically significant χ^2 statistic and lower MECVI and AIC reflect better fit (Bentler, 1990; Schermelleh-Engel et al., 2003). Forth, the convergency of the scale was assessed through the analysis of the average variance extracted (AVE), i.e. the analysis of the amount of variance captured by the latent factor or construct in comparison to the amount of variance that may be caused by measurement error (Fornell & Larcker, 1981). An AVE higher than .50 indicates that the variance explained by the factor or construct is higher than the variance from measurement error (Fornell & Larcker, 1981; Hair et al., 2009), and convergence is deemed adequate. Fifth, we assessed the reliability of the FCV-19S-P by computing its’ internal consistency (Cronbach’s alpha), McDonald’s coefficient omega, and composite reliability (CR; Fornell & Larcker, 1981; Hair et al., 2009; McDonald, 1999), with coefficients higher than .70 and .80 indicating acceptable to good reliability, respectively. Inter-item correlations and corrected item-total correlations were also computed, with acceptable correlations ranging between .30 and .70 (Ferketich, 1991). We then analyzed the pattern of associations between the FCV-19S-P and criterion variables (i.e., depression, anxiety, and stress subscales of the DASS-21) by computing the respective Pearson correlation coefficients. Correlation coefficients were considered to be weak, moderate, and strong if absolute values ranged between .10 to .29, .30 to .49, or were higher than .50, respectively (Cohen, 1988). An independent sample t test was performed to assess potentially existing statistically significant gender differences in self-reported fear of COVID-19. Finally, we assessed FCV-19S-P multigroup (female vs. male participants) measurement invariance. To do so, CFA with maximum likelihood estimation was used. In step one, CFA was used to identify the baseline structure of the FCV-19S-P that best fitted participants of both genders (Byrne, 2016). In step

two, each group best fitting models were incorporated into two multigroup CFA models to test between-group measurement invariance (Byrne, 2004, 2016; Gregorich, 2006; Vandenberg and Lance, 2000). Four nested models were consecutively tested (Byrne, 2004, 2016; Gregorich, 2006; Schmitt and Kuljanin, 2008) to assess: (a) configural invariance (no equality constraints were imposed on parameters; it implies that the factor structure of the FCV-19S-P is the same in the two groups); (b) metric invariance (factor loadings were constrained; common factors have similar meaning across groups); (c) scalar invariance (factor loadings and item intercepts were constrained; the latent factors' scores are similar in both groups); and (d) residual/uniqueness invariance (factor loadings, item intercepts and item residual variances were constrained; the item intercepts and residual variances are similar across the two groups). Forms of measurement invariance that are subsequent to configural invariance are confirmed by comparing the fit of the subsequent nested model with the one of the preceding nested models, in which similar fit suggests a subsequent form of measurement invariance. To test the relative fit of nested models we computed the difference between alternative fit indices for nested models: χ^2/df , *CFI*, *RMSEA*, and *SRMR*, with $p < .05$

for $\Delta \chi^2/df$, a decrease in *CFI* of $\geq .01$ complemented by an increase in *RMSEA* of $\geq .015$ among nested invariance models indicating a significant worsening of fit (D. C. R. Chen et al., 2012; F. F. Chen, 2007; Meade et al., 2008). For the *SRMR* fit indices, an increase of $\geq .03$ in the configural model and an increase of $\geq .01$ in the following models would indicate a significant worsening of fit (F. F. Chen, 2007). Statistical analyses were executed using software IBM SPSS Statistics (v. 26) and AMOS statistical package (v. 26). Alpha was set at .05 for all statistical analyses.

Results

Descriptive Information

Descriptive statistics are presented in Table 2. As can be seen, the fear of COVID-19 ranged between 7 and 35 ($M = 13.4$, $SD = 5.1$). Answers to individual items ranged between one and five, with average scores ranging between 1.2 ($SD = 0.6$ for item 3 and 2.6 ($SD = 1.2$) for item 2. The entire five-point Likert scale was used by participants for all items, and the distribution of all items had acceptable skewness ($0.267 < Sk < 2.99$) and kurtosis ($-0.568 < Ku < 9.96$) values.

Table 2

Descriptive Statistics for FCV-19S-P and for FCV-19S-P Individual Items.

	<i>M</i>	<i>SD</i>	<i>Me</i>	<i>Min</i>	<i>Max</i>	<i>Sk</i>	<i>Ku</i>
FCV-19S-P Total	13.43	5.14	12	7	35	0.925	0.582
Item 1	2.20	1.10	2	1	5	0.566	-0.568
Item 2	2.59	1.24	2	1	5	0.267	-1.085
Item 3	1.23	0.60	1	1	5	2.99	9.96
Item 4	2.25	1.29	2	1	5	0.730	-0.641
Item 5	2.44	1.23	2	1	5	0.471	-0.811
Item 6	1.27	0.62	1	1	5	2.703	8.22
Item 7	1.44	0.87	1	1	5	2.204	4.452

Confirmatory Factor Analysis and Convergence of the FCV-19S-P

The model fit indexes for CFA are presented in Table 3, while Figure 1 shows the standardized factorial weights and individual items' reliability for the initial and final models. Only two out of four combined fit indices for the CFA supported the one-factor solution with acceptable fit ($\chi^2/df = 23.38$; *CFI* = .81; *RMSEA* = .20; *SRMR* = .09). Thus, the quality of fitness of the initial model was limited. The inspection of the FCV-19S-P items suggests that some of them might have similar content. This may be a possible explana-


tion for the limited fitness of the unidimensional solution on this sample. For instance, items three ("My hands become clammy when I think about coronavirus-19"), six ("I cannot sleep because I'm worrying about getting coronavirus-19"), and seven ("My heart races or palpitates when I think about getting coronavirus-19"), all seem to assess psychosomatic manifestations of fear. The analysis of the modification indices seems to suggest that the error terms of these three items should be correlated. These error terms were correlated sequentially resulting in a modified one-factor model that maintained the seven items of the original FCV-19S-P.

Table 3

Model Fit Indices for Confirmatory Factor Analysis.

	Number of parameters estimated	$\chi^2(df), p$ -value	χ^2/df	<i>CFI</i>	<i>RMSEA</i>	<i>SRMR</i>	<i>MECVI</i>	<i>AIC</i>	$\Delta \chi^2(df), p$ -value
Initial model	Weights: 14 Variances: 8 Covariances: 0	327.26 (14), $p < .001$	23.38	.814	.198	.087	.645	369.26	292.13 (3), $p < .001$
Final model	Weights: 14 Variances: 8 Covariances: 3	35.13 (11), $p < .001$	3.19	.986	.061	.023	.145	82.13	

Figure 1
Factor Structure of the Initial and Final Models, Respectively, of the FCV-19S-P.



The four combined fit indices suggested acceptable to good fit of this modified model ($\chi^2/df = 3.19$; $CFI = .99$; $RMSEA = .06$; $SRMR = .02$). This model showed a higher goodness of fit than the initial model [$\Delta\chi^2(3) = 292.13, p < .001$; $MECVI: .15$ vs. $.65$; $AIC: 82.13$ vs. 369.26). Six of the items (86%) showed loadings equal or higher than $.50$, with item six presenting a loading of $.48$, indicating that the corresponding latent dimensions explained, at least, 25% of the result of all items except for item six. FCV-19S-P's AVE

was estimated in $.42$, indicating a less than optimal convergence of this measure.

Reliability: Internal Consistency and Composite Reliability

Table 4 summarizes the results for the measure's reliability (see Appendix 3 for inter item-correlations). As can be seen, the overall scale showed a good internal consistency ($\alpha = .84$), coefficient omega ($\omega = .85$), and composite reliability ($CR = .84$), with internal consistency coefficients if single items are deleted being comparable to the overall Cronbach's alpha, indicating that no item detracts from the reliability of the FCV-19S-P (see Table 4). Inter-item correlation coefficients ranged from $.30$ to $.62$ and maybe considered significant and not redundant.

Table 4
Average Variance Extracted and Reliability Analysis.

	FCV-19S-P
AVE	.42
CR	.83
McDonald's omega	.85
Cronbach's alpha	.84
Cronbach's alpha if item deleted	
Item 1	.81
Item 2	.80
Item 3	.83
Item 4	.82
Item 5	.80
Item 6	.83
Item 7	.82

Association of FCV-19S-P with Gender, Depression, Anxiety, and Stress

Fear of COVID-19 presented weak to moderate positive and statistically significant associations with scores of depression ($r = .23, p < .001$), anxiety ($r = .25, p < .001$) and stress ($r = .31, p < .001$), as measured by the Portuguese version of DASS-21. There was a statistically significant gender difference on average fear of COVID-19, $t(570) = 6.056, p < .001, d = 4.98$, with women ($M = 14.22, SD = 5.17$) reporting greater fear of COVID-19 than their male counterparts ($M = 11.42, SD = 4.47$).

Multigroup Measurement Invariance of the FCV-19S-P

The model fit of the four nested models and the fit indices for the multigroup measurement invariance models (female vs. male participants) are presented in Table 5. The configural model showed acceptable ($\chi^2/df = 2.44$) to good fit ($CFI = .98, RMSEA = .05$, and $SRMR = .03$) in all fit indices. It was, therefore, considered suitable to be used as the basis for testing the constrained models.

Table 5
Models Fit Indices for the Multigroup Measurement Invariance Models: Female vs. Male

	$\chi^2(df), p\text{-value}$	χ^2/df	CFI	RMSEA	SRMR	
Configural invariance model	53.57(22), $p < .001$	2.44	.98	.05	.03	
Metric invariance model	57.34(28), $p = .001$	2.05	.98	.04	.03	
Scalar invariance model	70.56(34), $p < .001$	2.08	.98	.04	.03	
Residual invariance model	105.87(41), $p < .001$	2.58	.96	.05	.03	
Nested model comparisons						
	Δdf	$\Delta\chi^2$	$p\text{-value}$	ΔCFI	$\Delta RMSEA$	$\Delta SRMR$
<i>Assuming configural invariance model to be correct</i>						
Metric invariance model	6	3.77	.778	0	-.01	0
Scalar invariance model	12	16.99	.149	0	-.01	0
Residual invariance model	19	52.30	.001	.02	0	0
<i>Assuming metric invariance model to be correct</i>						
Scalar invariance model	6	13.22	.04	0	0	0
Residual invariance model	13	48.53	.001	.02	-.01	0
<i>Assuming scalar invariance model to be correct</i>						
Residual invariance model	7	35.31	.001	.02	-.01	0

In comparison to the unconstrained model, the metric invariance model presented no significant worsening of the model fit in all four fit indices [$\Delta\chi^2(df) = 3.77$, $\Delta df = 6$, $p = .778$; $\Delta CFI = 0$; $\Delta RMSEA = -.01$; $\Delta SRMR = 0$], which suggests that the magnitudes of factor loadings were similar across groups. Similarly, the scalar invariance model revealed no significant deterioration of model fit in three out of four fit indices [$\Delta\chi^2(df) = 13.22$, $\Delta df = 6$, $p = .04$; $\Delta CFI = 0$; $\Delta RMSEA = 0$; $\Delta SRMR = 0$], suggesting that item intercepts are similar across groups. The residual invariance model, however, resulted in a slight, but significant, worsening of model fit in two out of four fit indices comparing to the metric and to the scalar invariance models [$\Delta\chi^2(df) = 35.31$, $\Delta df = 7$, $p < .001$; $\Delta CFI = .02$; $\Delta RMSEA = -.01$; $\Delta SRMR = 0$], suggesting that item residual variances were not similar across groups.

The partial residual invariance was tested (i.e. to find the source of non-invariance in the residual invariance model). We implemented a backward method involving sequentially dropping the constraints that contributed greater chi-square valued to the model. As a result, we sequentially freed items 3, 6, and 7, so that the partial residual invariance model had no significant worsening in model fit as compared with the scalar invariance model for all fit indices [$\Delta\chi^2(df) = 5.64$, $\Delta df = 4$, $p = .228$; $\Delta CFI = 0$; $\Delta RMSEA = 0$; $\Delta SRMR = 0$]. This resulted in a partial residual invariance model with good fit ($\chi^2/df = 2.01$; $CFI = .98$; $RMSEA = .04$; $SRMR = .03$).

Discussion

The COVID-19 pandemic had a severe negative psychological and socioeconomic impact on societies and individuals (Bitan et al., 2020). Its rapid spread, the initial lack of knowledge in regards to how it is transmitted and its short/long-term effects, and the uncertainty relative to how would the pandemic unfold in the medium/long-term, nurtured a generalized fear of the COVID-19 (Bambra et al., 2020; Carlucci et al., 2020; Nicola et al., 2020). Such fear

seems to be associated with an aggravation of the pandemic's negative impact upon individuals' psychological and physical health, social support, and health behaviors (Bao et al., 2020; Ornell et al., 2020; Usher, Durkin, et al., 2020; Usher, Jackson, et al., 2020). Given the potentially noxious effect of fear of COVID-19, Ahorsu and colleagues (2020) have developed the FCV-19S. Their purpose was to provide a useful tool to monitor the fear of COVID-19 and its consequences. The FCV-19S has now been translated into various languages and studied in samples of adults in 32 different countries.

The current study aimed at assessing the psychometric properties of a European Portuguese version of the FCV-19S, testing if a one-factor solution, similar to the one found for the original version of the FCV-19S, would be yielded in Portuguese adults from the general population. A secondary aim of the study was to test the multigroup (female vs. male participants) measurement invariance of the FCV-19S-P. The study findings supported most of our results' expectations. First, as predicted, confirmatory factor analysis supported the unidimensional factor structure of seven items of the FCV-19S-P, similar to the factorial structure found for the original version of FCV-19S (Ahorsu et al., 2020). Second, as expected, the FCV-19S-P showed a good reliability on the range of the coefficients found in previous research studies examining the psychometric properties of other versions of the FCV-19S (Ahorsu et al., 2020; Alyami et al., 2020; Barbosa-Camacho et al., 2021; Barrios et al., 2020; Bitan et al., 2020; Cavalheiro & Sticca, 2020; Caycho-Rodríguez et al., 2020; Chang et al., 2020; Chi et al., 2021; El-Bardan & Lathabhavan, 2021; Elemo et al., 2020; Faro et al., 2020; Giordani et al., 2020, 2021; Huarcaya-Victoria et al., 2020; Iversen et al., 2021; Lathabhavan, 2021; Magano et al., 2021; Mahmood et al., 2020; Mailliez et al., 2021; Martínez-Lorca et al., 2020; Masuyama et al., 2020; Medeiros et al., 2021; Midorikawa et al., 2021; Nguyen et al., 2020; Pang et al., 2020; Perz et al., 2020; Pilch et al., 2021; Piqueras et al., 2021; Reznik et al., 2020; Sakib et al., 2020; Satıcı et al., 2020;

Soraci et al., 2020; Tsipropoulou et al., 2020; Winter et al., 2020). Third, fear of COVID-19, as measured by the FCV-19S-P, was positively correlated with measures of depression, anxiety and stress (Ahorsu et al., 2020; Bitan et al., 2020; Satici et al., 2020), as anticipated. Fourth, congruent with our expectations and with the findings from previous studies examining the psychometric properties of the FCV-19S, women reported greater fear of COVID-19 than men (Barbosa-Camacho et al., 2021; Basit et al., 2021; Bitan et al., 2020; Cavalheiro & Sticca, 2020; Chi et al., 2021; Giordani et al., 2020, 2021; Huarcaya-Victoria et al., 2020; Iversen et al., 2021; Magano et al., 2021; Mahmood et al., 2020; Midorikawa et al., 2021; Nguyen et al., 2020; Pilch et al., 2021; Piqueras et al., 2021; Reznik et al., 2020; Sakib et al., 2020; Tsipropoulou et al., 2020). Finally, the FCV-19S-P multigroup (female vs. male participants) scalar (and residual) invariance was (at least partially) supported. However, our findings provide only partial support for the convergency of the FCV-19S-P. In fact, the average variance extracted (*AVE*) lower than the proposed cut-off point seems to suggest that the variance associated with measurement error may be slightly higher than the one captured by the construct (Fornell & Larcker, 1981). Nonetheless, according to Fornell and Larcker (1981), given the measure's good reliability, as assessed through Cronbach's alpha, McDonald's omega, and composite reliability coefficients, convergency of the construct may still be considered adequate. Taken together, these findings seem to suggest that the FCV-19S-P is a reliable and valid measure of fear of COVID-19, that may be used in studies focusing on Portuguese adults from the general population, and to draw valid conclusions on between-gender comparisons.

Our results are consistent with those found in the validation study of the original version of the FCV-19S (Ahorsu et al., 2020), as well as with those found in other 25 (out of 38) translated versions of this measure (Alyami et al., 2020; Barbosa-Camacho et al., 2021; Cavalheiro & Sticca, 2020; Chang et al., 2020; El-Bardan & Lathabhavan, 2021; Elemo et al., 2020; Faro et al., 2020; Giordani et al., 2020, 2021; Lathabhavan, 2021; Mahmood et al., 2020; Mailliez et al., 2021; Martínez-Lorca et al., 2020; Medeiros et al., 2021; Nguyen et al., 2020; Pang et al., 2020; Perz et al., 2020; Pilch et al., 2021; Piqueras et al., 2021; Sakib et al., 2020; Satici et al., 2020; Soraci et al., 2020; Tsipropoulou et al., 2020; Wakashima et al., 2020; Winter et al., 2020). They differ, however, from the results found in 12 other versions (Barbosa-Camacho et al., 2021; Barrios et al., 2020; Basit et al., 2021; Bitan et al., 2020; Caycho-Rodríguez et al., 2020; Chi et al., 2021; Huarcaya-Victoria et al., 2020; Iversen et al., 2021; Lathabhavan, 2021; Masuyama et al., 2020; Midorikawa et al., 2021; Reznik et al., 2020) in regards to the factorial structure of the measure. Our results are also inconsistent with those found for the European Portuguese version of the FCV-19S proposed by Magano and colleagues (2021), with respect to the factorial structure of the two versions and to their multigroup measurement invariance. One possible explanation for the facto-

rial differences between our version and Magano and colleagues' (2021) version might be the existence of slight translation nuances which may originate a slightly different interpretation of the items by adults from the general population living in Portugal. The confirmation of the one-factor solution, as found for the FCV-19S-P version we developed and studied here – unlike the version of Magano and colleagues (2021) – is a first step for testing configural invariance relative to the original version and other 25 translated versions, as a single-sample confirmatory factor analysis replicating a previously supported measurement model is a precondition for further testing configural invariance (Byrne, 2016). This minimum form of measurement invariance, yet not strictly sufficient, constitutes a pre-requisite for defensible comparisons relative to results obtained with samples of participants from different cultures, languages, and countries (Gregorich, 2006; Vandenberg & Lance, 2000). Further cross-cultural research comparing different versions of the FCV-19S and examining measurement invariance is necessary to confirm that valid conclusions on cross-cultural/country comparisons may be drawn.

Unlike Magano and colleagues' (2021) version, the FCV-19S-P showed full configural, metric, and scalar invariance, suggesting that both genders perceive items and the underlying construct similarly. This suggests that gender comparisons relative to the estimated factor variances and covariances, and to the observed means of fear of COVID-19 are defensible. Though only partial residual invariance was supported by our findings, some authors argue that this type of invariance has limited practical value (Gregorich, 2006) and is thus not mandatory for multigroup comparisons. As a result, valid conclusions regarding meaningful gender comparisons relative to groups' estimated (co)variances and estimated/observed means could potentially be drawn. In our sample, women reported higher fear of COVID-19, a result that may be associated with a greater sensitivity/vulnerability to stress and anxiety of women (McLean & Anderson, 2009; Olatunji et al., 2005; Tolin & Foa, 2008).

Limitations

This study has a number of limitations that need to be taken into consideration when interpreting the results. Ours was a convenience sample, composed mostly of young highly educated adults, and women. In addition, we used a web-delivered administration method of the study measures, which may have resulted in a participant selection bias. Thus, the study sample is not representative of the Portuguese adult population (INE, 2020), which prevents the generalization of our findings to the study population. Future research with other samples is necessary to examine the reliability and generalizability of these findings. On the other hand, because of the cross-sectional design of this study, it was not possible neither to examine the test-retest reliability of the FCV-19S-P and its sensitivity to change, nor to assess the measure's longitudinal measurement invariance. Thus, future studies

with longitudinal design are necessary to verify the longitudinal measurement invariance and the stability over time of the FCV-19S-P. Finally, other measures assessing event-related fear were not administered. This would have helped us to further examine the FCV-19S-P convergency. Further research including alternative measures of fear is warranted.

Conclusions

Despite the study limitations, our findings support the reliability and validity of a one-factor European Portuguese version of the FCV-19S, as well as its multigroup/gender invariance. The FCV-19S-P is a useful tool that might be used to monitor the fear of COVID-19 among Portuguese adults and draw meaningful gender comparisons. Since fear can lead to various mental health disorders and be a potential predictor of vulnerable groups, the assessment of the level of fear among the general population by health entities and professionals can be a powerful ally in preventing mental health disorders and uncovering vulnerable groups (Holmes et al., 2020; Huarcaya-Victoria et al., 2020; Nitschke et al., 2020; Wang et al., 2021). This would allow for the develop-

ment of tailored interventions to reduce the fear of COVID-19 and help healthcare providers to identify those most vulnerable individuals at greater risk and that would benefit the most from rapid and timely psychological intervention. Future research should test longitudinal and cross-cultural multigroup measurement invariance to guarantee that the FCV-19S(-P) measures the same construct in the same way across time and populations.

Conflict of interest.- The authors declare no conflict of interest.

Financial support.- MJ is funded by national funds through FCT - Fundação para a Ciência e Tecnologia, I.P., under the FCT PhD grant (grant 2020.10251.BD). AFV is supported by national funds through FCT - Fundação para a Ciência e Tecnologia, I.P. and, when eligible, by COMPETE 2020 FEDER funds, under the FCT Post-doctoral grant (grant SFRH/BPD/121452/2016), as well as Scientific Employment Stimulus - Institutional Call (CEEC-INST/00070/2021; public notification Edital/0030/2022). The William James Center for Research, Ispa - University Institute is supported by FCT funding (reference UIDB/04810/2020). The Research Centre for Human Development is funded by FCT funding (reference UIDB/04872/2020).

References

- Adolphs, R. (2013). The biology of fear. *Current Biology*, 23(2), 79-93. <https://doi.org/10.1016/j.cub.2012.11.055>
- Ahorsu, D. K., Lin, C.-Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2020). The Fear of COVID-19 Scale: Development and initial validation. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00316-x>
- Alyami, M., Henning, M., Krägeloh, C. U., & Alyami, H. (2020). Psychometric evaluation of the Arabic version of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00316-x>
- Arbuckle, J. L. (2008). *Amos 17 users' guide*. Amos Development Corporation.
- Baker, T. A., Krok-Schoen, J. L., O'Connor, M. L., & Brooks, A. K. (2016). The influence of pain severity and interference on satisfaction with pain management among middle-aged and older adults. *Pain Research and Management*, 2016, Article 9561024. <https://doi.org/10.1155/2016/9561024>
- Bambra, C., Riordan, R., Ford, J., & Matthews, F. (2020). The COVID-19 pandemic and health inequalities. *Journal of Epidemiology and Community Health*, 74, 964-968. <https://doi.org/10.1136/jech-2020-214401>
- Bao, Y., Sun, Y., Meng, S., Shi, J., & Lu, L. (2020). 2019-nCoV epidemic: Address mental health care to empower society. *The Lancet*, 395(10224), e37-e38. [https://doi.org/10.1016/S0140-6736\(20\)30309-3](https://doi.org/10.1016/S0140-6736(20)30309-3)
- Barbosa-Camacho, F. J., García-Reyna, B., Cervantes-Cardona, G. A., Cervantes-Pérez, E., Chavarria-Avila, E., Pintor-Belmontes, K. J., Guzmán-Ramírez, B. G., Bernal-Hernández, A., Ibarrola-Peña, J. C., Fuentes-Orozco, C., González-Ojeda, A., & Cervantes-Guevara, G. (2021). Comparison of fear of COVID-19 in medical and nonmedical personnel in a public hospital in Mexico: A brief report. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-021-00600-4>
- Barrios, I., Ríos-González, C., O'Higgins, M., González, I., García, O., Díaz, N. R., Castaldelli-Maia, J. M., Ventriglio, A., & Torales, J. (2020). Psychometric properties of the Spanish version of the Fear of COVID-19 Scale (FCV-19S). *Research Square*. <https://doi.org/10.21203/rs.3.rs-33345/v1>
- Basit, K. A., Zafar, A. B., Fawwad, A., Waris, N., Shaheen, F., & Basit, A. (2021). Psychometric analysis for Fear of COVID-19 Scale (FCV-19S) and its association with depression in patients with diabetes: A cross sectional study from a tertiary care centre in Karachi, Pakistan. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 15(3), 733-737. <https://doi.org/10.1016/j.dsx.2021.03.008>
- Beaton, D. E., Bombardier, C., Guillemin, F., & Ferraz, M. B. (2000). Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine*, 25(24), 3186-3191. [10.1097/00007632-200012150-00014](https://doi.org/10.1097/00007632-200012150-00014)
- Bentler, P. M. (1990). Quantitative methods in psychology. *Psychological Bulletin*, 107(2), 238-246. <http://dx.doi.org/10.1037/0033-2909.107.2.238>
- Bitan, D. T., Grossman-Giron, A., Bloch, Y., Mayer, Y., Shiffman, N., & Mendlovic, S. (2020). Fear of COVID-19 scale: Psychometric characteristics, reliability and validity in the Israeli population. *Psychiatry Research*, 289, Article 113100. <https://doi.org/10.1016/j.psychres.2020.113100>
- Bjursell, C. (2020). The COVID-19 pandemic as disjuncture: Lifelong learning in a context of fear. *International Review of Education*, 66, 673-689. <https://doi.org/10.1007/s11159-020-09863-w>
- Brown, T. A. (2006). *Confirmatory factor analysis for applied research*. Guilford Press.
- Byrne, B. M. (2004). Testing for multigroup invariance using AMOS graphics: A road less traveled. *Structural Equation Modeling: A Multidisciplinary Journal*, 11(2), 272-300. https://doi.org/10.1207/s15328007sem1102_8
- Byrne, B. M. (2016). *Structural equation modeling with AMOS: Basic concepts, applications, and programming* (3rd ed.). Routledge.
- Carlucci, L., D'Ambrosio, I., & Balsamo, M. (2020). Demographic and attitudinal factors of adherence to quarantine guidelines during COVID-19: The Italian model. *Frontiers in Psychology*, 11, Article 559288. <https://doi.org/10.3389/fpsyg.2020.559288>

- Cavalheiro, F. R. S., & Sticca, M. G. (2020). Adaptation and validation of the Brazilian version of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00415-9>
- Cawcutt, K. A., Starlin, R., & Rupp, M. E. (2020). Fighting fear in healthcare workers during the COVID-19 pandemic. *Infection Control & Hospital Epidemiology*, *41*(10), 1192-1193. <https://doi.org/10.1017/ice.2020.315>
- Caycho-Rodríguez, T., Vilca, L. W., Cervigni, M., Gallegos, M., Martino, P., Portillo, N., Barés, I., Calandra, M., & Burgos Videla, C. (2020). Fear of COVID-19 Scale: Validity, reliability and factorial invariance in Argentina's general population. *Death Studies*, *46*(4), 1-10. <https://doi.org/10.1080/07481187.2020.1836071>
- Centers for Disease Prevention and Control (2019). *CERC: Psychology of a Crisis*. U.S. Department of Health and Human Services, Centers for Disease Prevention and Control (CDC).
- Chang, K.-C., Hou, W.-L., Pakpour, A. H., Lin, C.-Y., & Griffiths, M. D. (2020). Psychometric testing of three COVID-19-related scales among people with mental illness. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00361-6>
- Chen, D. C. R., Kirshenbaum, D. S., Yan, J., Kirshenbaum, E., & Aseltine, R. H. (2012). Characterizing changes in student empathy throughout medical school. *Medical Teacher*, *34*(4), 305-311. <https://doi.org/10.3109/0142159X.2012.644600>
- Chen, F. F. (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, *14*(3), 464-504. <https://doi.org/10.1080/10705510701301834>
- Chi, X., Chen, S., Chen, Y., Chen, D., Yu, Q., Guo, T., Cao, Q., Zheng, X., Huang, S., Hossain, M. M., Stubbs, B., Yeung, A., & Zou, L. (2021). Psychometric evaluation of the Fear of COVID-19 Scale among Chinese population. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00441-7>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Lawrence Erlbaum.
- Colizzi, M., Bortoletto, R., Silvestri, M., Mondini, F., Puttini, E., Cainelli, C., Gaudino, R., Ruggieri, M., & Zocante, L. (2020). Medically unexplained symptoms in the times of COVID-19 pandemic: A case-report. *Brain, Behavior, & Immunity - Health*, *5*, Article 100073. <https://doi.org/10.1016/j.bbih.2020.100073>
- DeVellis, R. F. (2006). Classical test theory. *Medical Care*, *44*(11 Suppl 3), S50-S59. [10.1097/01.mlr.0000245426.10853.30](https://doi.org/10.1097/01.mlr.0000245426.10853.30)
- El-Bardan, M. F., & Lathabhavan, R. (2021). Fear of COVID-19 Scale: Psychometric properties, reliability and validity in Egyptian population. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, *15*(4), Article 102153. <https://doi.org/10.1016/j.dsx.2021.05.026>
- Elemo, A. S., Satici, S. A., & Griffiths, M. D. (2020). The Fear of COVID-19 Scale: Psychometric properties of the Ethiopian Amharic version. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00448-0>
- Faro, A., Silva, L. dos S., Santos, D. N. dos, & Feitosa, A. L. B. (2020). Adaptação e validação da Escala de Medo da COVID-19. *SciELO Preprints*. <https://doi.org/10.1590/SciELOPreprints.898>
- Ferketich, S. (1991). Focus on psychometrics. Aspects of item analysis. *Research in Nursing & Health*, *14*(2), 165-168. <https://doi.org/10.1002/nur.4770140211>
- Ferrara, P., & Albano, L. (2020). COVID-19 and healthcare systems: What should we do next? *Public Health*, *185*, 1-2. <https://doi.org/10.1016/j.puhe.2020.05.014>
- Fofana, N. K., Latif, F., Sarfraz, S., Bilal, Bashir, M. F., & Komal, B. (2020). Fear and agony of the pandemic leading to stress and mental illness: An emerging crisis in the novel coronavirus (COVID-19) outbreak. *Psychiatry Research*, *291*, Article 113230. <https://doi.org/10.1016/j.psychres.2020.113230>
- Fornell, C., & Larcker, D. F. (1981). Evaluating structural equation models with unobservable variables and measurement error. *Journal of Marketing Research*, *18*(1), 39-50. <https://doi.org/10.2307/3151312>
- Giordani, R. C. F., Giolo, S. R., Muhl, C., Estavela, A. J., & Mabuie Gove, J. I. (2021). Validation of the FCV-19 Scale and assessment of fear of COVID-19 in the population of Mozambique, East Africa. *Psychology Research and Behavior Management*, *14*, 345-354. <https://doi.org/10.2147/PRBM.S298948>
- Giordani, R. C. F., Zanoni da Silva, M., Muhl, C., & Giolo, S. R. (2020). Fear of COVID-19 Scale: Assessing fear of the coronavirus pandemic in Brazil. *Journal of Health Psychology*, *0*(0), 1-12. <https://doi.org/10.1177/1359105320982035>
- Gregorich, S. E. (2006). Do self-report instruments allow meaningful comparisons across diverse population groups?: Testing measurement invariance using the confirmatory factor analysis framework. *Medical Care*, *44*(Suppl 3), S78-S94. <https://doi.org/10.1097/01.mlr.0000245454.12228.8f>
- Hair, J., Black, W., Babin, B., & Anderson, R. (2009). *Multivariate data analysis*. Prentice Hall.
- Harper, C. A., Satchell, L. P., Fido, D., & Latzman, R. D. (2020). Functional fear predicts public health compliance in the COVID-19 pandemic. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00281-5>
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., Ballard, C., Christensen, H., Cohen Silver, R., Everall, I., Ford, T., John, A., Kabir, T., King, K., Madan, I., Michie, S., Przybylski, A. K., Shafran, R., Sweeney, A., ... Bullmore, E. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: A call for action for mental health science. *The Lancet Psychiatry*, *7*(6), 547-560. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1)
- Hoyle, R. H. (1995). *Structural equation modeling: Concepts, issues, and applications* (1st ed.). Sage Publications.
- Huarcaya-Victoria, J., Villarreal-Zegarra, D., Podestà, A., & Luna-Cuadros, M. A. (2020). Psychometric properties of a Spanish version of the Fear of COVID-19 Scale in general population of Lima, Peru. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00354-5>
- INE (2020). *População residente com 15 e mais anos: Total e por nível de escolaridade completo mais elevado*. PORDATA. <https://www.pordata.pt/Portugal/Popula%C3%A7%C3%A3o+residente+com+15+e+mais+anos+total+e+por+n%C3%ADvel+de+escolaridade+completo+mais+elevado-2101-169769>
- Iversen, M. M., Norekvål, T. M., Oterhals, K., Fadnes, L. T., Måland, S., Pakpour, A. H., & Breivik, K. (2021). Psychometric properties of the Norwegian version of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*, *20*(3), 1446-1464. <https://doi.org/10.1007/s11469-020-00454-2>
- Kim, K. H., & Zabelina, D. (2015). Cultural bias in assessment: Can creativity assessment help?. *International Journal of Critical Pedagogy*, *6*(2), 129-148.
- Kline, P. (2000). *The handbook of psychological testing* (2nd ed.). Routledge.
- Koçak, O., Koçak, Ö. E., & Younis, M. Z. (2021). The psychological consequences of COVID-19 fear and the moderator effects of individuals' underlying illness and witnessing infected friends and family. *International Journal of Environmental Research and Public Health*, *18*, Article 1836. <https://doi.org/10.3390/ijerph18041836>
- Lathabhavan, R. (2021). A psychometric analysis of Fear of COVID-19 Scale in India. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-021-00657-1>
- Lee, S. A., Jobe, M. C., Mathis, A. A., & Gibbons, J. A. (2020). Incremental validity of coronaphobia: Coronavirus anxiety explains depression, generalized anxiety, and death anxiety. *Journal of Anxiety Disorders*, *74*, Article 102268. <https://doi.org/10.1016/j.janxdis.2020.102268>

- Leventhal, H. (1970). Findings and theory in the study of fear communications. *Advances in Experimental Social Psychology*, 5, 119–186. [https://doi.org/10.1016/S0065-2601\(08\)60091-X](https://doi.org/10.1016/S0065-2601(08)60091-X)
- Lin, C.-Y. (2020). Social reaction toward the 2019 novel coronavirus (COVID-19). *Social Health and Behavior*, 3(1), 1–2. https://doi.org/10.4103/SHB.SHB_11_20
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy*, 33(3), 335–342. [https://doi.org/10.1016/0005-7967\(94\)00075-u](https://doi.org/10.1016/0005-7967(94)00075-u)
- Magano, J., Vidal, D. G., Sousa, H. F. P. e, Dinis, M. A. P., & Leite, Á. (2021). Validation and psychometric properties of the Portuguese version of the Coronavirus Anxiety Scale (CAS) and Fear of COVID-19 Scale (FCV-19S) and associations with travel, tourism and hospitality. *International Journal of Environmental Research and Public Health*, 18(2), 427. <https://doi.org/10.3390/ijerph18020427>
- Mahmood, Q. K., Jafree, S. R., & Qureshi, W. A. (2020). The psychometric validation of FCV19S in Urdu and socio-demographic association with fear in the people of the Khyber Pakhtunkhwa (KPK) province in Pakistan. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00371-4>
- Mailliez, M., Griffiths, M. D., & Carre, A. (2021). Validation of the French version of the Fear of COVID-19 Scale and its associations with depression, anxiety, and differential emotions. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-021-00499-x>
- Martínez-Lorca, M., Martínez-Lorca, A., Criado-Álvarez, J. J., Armesilla, M. D. C., & Latorre, J. M. (2020). The Fear of COVID-19 Scale: Validation in Spanish university students. *Psychiatry Research*, 293, Article 113350. <https://doi.org/10.1016/j.psychres.2020.113350>
- Masuyama, A., Shinkawa, H., & Kubo, T. (2020). Validation and psychometric properties of the Japanese version of the Fear of COVID-19 Scale among adolescents. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00368-z>
- McDonald, R. P. (1999). Test theory: A unified treatment. Lawrence Erlbaum Associates Publishers.
- McLean, C. P., & Anderson, E. R. (2009). Brave men and timid women? A review of the gender differences in fear and anxiety. *Clinical Psychology Review*, 29(6), 496–505. <https://doi.org/10.1016/j.cpr.2009.05.003>
- Meade, A. W., Johnson, E. C., & Braddy, P. W. (2008). Power and sensitivity of alternative fit indices in tests of measurement invariance. *Journal of Applied Psychology*, 93(3), 568–592. <https://doi.org/10.1037/0021-9010.93.3.568>
- Medeiros, E. D., Reis, L. M., Guimarães, C. L. C., da Silva, P. G. N., Monteiro, R. P., Coelho, G. L. de H., Guimarães, C. M. C., Martins, E. R. dos S., & de França, L. L. A. (2021). Psychometric properties of the Brazilian version of the Fear of COVID-19 Scale (FCV-19S). *Current Psychology*, 20, 1–10. <https://doi.org/10.1007/s12144-021-01476-2>
- Midorikawa, H., Aiba, M., Lebowitz, A., Taguchi, T., Shiratori, Y., Ogawa, T., Takahashi, A., Takahashi, S., Nemoto, K., Arai, T., & Tachikawa, H. (2021). Confirming validity of the Fear of COVID-19 Scale in Japanese with a nationwide large-scale sample. *PLoS ONE*, 16(2), Article e0246840. <https://doi.org/10.1371/journal.pone.0246840>
- Nguyen, N. P. T., Hoang, T. D., Tran, V. T., Vu, C. T., Siewe Fodjo, J. N., Colebunders, R., Dunne, M. P., & Vo, T. V. (2020). Preventive behavior of Vietnamese people in response to the COVID-19 pandemic. *PLoS ONE*, 15(9), Article e0238830. <https://doi.org/10.1371/journal.pone.0238830>
- Nicola, M., Alsafi, Z., Sohrabi, C., Kerwan, A., Al-Jabir, A., Iosifidis, C., Agha, M., & Agha, R. (2020). The socio-economic implications of the coronavirus pandemic (COVID-19): A review. *International Journal of Surgery*, 78, 185–193. <https://doi.org/10.1016/j.ijsu.2020.04.018>
- Nitschke, J. P., Forbes, P. A. G., Ali, N., Cutler, J., Apps, M. A. J., Lockwood, P. L., & Lamm, C. (2020). Resilience during uncertainty? Greater social connectedness during COVID-19 lockdown is associated with reduced distress and fatigue. *British Journal of Health Psychology*, 26(2), 553–569. <https://doi.org/10.1111/bjhp.12485>
- Okereke, M., Ukor, N. A., Adebisi, Y. A., Ogunkola, I. O., Favour Iyagbaye, E., Adiola Owhor, G., & Lucero-Prisno, D. E. (2021). Impact of COVID-19 on access to healthcare in low- and middle-income countries: Current evidence and future recommendations. *The International Journal of Health Planning and Management*, 36(1), 13–17. <https://doi.org/10.1002/hpm.3067>
- Olatunji, B. O., Sawchuk, C. N., Arrindell, W. A., & Lohr, J. M. (2005). Disgust sensitivity as a mediator of the sex differences in contamination fears. *Personality and Individual Differences*, 38(3), 713–722. <https://doi.org/10.1016/j.paid.2004.05.025>
- Ornell, F., Schuch, J. B., Sordi, A. O., & Kessler, F. H. P. (2020). “Pandemic fear” and COVID-19: Mental health burden and strategies. *Brazilian Journal of Psychiatry*, 42(3), 232–235. <https://doi.org/10.1590/1516-4446-2020-0008>
- Pais-Ribeiro, J., Honrado, A., & Leal, I. (2004). Contribuição para o estudo da adaptação Portuguesa das Escalas de Ansiedade, Depressão e Stress (EADS) de 21 Itens de Lovibond e Lovibond. *Psicologia, Saúde e Doenças*, 5(2), 229–239.
- Pang, N. T. P., Kamu, A., Hambali, N. L. B., Mun, H. C., Kassim, M. A., Mohamed, N. H., Ayu, F., Rahim, S. S. S. A., Omar, A., & Jeffree, M. S. (2020). Malay version of the Fear of COVID-19 Scale: Validity and reliability. *International Journal of Mental Health and Addiction*, 20(1), 263–272. <https://doi.org/10.1007/s11469-020-00355-4>
- Perz, C. A., Lang, B. A., & Harrington, R. (2020). Validation of the Fear of COVID-19 Scale in a US college sample. *International Journal of Mental Health and Addiction*, 20(1), 273–283. <https://doi.org/10.1007/s11469-020-00356-3>
- Pilch, I., Kurasz, Z., & Turska-Kawa, A. (2021). Experiencing fear during the pandemic: Validation of the Fear of COVID-19 Scale in Polish. *PeerJ*, 9, Article e11263. <https://doi.org/10.7717/peerj.11263>
- Piqueras, J. A., Gomez-Gomez, M., Marzo, J. C., Gomez-Mir, P., Falco, R., Valenzuela, B., PSICORECUR-SOS COVID-19 study group, Falcó, R., Lopez-Nuñez, A., Martínez-González, A. E., Marzo, J. C., Mateu, O., & Moreno-Amador, B. (2021). Validation of the Spanish version of Fear of COVID-19 Scale: Its association with acute stress and coping its association with acute stress and coping. *International Journal of Mental Health and Addiction*, 8, 1–18. <https://doi.org/10.1007/s11469-021-00615-x>
- Pyszczynski, T., Lockett, M., Greenberg, J., & Solomon, S. (2020). Terror management theory and the COVID-19 pandemic. *Journal of Humanistic Psychology*, 61(2), 173–189. <https://doi.org/10.1177/0022167820959488>
- Reznik, A., Gritsenko, V., Konstantinov, V., Khamenka, N., & Isralowitz, R. (2020). COVID-19 fear in Eastern Europe: Validation of the Fear of COVID-19 Scale. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00283-3>
- Robinson, O. J., Vytal, K., Cornwell, B. R., & Grillon, C. (2013). The impact of anxiety upon cognition: Perspectives from human threat of shock studies. *Frontiers in Human Neuroscience*, 7, 1–21. <https://doi.org/10.3389/fnhum.2013.00203>
- Roest, A. M., Jonge, P., Lim, C., Stein, D. J., Al-Hamzawi, A., Alonso, J., Benjet, C., Bruffaerts, R., Bunting, B., Caldas-de-Almeida, J. M., Ciutan, M., Girolamo, G., Hu, C., Levinson, D., Nakamura, Y., Navarro-Mateu, F., Piazza, M., Posada-Villa, J., Torres, Y., ... Scott, K. M. (2017). Fear and distress disorders as predictors of heart disease: A temporal perspective. *Journal of Psychosomatic Research*, 96, 65–75. <https://doi.org/10.1016/j.jpsychores.2017.03.015>

- Ropeik, D. (2004). The consequences of fear: Our modern world is a risky place and evokes many well-founded fears. But these fears themselves create a new risk for our health and well-being that needs to be addressed [Special issue]. *EMBO Reports*, 5, 56–60. <https://doi.org/10.1038/sj.embor.7400228>
- Rosenberg, J. (2017). *The effects of chronic fear on a person's health*. The American Journal of Managed Care. <https://www.ajmc.com/view/the-effects-of-chronic-fear-on-a-persons-health>
- Sakib, N., Bhuiyan, A. K. M. I., Hossain, S., Al Mamun, F., Hosen, I., Abdullah, A. H., Sarker, Md. A., Mohiuddin, M. S., Rayhan, I., Hossain, M., Sikder, Md. T., Gozal, D., Muhit, M., Islam, S. M. S., Griffiths, M. D., Pakpour, A. H., & Mamun, M. A. (2020). Psychometric validation of the Bangla Fear of COVID-19 Scale: Confirmatory factor analysis and rasch analysis. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00289-x>
- Satici, B., Gocet-Tekin, E., Deniz, M. E., & Satici, S. A. (2020). Adaptation of the Fear of COVID-19 Scale: Its association with psychological distress and life satisfaction in Turkey. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00294-0>
- Schermelleh-Engel, K., Moosbrugger, H., & Müller, H. (2003). Evaluating the fit of structural equation models: Tests of significance and descriptive goodness-of-fit measures. *Methods of Psychological Research Online*, 8(2), 23–74.
- Schmitt, N., & Kuljanin, G. (2008). Measurement invariance: Review of practice and implications. *Human Resource Management Review*, 18(4), 210–222. <https://doi.org/10.1016/j.hrmmr.2008.03.003>
- Seegerstrom, S. C., & Miller, G. E. (2004). Psychological stress and the human immune system: A meta-analytic study of 30 years of inquiry. *Psychological Bulletin*, 130(4), 601–630. <https://doi.org/10.1037/0033-2909.130.4.601>
- Solano-Flores, G., & Nelson-Barber, S. (2001). On the cultural validity of science assessments. *Journal of Research in Science Teaching*, 38(5), 553–573. <https://doi.org/10.1002/tea.1018>
- Soper, D. (2018). *Free statistics calculators*. DanielSoper.com. <https://www.danielsoper.com/statcalc/default.aspx>
- Soraci, P., Ferrari, A., Abbiati, F. A., Del Fante, E., De Pace, R., Urso, A., & Griffiths, M. D. (2020). Validation and psychometric evaluation of the Italian version of the fear of COVID-19 scale. *International Journal of Mental Health and Addiction*, 4, 1–10. <https://doi.org/10.1007/s11469-020-00277-1>
- Suzuki, L. A., & Ponterotto, J. G. (2007). *Handbook of multicultural assessment: Clinical, psychological, and educational applications* (3rd ed.). Jossey-Bass.
- Tolin, D. F., & Foa, E. B. (2008). Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychological Trauma: Theory, Research, Practice, and Policy*, 5(1), 37–85. <https://doi.org/10.1037/1942-9681.S.1.37>
- Tsipropoulou, V., Nikopoulou, V. A., Holeva, V., Nasika, Z., Diakogiannis, I., Sakka, S., Kostikidou, S., Varvara, C., Spyridopoulou, E., & Parlapani, E. (2020). Psychometric properties of the Greek version of FCV-19S. *International Journal of Mental Health and Addiction*. <https://doi.org/10.1007/s11469-020-00319-8>
- Usher, K., Durkin, J., & Bhullar, N. (2020). The COVID-19 pandemic and mental health impacts. *International Journal of Mental Health Nursing*, 29, 315–318. <https://doi.org/10.1111/inm.12726>
- Usher, K., Jackson, D., Durkin, J., Gyamfi, N., & Bhullar, N. (2020). Pandemic-related behaviours and psychological outcomes; A rapid literature review to explain COVID-19 behaviours. *International Journal of Mental Health Nursing*, 29(6), 1018–1034. <https://doi.org/10.1111/inm.12790>
- Vandenberg, R. J., & Lance, C. E. (2000). Review and synthesis of the measurement invariance literature: Suggestions, practices, and recommendations for organizational research. *Organizational Research Methods*, 3(1), 4–70. <https://doi.org/10.1177/109442810031002>
- Wakashima, K., Asai, K., Kobayashi, D., Koiwa, K., Kamoshida, S., & Sakuraba, M. (2020). The Japanese version of the fear of COVID-19 scale: Reliability, validity, and relation to coping behavior. *PLoS One*, 15(11), Article e0241958. <https://doi.org/10.1371/journal.pone.0241958>
- Wang, Y., Shi, L., Que, J., Lu, Q., Liu, L., Lu, Z., Xu, Y., Liu, J., Sun, Y., Meng, S., Yuan, K., Ran, M., Lu, L., Bao, Y., & Shi, J. (2021). The impact of quarantine on mental health status among general population in China during the COVID-19 pandemic. *Molecular Psychiatry*, 26(9), 4813–4822. <https://doi.org/10.1038/s41380-021-01019-y>
- Wheaton, B. (1987). Assessment of fit in overidentified models with latent variables. *Sociological Methods & Research*, 16(1), 118–154. <https://doi.org/10.1177/0049124187016001005>
- Winter, T., Riordan, B. C., Pakpour, A. H., Griffiths, M. D., Mason, A., Poulgrain, J. W., & Scarf, D. (2020). Evaluation of the English version of the fear of COVID-19 scale and its relationship with behavior change and political beliefs. *International Journal of Mental Health and Addiction*, 15, 1–11. <https://doi.org/10.1007/s11469-020-00342-9>
- Worldometer. (2020). *COVID-19 coronavirus pandemic*. <https://www.worldometers.info/coronavirus/>

Appendix 1

Summary of Previous Research on FCV-19S Psychometric Properties

Reference	Country	<i>n</i>	PCA (Varimax rotation)	CFA (Maximum likelihood estimation)	Cronbach's Alpha	F vs. M
Ahorsu et al. (2020)	Iran	717	1 factor Variance explained: .51 Factor loadings > .66	-	.82	-
Alyami et al. (2020)	Saudi Arabia	639	-	1 factor χ^2/df = not reported; CFI = 1; RMSEA = .06; SRMR = .02; Saturation levels > .62	.88	-
Barbosa-Camacho et al. (2021) ^a	Mexico	1216	-	1 factor χ^2/df = 1.32; CFI = .99; TLI = 0.99; RMSEA = .02; SRMR = .01; AIC = 51.23 Saturation levels > .53	1 factor = .85	F > M
Barbosa-Camacho et al. (2021) ^a	Mexico	1216	-	2 factors χ^2/df = 2.02; CFI = .99; TLI = 0.99; RMSEA = .03; SRMR = .01; AIC = 56.17 Saturation levels > .56	Factor 1 = not reported Factor 2 = not reported	F > M
Barrios et al. (2020)	Paraguay	1077	2 factors Variance explained: .61 Factor loadings > .64	2 factors χ^2/df = 1.53; CFI = 1; NFI = .99; GFI = 1; TLI = 0.99; RMSEA = .03 Saturation levels > .54 <i>r</i> between factors: .72 ≤ <i>r</i> ≤ .95	Global = .86 Factor 1 = .84 Factor 2 = .82	-
Basit et al. (2021)	Pakistan	380	3 factors Variance explained: .96 Factor loadings > .52	-	Global = .88 Factor 1 = .90 Factor 2 = .92 Factor 3 = .68	F > M
Bitan et al. (2020)	Israel	639	2 factors Variance explained: .66 Factor loadings > .57	-	Factor 1 = .77 Factor 2 = .80	F > M
Cavalheiro & Sticca (2020)	Brazil	354	1 factor Variance explained: .54 Factor loadings > .65	1 factor χ^2/df = 3.29; CFI = .99; GFI = .98; TLI = 0.98; NFI = .98; RMSEA = .06 Saturation levels > .57	.88	F > M
Caycho-Rodríguez et al. (2020)	Argentina	1291	-	2 factors χ^2/df = 6.82; CFI = .98; TLI = 0.97; RMSEA = .07; SRMR = .03 Saturation levels > .60 <i>r</i> between factors: not reported	Factor 1 = .80 Factor 2 = .73	-
Chang et al. (2020)	Taiwan	400	-	1 factor χ^2/df = 2.26; CFI = .99; RMSEA = .06; SRMR = .06 Saturation levels > .68	.93	-
Chi et al. (2021)	China	793	-	2 factors χ^2/df = 5.36; CFI = .99; TLI = 0.98; RMSEA = .07; SRMR = .03 Saturation levels > .56 <i>r</i> between factors: .72	Global = .92 Factor 1 = .89 Factor 2 = .84	F > M
El-Bardan & Lathabhavan (2021)	Egypt	1832	-	1 factor χ^2/df = 2.26; CFI = .92; TLI = 0.89; GFI = .92; AGFI = .91; RMSEA = .07; SRMR = .04 Saturation levels > .66	.87	-
Elemo et al. (2020)	Ethiopia	307	-	1 factor χ^2/df = 5.92; CFI = .93; NFI = .92; IFI = .93; SRMR = .59 Saturation levels > .45	.87	-
Faro et al. (2020)	Brazil	1000	-	1 factor χ^2/df = 5.92; CFI = .99; GFI = .99; TLI = 0.98; RMSEA = .07; SRMR = .06 Saturation levels > .57	.86	-

Reference	Country	<i>n</i>	PCA (Varimax rotation)	CFA (Maximum likelihood estimation)	Cronbach's Alpha	F vs. M
Giordani et al. (2020)	Brazil	7430	1 factor Variance explained: .55 Factor loadings > .71	-	.86	F > M
Giordani et al. (2021)	Mozambique	387	-	1 factor $\chi^2/df = 3.54$; CFI = .98; TLI = 0.97; RMSEA = .08; SRMR = .08 Saturation levels > .69	.87	F > M
Huarcaya-Victoria et al. (2020) ^b	Peru	832	2 factors Variance explained: not reported Factor loadings > .49	2 factors $\chi^2/df = 5.69$; CFI = .99; TLI = 0.96; RMSEA = .08; SRMR = .02; AIC = 15,008.8; BIC = 15,108 Saturation levels > .53 <i>r</i> between factors: not reported	Factor 1 = .85 Factor 2 = .89	F > M
Iversen et al. (2021)	Norway	1063	-	2 factors $\chi^2/df = 3.64$; CFI = .99; RMSEA = .05; SRMR = .02 Saturation levels > .69 <i>r</i> between factors: .84	Global = .88 Factor 1 = not reported Factor 2 = not reported	F > M
Lathabhavan (2021) ^c	India	992	-	1 factor $\chi^2/df = 2.31$; CFI = .94; GFI = .93; AGFI = .87; TLI = 0.91; RMSEA = .10; SRMR = .05 Saturation levels > .57	.85	-
Lathabhavan (2021) ^c	India	992	-	2 factors $\chi^2/df = 1.74$; CFI = .96; GFI = .95; AGFI = .89; TLI = 0.95; RMSEA = .07; SRMR = .04 Saturation levels > .71	Factor 1 = .82 Factor 2 = .86	-
Magano et al. (2021)	Portugal	561	2 factors Variance explained: .71 Factor loadings > .71	2 factors $\chi^2/df = 3.58$; CFI = .99; TLI = 0.97; RMSEA = .07; SRMR = .03; Saturation levels > .67 <i>r</i> between factors: not reported	Factor 1 = .83 Factor 2 = .82	F > M
Mahmood et al. (2020)	Pakistan	501	-	1 factor $\chi^2/df = 3.57$; CFI = .99; GFI = .98; AGFI = .94; TLI = 0.97; RMSEA = .07; RMSR = .04 Saturation levels > .59	.85	F > M
Maillez et al. (2021)	France	316	-	1 factor $\chi^2/df = 5.16$; CFI = 1; TLI = 1.03; RMSEA = 0; SRMR = .02 Saturation levels > .58	.87	-
Martínez-Lorca et al. (2020)	Spain	606	1 factor Variance explained: .49 Factor loadings > .61	-	.86	-
Masuyama et al. (2020)	Japan	629	-	2 factors $\chi^2/df = 1.72$; CFI = 1; RMSEA = .03; SRMR = .04 Saturation levels > .56 <i>r</i> between factors: not reported	Global = .82 Factor 1 = .71 Factor 2 = .82	-
Medeiros et al. (2020) ^d	Brazil	230	1 factor Variance explained: .65 Factor loadings > .70	-	.91	-
Medeiros et al. (2020) ^d	Brazil	302	-	1 factor $\chi^2/df =$ not reported; CFI = .99; TLI = 0.98; RMSEA = .07; AVE = .50; CR = .87 Saturation levels > .55	.87	-
Midorikawa et al. (2020)	Japan	6750	-	2 factors $\chi^2/df = 13.68$; CFI = .99; TLI = 0.99; RMSEA = .04; AIC = 196.15 Saturation levels > .51 <i>r</i> between factors: .67	Factor 1 = .77 Factor 2 = .83	F > M

Reference	Country	<i>n</i>	PCA (Varimax rotation)	CFA (Maximum likelihood estimation)	Cronbach's Alpha	F vs. M
Nguyen et al. (2020)	Vietnam	5423	1 factor Variance explained: .62 Factor loadings > .69	-	.90	F > M
Pang et al. (2020)	Malaysia	228	-	1 factor AVE = .40; CR = .80 Saturation levels > not reported	.89	-
Perz et al. (2020)	USA	237	1 factor Variance explained: .66 Factor loadings > .70	-	.91	F = M
Pilch et al. (2021) ^e	Poland	708	-	1 factor $\chi^2/df = 5.05$; CFI = .98; TLI = 0.97; RMSEA = .07; GFI = .99 Saturation levels > not reported	Sample 1 = .89 Sample 2 = .85	F > M
Piqueras et al. (2021)	Spain & Dominican Republic	1146	-	1 factor $\chi^2/df = 6.14$; R-CFI = .98; TLI = 0.97; R-RMSEA = .06; SRMR = .02 AVE = .44; CR = .85 Saturation levels > .54	.86	F > M
Reznik et al. (2020)	Russia & Belarus	850	2 factors Variance explained: .49 Factor loadings > .57	-	Global = .81	F > M
Sakib et al. (2020)	Bangladesh	8550	-	1 factor $\chi^2/df = 39.63$; CFI = .96; TLI = 0.95; RMSEA = .07; WRMSR = 0.89 Saturation levels > .72	.87	F > M
Satici et al. (2020)	Turkey	1304	-	1 factor $\chi^2/df = 23.04$; CFI = .92; GFI = .94; NFI = .91; IFI = .92; SRMR = .06 Saturation levels > .62	.85	-
Soraci et al. (2020)	Italy	249	-	1 factor $\chi^2/df = 2.16$; CFI = .99; TLI = 0.99; RMSEA = .07; SRMR = .05 Saturation levels > .69	.87	-
Tsipropoulou et al. (2020)	Greece	2970	-	1 factor $\chi^2/df = 33.91$; CFI = .89; TLI = 0.83; RMSEA = .11; SRMR = .06 Saturation levels > .45	.87	F > M
Wakashima et al. (2020)	Japan	450	-	1 factor $\chi^2/df = 5.92$; CFI = .94; RMSEA = .11; SRMR = .05; BIC = 7,817.05 Saturation levels > .45	.87	F = M
Winter et al. (2020) ^f	New Zealand	1397	-	1 factor $\chi^2/df =$ not reported; CFI = .90; RMSEA = .16 Saturation levels > .64	.89	-
Winter et al. (2020) ^f	New Zealand	1023	-	1 factor $\chi^2/df =$ not reported; CFI = .92; RMSEA = .13 Saturation levels > .67	.88	-

Notes:

^aBarbosa-Camacho and colleagues (2021) validated two FCV-19Ss with different factor structures using the same sample. ^bTo perform EFA and CFA, Huarcaya-Victoria and colleagues (2020) split the sample in the half.

^cLathabhavan (2021) validated two FCV-19Ss with different factor structures using the same sample.

^dMedeiros and colleagues (2020) analyzed both EFA and CFA using two different samples.

^ePilch and colleagues (2021) performed CFA using the total sample ($n = 708$), but calculated Cronbach's alpha for sample 1 ($n = 383$) and sample 2 ($n = 325$).

^fWinter and colleagues (2020) performed CFA using two different samples.

Appendix 2

European Portuguese version of the Fear of COVID-19 Scale (FCV-19S-P)

Usando uma escala de 1 a 5, em que 1 indica "Discordo totalmente" e 5 indica "Concordo totalmente", indique o quanto concorda com cada uma das afirmações:

1. Aquilo de que tenho mais medo é da COVID-19.
2. Pensar na COVID-19 deixa-me desconfortável.
3. Fico com as mãos suadas quando penso na COVID-19.
4. Tenho medo de perder a minha vida por causa da COVID-19.
5. Fico nervoso/a ou ansioso/a quando vejo as notícias e histórias sobre a COVID-19 nos meios de comunicação social.
6. Não consigo dormir por causa da preocupação de vir a apanhar COVID-19.
7. O meu coração dispara ou tenho palpitações quando penso em apanhar COVID-19.

Appendix 3

Inter-Items Correlations of the FCV-19S-P

	Item 1	Item 2	Item 3	Item 4	Item 5	Item 6
Item 1	-	-	-	-	-	-
Item 2	.610*	-	-	-	-	-
Item 3	.346*	.392*	-	-	-	-
Item 4	.511*	.470*	.303*	-	-	-
Item 5	.491*	.619*	.384*	.488*	-	-
Item 6	.326*	.341*	.575*	.297*	.395*	-
Item 7	.386*	.415*	.566*	.380*	.432*	.644*

* $p < .01$.