

QUILLEN EQUIVALENCES FOR STABLE CATEGORIES

GEORGIOS DALEZIOS, SERGIO ESTRADA, AND HENRIK HOLM

ABSTRACT. For an abelian category \mathcal{A} we investigate when the stable categories $\underline{\text{GProj}}(\mathcal{A})$ and $\underline{\text{GInj}}(\mathcal{A})$ are triangulated equivalent. To this end, we realize these stable categories as homotopy categories of certain (non-trivial) model categories and give conditions on \mathcal{A} that ensure the existence of a Quillen equivalence between the model categories in question. We also study when such a Quillen equivalence transfers from \mathcal{A} to the category of chain complexes in \mathcal{A} .

1. INTRODUCTION

Over an Iwanaga–Gorenstein ring A , that is, a ring which is noetherian and has finite injective dimension from both sides, the category $\text{MCM}(A)$ of (finitely generated) maximal Cohen–Macaulay A -modules¹ is a Frobenius category in which the projective-injective objects are precisely the finitely generated projective A -modules. The associated stable category $\underline{\text{MCM}}(A)$ is therefore triangulated, and a classic result of Buchweitz [8, Thm. 4.4.1] shows that $\underline{\text{MCM}}(A)$ is triangulated equivalent to the singularity category² $\mathcal{D}_{\text{sg}}(A)$, which is an important mathematical object that has been studied by many authors; see [5, 6, 27, 33].

If A is not Iwanaga–Gorenstein, then the category $\text{MCM}(A)$ is, in general, not Frobenius. However, over any ring A one can always consider the category $\text{GProj}(A)$ of so-called Gorenstein projective modules (which are not assumed to be finitely generated); this category is always Frobenius and the associated stable category $\underline{\text{GProj}}(A)$ is triangulated. In the case where A is Iwanaga–Gorenstein, an A -module is maximal Cohen–Macaulay if and only if it is finitely generated and Gorenstein projective, and hence $\underline{\text{MCM}}(A)$ can be identified with the finitely generated modules in $\underline{\text{GProj}}(A)$. This explains the interest in the category $\underline{\text{GProj}}(A)$ for general ring A . Its injective counterpart $\underline{\text{GInj}}(A)$, the stable category of Gorenstein injective A -modules, is equally important and has been studied in e.g [7, 26].

Our work is motivated by a recent result of Zheng and Huang [37] which asserts that for many rings A , the categories $\underline{\text{GProj}}(A)$ and $\underline{\text{GInj}}(A)$ are equivalent as triangulated categories. As it makes sense to consider the stable categories $\underline{\text{GProj}}(\mathcal{A})$ and $\underline{\text{GInj}}(\mathcal{A})$ for any

2010 *Mathematics Subject Classification.* 18E10, 18E30, 18G55, 13D02, 16E05.

The first author is supported by the Fundación Séneca of Murcia 19880/GERM/15. The second author is supported by the grant MTM2016-77445-P and FEDER funds and the grant 18394/JLI/13 by the Fundación Séneca-Agencia de Ciencia y Tecnología de la Región de Murcia in the framework of III PCTRM 2011-2014 .

¹In the important special case where A is a quasi-Frobenius ring, for example, if $A = kG$ is the group algebra of a finite group G with coefficients in a field k , the category $\text{MCM}(A)$ is just the category $\text{mod}(A)$ of all finitely generated A -modules.

²The singularity category $\mathcal{D}_{\text{sg}}(A)$ is defined to be the Verdier quotient $\mathcal{D}^b(A)/\mathcal{D}_{\text{perf}}^b(A)$ of the bounded derived category $\mathcal{D}^b(A)$, whose objects are complexes of A -modules with bounded and finitely generated homology, by the subcategory $\mathcal{D}_{\text{perf}}^b(A)$, whose objects are isomorphic (in $\mathcal{D}^b(A)$) to a perfect complex, that is, to a bounded complex of finitely generated projective A -modules. The name *singularity category* and the symbol $\mathcal{D}_{\text{sg}}(A)$ seem to be the popular choices nowadays, however, in the work of Buchweitz [8, Def. 1.2.2], this category is called the *stabilized derived category* and denoted by $\overline{\mathcal{D}^b(A)}$, and in the work of Orlov [29], it is called the *triangulated category of singularities* and denoted by $\mathcal{D}_{\text{sg}}(A)$.

bicomplete abelian category \mathcal{A} with enough projectives and injectives (see Section 2 for details), the following question naturally arises:

Question. For which abelian categories \mathcal{A} (assumed to be bicomplete with enough projectives and injectives) are $\underline{\mathbf{GProj}}(\mathcal{A})$ and $\underline{\mathbf{GInj}}(\mathcal{A})$ equivalent as triangulated categories?

Every Frobenius category \mathcal{E} , in particular, $\mathbf{GProj}(\mathcal{A})$ and $\mathbf{GInj}(\mathcal{A})$, can be equipped with a canonical model structure which has the property that the associated homotopy category $\mathrm{Ho}(\mathcal{E})$ is equivalent to the stable category $\underline{\mathcal{E}}$; see e.g. [18, Prop. 4.1]. Thus, if the Frobenius categories $\mathbf{GProj}(\mathcal{A})$ and $\mathbf{GInj}(\mathcal{A})$, equipped with these canonical model structures, happen to be Quillen equivalent, then we get an affirmative answer to the question above. However, the model categories $\mathbf{GProj}(\mathcal{A})$ and $\mathbf{GInj}(\mathcal{A})$, and even the underlying ordinary categories, will rarely be (Quillen) equivalent. In this paper, we consider instead the categories

$$\mathcal{U}^\pi = \{M \in \mathcal{A} \mid \mathrm{Gpd}_{\mathcal{A}}(M) < \infty\} \quad \text{and} \quad \mathcal{U}^l = \{N \in \mathcal{A} \mid \mathrm{Gid}_{\mathcal{A}}(N) < \infty\}$$

and show in Theorems 3.7 and 3.9 that \mathcal{U}^π and \mathcal{U}^l can be equipped with model structures for which the associated homotopy categories $\mathrm{Ho}(\mathcal{U}^\pi)$ and $\mathrm{Ho}(\mathcal{U}^l)$ are the stable categories $\underline{\mathbf{GProj}}(\mathcal{A})$ and $\underline{\mathbf{GInj}}(\mathcal{A})$. The advantage of having these realizations of the stable categories is that in several cases the model categories \mathcal{U}^π and \mathcal{U}^l will be Quillen equivalent—even though $\mathbf{GProj}(\mathcal{A})$ and $\mathbf{GInj}(\mathcal{A})$ are not—and in such cases we therefore get an affirmative answer (for a strong reason) to the question above³. To investigate when \mathcal{U}^π and \mathcal{U}^l will be Quillen equivalent, we introduce the notion of a Sharp–Foxby adjunction (Definition 3.4). We prove in Theorem 3.11 and Corollary 3.12 that if \mathcal{A} admits such an adjunction, then \mathcal{U}^π and \mathcal{U}^l will be Quillen equivalent:

Theorem A. *A Sharp–Foxby adjunction (S, T) on \mathcal{A} induces a Quillen equivalence between the model categories \mathcal{U}^π and \mathcal{U}^l . Thus the total (left/right) derived functors of S and T yield an adjoint equivalence of the corresponding homotopy categories,*

$$\underline{\mathbf{GProj}}(\mathcal{A}) \simeq \mathrm{Ho}(\mathcal{U}^\pi) \begin{array}{c} \xrightarrow{\mathbf{LS}} \\ \xleftarrow{\mathbf{RT}} \end{array} \mathrm{Ho}(\mathcal{U}^l) \simeq \underline{\mathbf{GInj}}(\mathcal{A}).$$

In fact, this is an equivalence of triangulated categories.

The choice to work with the categories \mathcal{U}^π and \mathcal{U}^l is historically motivated by classic results in commutative algebra by Sharp [31] and Foxby [14]. In the language of this paper, the results can be phrased as follows: If A is a Cohen–Macaulay ring with a dualizing module D , then the functors $S = D \otimes_A -$ and $T = \mathrm{Hom}_A(D, -)$ constitute a Sharp–Foxby adjunction on $\mathcal{A} = \mathrm{Mod}(A)$; see Example 3.6 for details. Thus, for such rings Theorem A improves the previously mentioned result of Zheng and Huang [37] to a triangulated equivalence between $\underline{\mathbf{GProj}}(\mathcal{A})$ and $\underline{\mathbf{GInj}}(\mathcal{A})$ induced by a Quillen equivalence.

In Section 4 we investigate to what extent a Sharp–Foxby adjunction on a category \mathcal{A} (and hence also a Quillen equivalence between the model categories \mathcal{U}^π and \mathcal{U}^l , see Theorem A) transfers to the category of chain complexes in \mathcal{A} . In 4.5 we obtain the following.

Theorem B. *Assume that (S, T) is a Sharp–Foxby adjunction on \mathcal{A} ; in particular, $\underline{\mathbf{GProj}}(\mathcal{A})$ and $\underline{\mathbf{GInj}}(\mathcal{A})$ are equivalent as triangulated categories by Theorem A. Assume furthermore that the finitistic projective and the finitistic injective dimensions of \mathcal{A} are finite.*

If $\mathcal{B} = \mathrm{Ch}(\mathcal{A})$, then degreewise application of S and T yields a Sharp–Foxby adjunction on \mathcal{B} ; in particular, $\underline{\mathbf{GProj}}(\mathcal{B})$ and $\underline{\mathbf{GInj}}(\mathcal{B})$ are equivalent as triangulated categories.

³In general, we do not expect every (triangulated) equivalence between $\underline{\mathbf{GProj}}(\mathcal{A})$ and $\underline{\mathbf{GInj}}(\mathcal{A})$, if such an equivalence even exist, to be induced from a Quillen equivalence between model categories. Indeed, it is well-known that there are examples of non Quillen equivalent model categories with equivalent homotopy categories.

2. PRELIMINARIES

Throughout this paper, \mathcal{A} denotes any bicomplete abelian category with enough projectives and enough injectives.

Gorenstein projective and Gorenstein injective modules (over any ring) were defined by Enochs and Jenda [10, §2], but the definition works for objects in any abelian category:

Definition 2.1. An acyclic (= exact) complex $P = \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow P_{-1} \rightarrow \cdots$ of projective objects in \mathcal{A} is called *totally acyclic* if for any projective object Q in \mathcal{A} the complex

$$\mathrm{Hom}_{\mathcal{A}}(P, Q) = \cdots \longrightarrow \mathrm{Hom}_{\mathcal{A}}(P_{-1}, Q) \longrightarrow \mathrm{Hom}_{\mathcal{A}}(P_0, Q) \longrightarrow \mathrm{Hom}_{\mathcal{A}}(P_1, Q) \longrightarrow \cdots$$

is acyclic. An object G in \mathcal{A} is called *Gorenstein projective* if it is a cycle of such a totally acyclic complex of projectives, that is, if $G = Z_j(P)$ for some integer j . We write $\mathrm{GProj}(\mathcal{A})$ for the full subcategory of \mathcal{A} consisting of all Gorenstein projective objects.

Dually, an acyclic complex $I = \cdots \rightarrow I_1 \rightarrow I_0 \rightarrow I_{-1} \rightarrow \cdots$ of injective objects in \mathcal{A} is called *totally acyclic* if for any injective object E in \mathcal{A} the complex

$$\mathrm{Hom}_{\mathcal{A}}(E, I) = \cdots \longrightarrow \mathrm{Hom}_{\mathcal{A}}(E, I_1) \longrightarrow \mathrm{Hom}_{\mathcal{A}}(E, I_0) \longrightarrow \mathrm{Hom}_{\mathcal{A}}(E, I_{-1}) \longrightarrow \cdots$$

is acyclic. An object H in \mathcal{A} is called *Gorenstein injective* if it is a cycle of such a totally acyclic complex of injectives, that is, if $H = Z_j(I)$ for some integer j . We write $\mathrm{GInj}(\mathcal{A})$ for the full subcategory of \mathcal{A} consisting of all Gorenstein injective objects.

The *Gorenstein projective dimension*, $\mathrm{Gpd}_{\mathcal{A}}(M)$, of an object M in \mathcal{A} is defined by declaring that one has $\mathrm{Gpd}_{\mathcal{A}}(M) \leq n$ (for $n \in \mathbb{N}_0$) if and only if there exists an exact sequence $0 \rightarrow G_n \rightarrow G_{n-1} \rightarrow \cdots \rightarrow G_0 \rightarrow M \rightarrow 0$ in \mathcal{A} with $G_0, \dots, G_n \in \mathrm{GProj}(\mathcal{A})$. The *Gorenstein injective dimension*, $\mathrm{Gid}_{\mathcal{A}}(M)$, of M is defined analogously.

Recall that a *Frobenius category* is an exact category \mathcal{E} with enough (relative) projectives and enough (relative) injectives and where the classes of projectives and injectives coincide; such objects are called *projective-injective* (or just *pro-injective*) objects. The *stable category* $\underline{\mathcal{E}}$ is the quotient category \mathcal{E}/\sim where the relation “ \sim ” is defined by $f \sim g$ (here f and g are parallel morphisms in \mathcal{E}) if $f - g$ factors through a projective-injective object. The category $\underline{\mathcal{E}}$ is triangulated as described in Happel [21, Chap. I§2] (see also 2.5).

The following result is well-known, but for completeness we include a short proof.

Proposition 2.2. *The category $\mathrm{GProj}(\mathcal{A})$ is Frobenius and the projective-injective objects herein are the projective objects in \mathcal{A} . Thus, the stable category $\underline{\mathrm{GProj}}(\mathcal{A})$ is triangulated.*

The category $\mathrm{GInj}(\mathcal{A})$ is Frobenius and the projective-injective objects herein are the injective objects in \mathcal{A} . Thus, the stable category $\underline{\mathrm{GInj}}(\mathcal{A})$ is triangulated.

Proof. We only show the claims about the category $\mathrm{GProj}(\mathcal{A})$, as the claims about $\mathrm{GInj}(\mathcal{A})$ are proved similarly. The proof only uses basic properties of Gorenstein projective objects. In the case of modules, that is, if $\mathcal{A} = \mathrm{Mod}(A)$ for a ring A , these properties are recorded in [23], however, the reader easily verifies that the same properties hold for Gorenstein projective objects in any abelian category \mathcal{A} with enough projectives.

First of all, by [23, Thm. 2.5] the class $\mathrm{GProj}(\mathcal{A})$ is an additive extension-closed subcategory of the abelian category \mathcal{A} , and thus $\mathrm{GProj}(\mathcal{A})$ is an exact category. Clearly, every (categorical) projective object P in \mathcal{A} is a (relative) projective object in $\mathrm{GProj}(\mathcal{A})$, but it is also (relative) injective since every short exact sequence $0 \rightarrow P \rightarrow G \rightarrow G' \rightarrow 0$ in \mathcal{A} with $G, G' \in \mathrm{GProj}(\mathcal{A})$ splits; indeed by [23, Prop. 2.3] one has $\mathrm{Ext}_{\mathcal{A}}^1(G', P) = 0$. By the definition of Gorenstein projective objects, every $G \in \mathrm{GProj}(\mathcal{A})$ fits into short exact sequences $0 \rightarrow H \rightarrow P \rightarrow G \rightarrow 0$ and $0 \rightarrow G \rightarrow P' \rightarrow H' \rightarrow 0$ in \mathcal{A} where P, P' are (categorical) projective and H, H' are Gorenstein projective. It follows that if G is (relative) projective or

(relative) injective, then G is a direct summand of a (categorical) projective object, P or P' , and hence G is (categorical) projective. It also follows that $\text{GProj}(\mathcal{A})$ has enough (relative) projectives and enough (relative) injectives. \square

In Theorems 3.7 and 3.9 we construct certain model categories \mathcal{U}^π and \mathcal{U}^t for which the associated homotopy categories $\text{Ho}(\mathcal{U}^\pi)$ and $\text{Ho}(\mathcal{U}^t)$ are $\text{GProj}(\mathcal{A})$ and $\text{GInj}(\mathcal{A})$.

The standard references for the theory of cotorsion pairs are Enochs and Jenda [11] and Göbel and Trlifaj [20]. Below we recall a few notions that we need.

2.3. A pair $(\mathcal{X}, \mathcal{Y})$ of classes of objects in \mathcal{A} is a *cotorsion pair* if $\mathcal{X}^\perp = \mathcal{Y}$ and $\mathcal{X} = {}^\perp\mathcal{Y}$. Here, given a class \mathcal{C} of objects in \mathcal{A} , the right orthogonal \mathcal{C}^\perp is defined to be the class of all $Y \in \mathcal{A}$ such that $\text{Ext}_{\mathcal{A}}^1(C, Y) = 0$ for all $C \in \mathcal{C}$. The left orthogonal ${}^\perp\mathcal{C}$ is defined similarly. A cotorsion pair $(\mathcal{X}, \mathcal{Y})$ is *hereditary* if $\text{Ext}_{\mathcal{A}}^i(X, Y) = 0$ for all $X \in \mathcal{X}$, $Y \in \mathcal{Y}$, and $i \geq 1$. A cotorsion pair $(\mathcal{X}, \mathcal{Y})$ is *complete* if it has *enough projectives* and *enough injectives*, i.e. for each $A \in \mathcal{A}$ there exist short exact sequences $0 \rightarrow Y \rightarrow X \rightarrow A \rightarrow 0$ (enough projectives) and $0 \rightarrow A \rightarrow Y' \rightarrow X' \rightarrow 0$ (enough injectives) with $X, X' \in \mathcal{X}$ and $Y, Y' \in \mathcal{Y}$.

In order for the above to make sense, the category \mathcal{A} only needs to be exact (not necessarily abelian), so that one has a notion of “short exact sequences” (often called *conflations*) and hence also of (Yoneda) $\text{Ext}_{\mathcal{A}}$.

Cotorsion pairs are related to relative homological algebra, see [11], and due to work of Hovey [25] they are also related to abelian (or exact) model category structures.

2.4. An *abelian model structure* on \mathcal{A} , that is, a model structure on \mathcal{A} which is compatible with the abelian structure in the sense of [25, Def. 2.1], corresponds by Thm. 2.2 in *loc. cit.* to a triple $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ of classes of objects in \mathcal{A} for which \mathcal{W} is thick⁴ and $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{W} \cap \mathcal{F})$ are complete cotorsion pairs in \mathcal{A} . Such a triple $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ is called a *Hovey triple* in \mathcal{A} . In the model structure on \mathcal{A} determined by such a Hovey triple, \mathcal{C} is precisely the class of cofibrant objects, \mathcal{F} is precisely the class of fibrant objects, and \mathcal{W} is precisely the class of trivial objects (that is, objects weakly equivalent to zero). A *hereditary Hovey triple* is a Hovey triple $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ for which the associated complete cotorsion pairs $(\mathcal{C} \cap \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}, \mathcal{W} \cap \mathcal{F})$ are both hereditary (as defined in 2.3).

Gillespie extends in [17, Thm. 3.3] Hovey’s correspondance, mentioned above, from the realm of abelian categories to the realm of weakly idempotent complete exact categories. More precisely, if \mathcal{A} is just an exact category (not necessarily abelian), then an *exact model structure* on \mathcal{A} is a model structure on \mathcal{A} which is compatible with the exact structure in the sense of [17, Def. 3.1]. If, in addition, \mathcal{A} is weakly idempotent complete ([17, Def. 2.2]), then exact model structures on \mathcal{A} correspond precisely to Hovey triples $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ in \mathcal{A} .

Recall from [24, Cor. 1.2.7 and Thm. 1.2.10(i)] that if \mathcal{C} is any model category, then the inclusion $\mathcal{C}_{\text{cf}} \rightarrow \mathcal{C}$ induces an equivalence $\mathcal{C}_{\text{cf}}/\sim \rightarrow \text{Ho}(\mathcal{C})$. Here \mathcal{C}_{cf} is the full subcategory of \mathcal{C} whose objects are both cofibrant and fibrant, “ \sim ” is the (abstract) homotopy relation from [24, Def. 1.2.4], and $\text{Ho}(\mathcal{C})$ is the homotopy category of the model category \mathcal{C} (that is, the localization of \mathcal{C} with respect to the collection of weak equivalences).

2.5. Let \mathcal{A} be a weakly idempotent complete exact category equipped with an exact model structure coming from a *hereditary Hovey triple* $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ in \mathcal{A} . As explained in 2.4, one has $\mathcal{A}_{\text{cf}} = \mathcal{C} \cap \mathcal{F}$, which by [17, Prop. 5.2(4)]/[32, Thm. 6.21(1)] is a Frobenius category with $\mathcal{C} \cap \mathcal{W} \cap \mathcal{F}$ as the class of projective-injective objects. By [17, Prop. 4.4(5)]/[32, Lem.

⁴Recall that a class \mathcal{W} in an abelian (or, more generally, in an exact) category \mathcal{A} is *thick* if it is closed under retracts and satisfies that whenever two out of three terms in a short exact sequence are in \mathcal{W} , then so is the third.

6.16(3)] two parallel morphisms in $\mathcal{A}_{\text{cf}} = \mathcal{C} \cap \mathcal{F}$ are homotopic, in the (abstract) model categorical sense, if and only their difference factors through an object in $\mathcal{C} \cap \mathcal{W} \cap \mathcal{F}$. Thus, $\mathcal{A}_{\text{cf}}/\sim$ is nothing but the stable category $\underline{\mathcal{A}}_{\text{cf}}$ of the Frobenius category \mathcal{A}_{cf} (see the remarks preceding Proposition 2.2), so the category $\mathcal{A}_{\text{cf}}/\sim$ carries a natural triangulated structure. As mentioned above, one has an equivalence of categories $\text{Ho}(\mathcal{A}) \simeq \mathcal{A}_{\text{cf}}/\sim$, and via this equivalence the homotopy category $\text{Ho}(\mathcal{A})$ inherits a triangulated structure from $\mathcal{A}_{\text{cf}}/\sim$. More precisely, the distinguished triangles in $\text{Ho}(\mathcal{A})$ are, up to isomorphism, the images in $\text{Ho}(\mathcal{A})$ of distinguished triangles in $\underline{\mathcal{A}}_{\text{cf}} = \mathcal{A}_{\text{cf}}/\sim$ under the equivalence $\mathcal{A}_{\text{cf}}/\sim \rightarrow \text{Ho}(\mathcal{A})$. It is evident that when $\text{Ho}(\mathcal{A})$ is equipped with this triangulated structure, then the equivalence $\text{Ho}(\mathcal{A}) \simeq \mathcal{A}_{\text{cf}}/\sim$ (of ordinary categories *a priori*) becomes an equivalence of triangulated categories, that is, the functors $\text{Ho}(\mathcal{A}) \rightleftarrows \mathcal{A}_{\text{cf}}/\sim$ are triangulated.

3. SHARP–FOXBY ADJUNCTIONS

Recall from the beginning of Section 2 that \mathcal{A} always denotes any bicomplete abelian category with enough projectives and enough injectives. In this section, we give conditions on \mathcal{A} which ensure that $\underline{\text{GProj}}(\mathcal{A})$ and $\underline{\text{GInj}}(\mathcal{A})$ are equivalent as triangulated categories.

Definition 3.1. Let \mathcal{U}^π be the full subcategory of \mathcal{A} whose objects are given by

$$\mathcal{U}^\pi = \{M \in \mathcal{A} \mid \text{Gpd}_{\mathcal{A}}(M) < \infty\}.$$

Let and \mathcal{C}^π , \mathcal{W}^π , and \mathcal{F}^π be the following subclasses of \mathcal{U}^π :

$$\mathcal{C}^\pi = \text{GProj}(\mathcal{A}), \quad \mathcal{W}^\pi = \{M \in \mathcal{A} \mid \text{pd}_{\mathcal{A}}(M) < \infty\}, \quad \text{and} \quad \mathcal{F}^\pi = \mathcal{U}^\pi.$$

The classes \mathcal{U}^π , \mathcal{C}^π , \mathcal{W}^π , and \mathcal{F}^π depend on \mathcal{A} , and if necessary we use the more detailed notation $\mathcal{U}_{\mathcal{A}}^\pi$, $\mathcal{C}_{\mathcal{A}}^\pi$, $\mathcal{W}_{\mathcal{A}}^\pi$, and $\mathcal{F}_{\mathcal{A}}^\pi$ instead. (The superscript “ π ” is supposed to give the reader associations to the word “projective”.)

Definition 3.2. Let \mathcal{U}^i be the full subcategory of \mathcal{A} whose objects are given by

$$\mathcal{U}^i = \{N \in \mathcal{A} \mid \text{Gid}_{\mathcal{A}}(N) < \infty\}.$$

Let and \mathcal{C}^i , \mathcal{W}^i , and \mathcal{F}^i be the following subclasses of \mathcal{U}^i :

$$\mathcal{C}^i = \mathcal{U}^i, \quad \mathcal{W}^i = \{N \in \mathcal{A} \mid \text{id}_{\mathcal{A}}(N) < \infty\}, \quad \text{and} \quad \mathcal{F}^i = \text{GInj}(\mathcal{A}).$$

The classes \mathcal{U}^i , \mathcal{C}^i , \mathcal{W}^i , and \mathcal{F}^i depend on \mathcal{A} , and if necessary we use the more detailed notation $\mathcal{U}_{\mathcal{A}}^i$, $\mathcal{C}_{\mathcal{A}}^i$, $\mathcal{W}_{\mathcal{A}}^i$, and $\mathcal{F}_{\mathcal{A}}^i$ instead. (The superscript “ i ” is supposed to give the reader associations to the word “injective”.)

Lemma 3.3. *The categories \mathcal{U}^π and \mathcal{U}^i are additive and extension-closed subcategories of the abelian category \mathcal{A} ; hence they are exact categories. Furthermore, \mathcal{U}^π and \mathcal{U}^i are closed under direct summands in \mathcal{A} ; hence they are idempotent complete.*

Proof. In the case where $\mathcal{A} = \text{Mod}(A)$ for a ring A , the assertions follow from [23, Prop. 2.19 and Thm. 2.24] (and the dual statements about Gorenstein injective modules). By inspection, one verifies that the same proofs work in any bicomplete abelian category \mathcal{A} with enough projectives and enough injectives. \square

We show in Theorems 3.7 and 3.9 that $(\mathcal{C}^\pi, \mathcal{W}^\pi, \mathcal{F}^\pi)$ and $(\mathcal{C}^i, \mathcal{W}^i, \mathcal{F}^i)$ are Hovey triples (see 2.4) in the idempotent complete exact categories \mathcal{U}^π and \mathcal{U}^i .

Definition 3.4. A *Sharp–Foxby adjunction* on \mathcal{A} is an adjunction (S, T) of endofunctors on \mathcal{A} for which the following properties hold:

(SF1) S maps \mathcal{U}^π to \mathcal{U}^i and it maps \mathcal{W}^π to \mathcal{W}^i .

- (SF2) The restriction of S to \mathcal{U}^π is exact: if $0 \rightarrow X' \rightarrow X \rightarrow X'' \rightarrow 0$ is an exact sequence in \mathcal{A} with $X', X, X'' \in \mathcal{U}^\pi$, then the sequence $0 \rightarrow SX' \rightarrow SX \rightarrow SX'' \rightarrow 0$ is exact.
- (SF3) T maps \mathcal{U}^t to \mathcal{U}^π and it maps \mathcal{W}^t to \mathcal{W}^π .
- (SF4) The restriction of T to \mathcal{U}^t is exact: if $0 \rightarrow Y' \rightarrow Y \rightarrow Y'' \rightarrow 0$ is an exact sequence in \mathcal{A} with $Y', Y, Y'' \in \mathcal{U}^t$, then the sequence $0 \rightarrow TY' \rightarrow TY \rightarrow TY'' \rightarrow 0$ is exact.
- (SF5) The unit of adjunction $\eta_X: X \rightarrow TSX$ is an isomorphism for every $X \in \mathcal{U}^\pi$.
- (SF6) The counit of adjunction $\varepsilon_Y: STY \rightarrow Y$ is an isomorphism for every $Y \in \mathcal{U}^t$.

Remark 3.5. By (SF1), (SF3), (SF5), and (SF6) a Sharp–Foxby adjunction $S: \mathcal{A} \rightleftarrows \mathcal{A}: T$ restricts to adjoint equivalences of categories $\mathcal{U}^\pi \rightleftarrows \mathcal{U}^t$ and $\mathcal{W}^\pi \rightleftarrows \mathcal{W}^t$. By Lemma 3.3 the categories \mathcal{U}^π and \mathcal{U}^t have natural exact structures. Conditions (SF2) and (SF4) imply that the induced adjoint equivalence $\mathcal{U}^\pi \rightleftarrows \mathcal{U}^t$ preserves the exact structure, i.e. the functors are exact; thus it is an adjoint equivalence of exact categories.⁵

The following example explains the terminology in Definition 3.4.

Example 3.6. Let A be a commutative noetherian local Cohen–Macaulay ring with a dualizing module D . Foxby considered in [14, §1] two classes $\mathbf{A}(A)$ and $\mathbf{B}(A)$ of A -modules⁶:

A module M is in $\mathbf{A}(A)$ if and only if $\mathrm{Tor}_i^A(D, M) = 0$ and $\mathrm{Ext}_A^i(D, D \otimes_A M) = 0$ for all $i > 0$ and the natural homomorphism $\eta_M: M \rightarrow \mathrm{Hom}_A(D, D \otimes_A M)$ is an isomorphism.

A module N is in $\mathbf{B}(A)$ if and only if $\mathrm{Ext}_A^i(D, N) = 0$ and $\mathrm{Tor}_i^A(D, \mathrm{Hom}_A(D, N)) = 0$ for all $i > 0$ and the natural homomorphism $\varepsilon_N: D \otimes_A \mathrm{Hom}_A(D, N) \rightarrow N$ is an isomorphism.

Foxby [14] proved that the adjunction $(D \otimes_A -, \mathrm{Hom}_A(D, -))$ on $\mathrm{Mod}(A)$ restricts to an adjoint equivalence $\mathbf{A}(A) \rightleftarrows \mathbf{B}(A)$ and further to an adjoint equivalence $\mathcal{W}_{\mathrm{Mod}(A)}^\pi \rightleftarrows \mathcal{W}_{\mathrm{Mod}(A)}^t$ (see Definitions 3.1 and 3.2). The latter is an extension of a result [31, Thm. (2.9)] by Sharp, which asserts that $D \otimes_A -$ and $\mathrm{Hom}_A(D, -)$ restrict to an adjoint equivalence between the categories of *finitely generated* A -modules with finite projective dimension and *finitely generated* A -modules with finite injective dimension. Note that it is evident from the definitions that the restriction of $D \otimes_A -$ to $\mathbf{A}(A)$ and of $\mathrm{Hom}_A(D, -)$ to $\mathbf{B}(A)$ are exact functors.

By Enochs, Jenda, and Xu [12, Cor. 2.4 and 2.6] an A -module belongs to $\mathbf{A}(A)$, respectively, $\mathbf{B}(A)$, if and only if it has finite Gorenstein projective dimension, respectively, finite Gorenstein injective dimension. Thus, in the notation from 3.1 and 3.2 we have:

$$\mathbf{A}(A) = \mathcal{U}_{\mathrm{Mod}(A)}^\pi \quad \text{and} \quad \mathbf{B}(A) = \mathcal{U}_{\mathrm{Mod}(A)}^t.$$

Consequently, $(S, T) = (D \otimes_A -, \mathrm{Hom}_A(D, -))$ is a Sharp–Foxby adjunction on $\mathrm{Mod}(A)$. In view of [9, Thms. 4.1 and 4.4] this remains to be true if A is any two-sided noetherian ring with a dualizing module D , that is, a dualizing complex concentrated in degree zero.

Theorem 3.7. *Consider the idempotent complete exact category \mathcal{U}^π from Lemma 3.3. The triple $(\mathcal{C}^\pi, \mathcal{W}^\pi, \mathcal{F}^\pi)$ from Definition 3.1 is a hereditary Hovey triple in \mathcal{U}^π (see 2.4). In particular, \mathcal{U}^π has an exact model structure for which:*

⁵ If \mathcal{E} and \mathcal{E}' are exact categories and $F: \mathcal{E} \rightleftarrows \mathcal{E}': G$ is an adjoint equivalence of the underlying (ordinary) categories, then it does not automatically follow that the functors F and G are exact. Indeed, if \mathcal{E} and \mathcal{E}' have the same underlying category and the exact structure on \mathcal{E} is coarser than that on \mathcal{E}' (that is, every sequence which is exact in \mathcal{E} is also exact in \mathcal{E}' — for example, \mathcal{E} could have the trivial exact structure, in which the only “exact” sequences are the split exact ones, whereas \mathcal{E}' could have any exact structure), then the identity functors $\mathcal{E} \rightleftarrows \mathcal{E}'$ constitute an adjoint equivalence of the underlying categories where only $\mathcal{E} \rightarrow \mathcal{E}'$ is exact (but $\mathcal{E} \leftarrow \mathcal{E}'$ is not).

⁶ In the literature, the classes $\mathbf{A}(A)$ and $\mathbf{B}(A)$ are referred to as *Foxby classes*. Sometimes, $\mathbf{A}(A)$ is called the *Auslander class* and $\mathbf{B}(A)$ is called the *Bass class*. Foxby himself [14] used the symbols Φ_D and Ψ_D for these classes, but in the paper [12] by Enochs, Jenda, and Xu they are denoted by \mathcal{G}_0 and \mathcal{J}_0 . We have adopted the symbols $\mathbf{A}(A)$ and $\mathbf{B}(A)$ from the joint work of Avramov and Foxby; see for example [1, §3].

- The cofibrant objects in \mathcal{U}^π are the Gorenstein projective objects in \mathcal{A} .
- The trivial objects in \mathcal{U}^π are the objects in \mathcal{A} with finite projective dimension.
- All objects in \mathcal{U}^π are fibrant.

The homotopy category of this model category is equivalent, as a triangulated category, to the stable category of Gorenstein projective objects in \mathcal{A} ; in symbols:

$$\mathrm{Ho}(\mathcal{U}^\pi) \simeq \underline{\mathrm{GProj}}(\mathcal{A}).$$

Remark 3.8. A number of fundamental properties of Gorenstein projective modules, i.e. Gorenstein projective objects in the category $\mathcal{A} = \mathrm{Mod}(A)$ where A is a ring, are recorded in e.g. [9, 23]. The results we need about Gorenstein projective objects in a general abelian category (still bicomplete with enough projectives and enough injectives) can be proved as it is done for modules. We leave it to the reader to inspect the relevant proofs.

Proof of Theorem 3.7. It is well-known that \mathcal{W}^π is a thick subcategory of \mathcal{A} (and hence also of \mathcal{U}^π). By [23, Prop. 2.27] the intersection $\mathcal{C}^\pi \cap \mathcal{W}^\pi$ equals the class $\mathrm{Proj} \mathcal{A}$ of projective objects in \mathcal{A} . Thus the pair $(\mathcal{C}^\pi \cap \mathcal{W}^\pi, \mathcal{F}^\pi)$ is equal to $(\mathrm{Proj} \mathcal{A}, \mathcal{U}^\pi)$, which we now argue is a complete hereditary cotorsion pair in \mathcal{U}^π . As $\mathrm{Ext}_{\mathcal{A}}^{\geq 1}(P, A) = 0$ for all $P \in \mathrm{Proj} \mathcal{A}$ and all $A \in \mathcal{U}^\pi$ (even all $A \in \mathcal{A}$), we get that $(\mathrm{Proj} \mathcal{A})^\perp = \mathcal{U}^\pi$ (as the “ \perp ” is only calculated inside of \mathcal{U}^π) and that $\mathrm{Proj} \mathcal{A} \subseteq {}^\perp \mathcal{U}^\pi$. To show that $\mathrm{Proj} \mathcal{A} \supseteq {}^\perp \mathcal{U}^\pi$ let $M \in {}^\perp \mathcal{U}^\pi$ ($\subseteq \mathcal{U}^\pi$). By assumption, \mathcal{A} has enough projectives, and hence there exists a short exact sequence in \mathcal{A} ,

$$(\#1) \quad 0 \longrightarrow A \longrightarrow P \longrightarrow M \longrightarrow 0,$$

where P is projective. As M belongs to \mathcal{U}^π , so does A by [23, Thm. 2.24]. By assumption, $\mathrm{Ext}_{\mathcal{A}}^1(M, A) = 0$, so (#1) splits and hence $M \in \mathrm{Proj} \mathcal{A}$. This shows that $(\mathrm{Proj} \mathcal{A}, \mathcal{U}^\pi)$ is a hereditary cotorsion pair. For completeness of this cotorsion pair, the sequence (#1) shows that the pair has enough projectives. The trivial exact sequence $0 \rightarrow M \rightarrow M \rightarrow 0 \rightarrow 0$ (for any M in \mathcal{U}^π) shows that the pair has enough injectives.

Next we show that $(\mathcal{C}^\pi, \mathcal{W}^\pi \cap \mathcal{F}^\pi) = (\mathrm{GProj} \mathcal{A}, \mathcal{W}^\pi)$ is a complete hereditary cotorsion pair in \mathcal{U}^π . By [23, Thm. 2.20] we have $\mathrm{Ext}_{\mathcal{A}}^{\geq 1}(G, A) = 0$ for all $G \in \mathrm{GProj} \mathcal{A}$ and $A \in \mathcal{W}^\pi$, and hence we get $\mathrm{GProj} \mathcal{A} \subseteq {}^\perp \mathcal{W}^\pi$ and $(\mathrm{GProj} \mathcal{A})^\perp \supseteq \mathcal{W}^\pi$. To show that $\mathrm{GProj} \mathcal{A} \supseteq {}^\perp \mathcal{W}^\pi$, let $M \in {}^\perp \mathcal{W}^\pi$ ($\subseteq \mathcal{U}^\pi$). By [23, Thm 2.10] there exists a short exact sequence

$$(\#2) \quad 0 \longrightarrow A \longrightarrow G \longrightarrow M \longrightarrow 0$$

with $G \in \mathrm{GProj} \mathcal{A}$ and $A \in \mathcal{W}^\pi$. By assumption, $\mathrm{Ext}_{\mathcal{A}}^1(M, A) = 0$, so (#2) splits and hence M is a direct summand in G . By [23, Thm 2.5] (see also Prop. 1.4 in *loc. cit.*) the class $\mathrm{GProj} \mathcal{A}$ is closed under direct summands (here we use our assumption that \mathcal{A} is cocomplete, or at least that \mathcal{A} has countable coproducts), and it follows that M itself belongs to $\mathrm{GProj} \mathcal{A}$. To show $(\mathrm{GProj} \mathcal{A})^\perp \subseteq \mathcal{W}^\pi$, assume that $M \in (\mathrm{GProj} \mathcal{A})^\perp$ ($\subseteq \mathcal{U}^\pi$). By [9, Lem. 2.17] there is a short exact sequence

$$(\#3) \quad 0 \longrightarrow M \longrightarrow A' \longrightarrow G' \longrightarrow 0$$

where $G' \in \mathrm{GProj} \mathcal{A}$ and $\mathrm{pd}_{\mathcal{A}}(A') = \mathrm{Gpd}_{\mathcal{A}}(M) < \infty$, that is, A' is in \mathcal{W}^π . By assumption, $\mathrm{Ext}_{\mathcal{A}}^1(G', M) = 0$, so (#3) splits and hence M also belongs to \mathcal{W}^π (which is thick). Thus $(\mathrm{GProj} \mathcal{A}, \mathcal{W}^\pi)$ is a hereditary cotorsion pair in \mathcal{U}^π , and the existence of the sequences (#2) and (#3) shows that this cotorsion pair is complete.

These arguments prove that $(\mathcal{C}^\pi, \mathcal{W}^\pi, \mathcal{F}^\pi)$ is a hereditary Hovey triple in \mathcal{U}^π . In view of the equalities $\mathcal{C}^\pi \cap \mathcal{F}^\pi = \mathrm{GProj} \mathcal{A}$ and $\mathcal{C}^\pi \cap \mathcal{W}^\pi \cap \mathcal{F}^\pi = \mathrm{Proj} \mathcal{A}$, where the latter is by [23, Prop 2.27], the rest of the theorem now follows from 2.4 and 2.5 (and Proposition 2.2). \square

Theorem 3.9. *Consider the idempotent complete exact category \mathcal{U}^t from Lemma 3.3. The triple $(\mathcal{C}^t, \mathcal{W}^t, \mathcal{F}^t)$ from Definition 3.2 is a hereditary Hovey triple in \mathcal{U}^t (see 2.4). In particular, \mathcal{U}^t has an exact model structure for which:*

- All objects in \mathcal{U}^t are cofibrant.
- The trivial objects in \mathcal{U}^t are the objects in \mathcal{A} with finite injective dimension.
- The fibrant objects in \mathcal{U}^t are the Gorenstein injective objects in \mathcal{A} .

The homotopy category of this model category is equivalent, as a triangulated category, to the stable category of Gorenstein injective objects in \mathcal{A} ; in symbols:

$$\mathrm{Ho}(\mathcal{U}^t) \simeq \underline{\mathrm{GInj}}(\mathcal{A}).$$

Proof. Dual to the proof of Theorem 3.1. □

Our next goal is to show that a Sharp–Foxby adjunction on \mathcal{A} induces a Quillen equivalence between the model categories \mathcal{U}^π and \mathcal{U}^t . To this end, the next result will be useful.

Proposition 3.10. *Let \mathcal{M} and \mathcal{M}' be two weakly idempotent complete exact model categories with associated Hovey triples $(\mathcal{C}, \mathcal{W}, \mathcal{F})$ and $(\mathcal{C}', \mathcal{W}', \mathcal{F}')$; see 2.4. Assume that (F, G) is a Quillen adjunction $\mathcal{M} \rightleftarrows \mathcal{M}'$ where the functors F and G are exact and satisfy $F(\mathcal{W}) \subseteq \mathcal{W}'$ and $G(\mathcal{W}') \subseteq \mathcal{W}$. Then (F, G) is a Quillen equivalence if and only if the unit $\eta_X: X \rightarrow GFX$ is a weak equivalence for every $X \in \mathcal{C}$ and the counit $\varepsilon_Y: FGY \rightarrow Y$ is a weak equivalence for every $Y \in \mathcal{F}'$.*

Proof. Write Q for the cofibrant replacement functor in \mathcal{M} and $q_X: QX \rightarrow X$ for the natural trivial fibration ($X \in \mathcal{M}$). Similarly, write R for the fibrant replacement functor in \mathcal{M}' and $r_Y: Y \rightarrow RY$ for the natural trivial cofibration ($Y \in \mathcal{M}'$). By [24, Prop. 1.3.13] we have that (F, G) is a Quillen equivalence if and only if the composite

$$X \xrightarrow{\eta_X} GFX \xrightarrow{Gr_{FX}} GRFX$$

is a weak equivalence for all $X \in \mathcal{C}$ and the composite

$$FQGY \xrightarrow{Fq_{GY}} FGY \xrightarrow{\varepsilon_Y} Y$$

is a weak equivalence for all $Y \in \mathcal{F}'$. We claim that the morphisms Gr_{FX} and Fq_{GY} are always weak equivalences for every $X \in \mathcal{M}$ and $Y \in \mathcal{M}'$ (which proves the assertion by the 2-out-of-3 property for weak equivalences). We only show that Gr_{FX} is a weak equivalence. The fact that $r_{FX}: FX \rightarrow RFX$ is a trivial cofibration means, by definition [17, Def. 3.1] of an exact model structure, that r_{FX} is an admissible monomorphism with a trivially cofibrant cokernel, that is, one has a conflation (a short exact sequence)

$$FX \xrightarrow{\simeq} r_{FX} \rightarrow RFX \xrightarrow{\pi} C$$

in \mathcal{M}' where C is trivially cofibrant, that is, $C \in \mathcal{C}' \cap \mathcal{W}'$ (and RFX is of course fibrant). By applying the exact functor G to the sequence above, we get a conflation in \mathcal{M} , which is the bottom row of the following pullback diagram:

$$\begin{array}{ccccc} GFX & \xrightarrow[\simeq]{\iota} & T & \xrightarrow{\varrho} & QGC \\ \parallel & & \downarrow \varphi & & \downarrow q_{GC} \\ GFX & \xrightarrow{Gr_{FX}} & GRFX & \xrightarrow{G\pi} & GC. \end{array}$$

Note that this pullback diagram really exists; indeed, by definition of an exact category, any pullback of an admissible epimorphism exists and admissible epimorphisms are stable

under pullbacks. In particular, ϱ is an admissible epimorphism (and ϱ has the same kernel as $G\pi$; cf. Freyd [15, Thm. 2.52]). Since $C \in \mathcal{W}'$ we have $GC \in \mathcal{W}$ by assumption. Since one always has $Q(\mathcal{W}) \subseteq \mathcal{W}$, it follows that $QGC \in \mathcal{W}$, and hence $QGC \in \mathcal{C} \cap \mathcal{W}$ (as QY is always cofibrant). This means that ι is a trivial cofibration. In any model category, the class of trivial fibrations is stable under pullbacks by [25, Cor. 1.1.11]; thus the fact that q_{GC} is a trivial fibration forces φ to be the same. As ι and φ are, in particular, weak equivalences, so is their composite $Gr_{FX} = \varphi \circ \iota$, as desired. \square

Theorem 3.11. *A Sharp–Foxby adjunction (S, T) on \mathcal{A} induces a Quillen equivalence between the model categories \mathcal{U}^π and \mathcal{U}^ι constructed in Theorems 3.7 and 3.9. Thus the total (left/right) derived functors of S and T yield an adjoint equivalence of the corresponding homotopy categories,*

$$(\#4) \quad \mathrm{Ho}(\mathcal{U}^\pi) \begin{array}{c} \xrightarrow{\mathbf{LS}} \\ \xleftarrow{\mathbf{RT}} \end{array} \mathrm{Ho}(\mathcal{U}^\iota).$$

In fact, this is an equivalence of triangulated categories.

Proof. As mentioned Remark 3.5, a Sharp–Foxby adjunction (S, T) on \mathcal{A} induces an exact adjoint equivalence between \mathcal{U}^π and \mathcal{U}^ι with $S(\mathcal{W}^\pi) \subseteq \mathcal{W}^\iota$ and $T(\mathcal{W}^\iota) \subseteq \mathcal{W}^\pi$. Hence the unit $\eta_X: X \rightarrow T S X$ is an isomorphism, and hence also a weak equivalence, for all $X \in \mathcal{U}^\pi$ (in particular for $X \in \mathcal{C}^\pi$); and the counit $\varepsilon_Y: S T Y \rightarrow Y$ is an isomorphism, and hence also a weak equivalence, for all $Y \in \mathcal{U}^\iota$ (in particular for $Y \in \mathcal{F}^\iota$). Thus, if we can show that (S, T) is a Quillen adjunction $\mathcal{U}^\pi \rightleftarrows \mathcal{U}^\iota$, then Proposition 3.10 will imply that it is in fact a Quillen equivalence (as claimed). To show this, it must be argued that $S: \mathcal{U}^\pi \rightarrow \mathcal{U}^\iota$ is a left Quillen functor (see [24, Def. 1.3.1]), that is, we must argue that S maps (trivial) cofibrations in \mathcal{U}^π to (trivial) cofibrations in \mathcal{U}^ι . Let f be a (trivial) cofibration in \mathcal{U}^π , that is, f is an admissible monomorphism with a (trivially) cofibrant cokernel C (see [17, Def. 3.1]). Since S is exact, it follows that Sf is an admissible monomorphism in \mathcal{U}^ι with cokernel SC . Hence, we only need to prove that S maps (trivially) cofibrant objects in \mathcal{U}^π to (trivially) cofibrant objects in \mathcal{U}^ι . However, this is clear as every object in \mathcal{U}^ι is cofibrant, see Theorem 3.9, and since we have $S(\mathcal{W}^\pi) \subseteq \mathcal{W}^\iota$.

Having established that (S, T) yields a Quillen equivalence $\mathcal{U}^\pi \rightleftarrows \mathcal{U}^\iota$, the adjoint equivalence of homotopy categories displayed in (#4) follows from [24, Prop. 1.3.13].

It remains to see that the functors \mathbf{LS} and \mathbf{RT} are triangulated. By [28, Lem. 5.3.6] it suffices to prove that \mathbf{LS} is triangulated, because then its right adjoint \mathbf{RT} will automatically be triangulated as well. Recall from 2.5 that the distinguished triangles in $\mathrm{Ho}(\mathcal{U}^\pi)$ are, up to isomorphism, the images in $\mathrm{Ho}(\mathcal{U}^\pi)$ of distinguished triangles in $\underline{\mathrm{GProj}}(\mathcal{A})$ under the equivalence $\underline{\mathrm{GProj}}(\mathcal{A}) \rightarrow \mathrm{Ho}(\mathcal{U}^\pi)$ (see also Theorem 3.7).

At this point we need to recall from [21, Chap. I§2.5] how the triangulated structure on the stable category $\underline{\mathrm{GProj}}(\mathcal{A})$ is defined. For every morphism $u: G \rightarrow G'$ in the Frobenius category $\mathrm{GProj}(\mathcal{A})$ choose a short exact sequence (a conflation) $G \xrightarrow{i} P \xrightarrow{p} \tilde{G}$ in $\mathrm{GProj}(\mathcal{A})$ where P is a projective-injective object, that is, $P \in \mathrm{Proj}(\mathcal{A})$. The object \tilde{G} is the suspension of G ; in symbols, $\tilde{G} = \Sigma G$ (the assignment $G \mapsto \tilde{G} = \Sigma G$ is not functorial on $\mathrm{GProj}(\mathcal{A})$, but it is functorial on $\underline{\mathrm{GProj}}(\mathcal{A})$). Then consider the pushout diagram in $\mathrm{GProj}(\mathcal{A})$,

$$(\#5) \quad \begin{array}{ccccc} G & \xrightarrow{i} & P & \xrightarrow{p} & \tilde{G} \\ \downarrow u & \text{pushout} & \downarrow \iota & & \parallel \\ G' & \xrightarrow{v} & G'' & \xrightarrow{w} & \tilde{G} \end{array}$$

The diagram

$$(\#6) \quad G \xrightarrow{u} G' \xrightarrow{v} G'' \xrightarrow{w} \tilde{G},$$

considered as a diagram in $\underline{\mathbf{GProj}}(\mathcal{A})$, is called a *standard triangle*. By definition, a distinguished triangle in $\underline{\mathbf{GProj}}(\mathcal{A})$ is a diagram in this category which is isomorphic to some standard triangle. The triangulated structure on $\underline{\mathbf{GInj}}(\mathcal{A})$ is defined similarly.

We must show that the functor \mathbf{LS} maps every distinguished triangle Δ in $\mathbf{Ho}(\mathcal{U}^\pi)$ to a distinguished triangle in $\mathbf{Ho}(\mathcal{U}^t)$. By the considerations above, we may assume that Δ is the image in $\mathbf{Ho}(\mathcal{U}^t)$ of a standard triangle (#6) in $\underline{\mathbf{GProj}}(\mathcal{A})$. By definition, see [24, Def. 1.3.6], the action of the functor \mathbf{LS} on an object X in $\mathbf{Ho}(\mathcal{U}^\pi)$ is $\mathbf{LS}(X) = SQX$ where QX is a cofibrant replacement of X . As the objects in (#6) are already cofibrant in \mathcal{U}^π , see Theorem 3.7, the diagram $\mathbf{LS}(\Delta)$ is nothing but

$$(\#7) \quad SG \xrightarrow{Su} SG' \xrightarrow{Sv} SG'' \xrightarrow{Sw} S\tilde{G},$$

which we must show is a distinguished triangle in $\mathbf{Ho}(\mathcal{U}^t)$. Since the pair $(\mathcal{C}^t \cap \mathcal{W}^t, \mathcal{F}^t) = (\mathcal{W}^t, \mathbf{GInj}\mathcal{A})$ is a hereditary cotorsion pair in \mathcal{U}^t , see Theorem 3.9 and Definition 3.2, it follows from [32, Lem. 6.20] that we can find a diagram in \mathcal{U}^t ,

$$(\#8) \quad \begin{array}{ccccc} SG & \xrightarrow{Si} & SP & \xrightarrow{Sp} & S\tilde{G} \\ \downarrow h & & \downarrow e & & \downarrow \tilde{h} \\ H & \xrightarrow{i_0} & E & \xrightarrow{p_0} & \tilde{H} \\ \downarrow & & \downarrow & & \downarrow \\ J & \longrightarrow & I & \longrightarrow & \tilde{J} \end{array}$$

whose rows and columns are conflations, where H, E, \tilde{H} are Gorenstein injective, and where J, I, \tilde{J} have finite injective dimension. As $P \in \mathbf{Proj}\mathcal{A} \subseteq \mathcal{W}^\pi$ we have $SP \in \mathcal{W}^t$, that is, SP has finite injective dimension. It follows from the middle column in (#8) that E has finite injective dimension, and since E is also Gorenstein injective it must be injective (this is immediate from the definition, 2.1, of Gorenstein injective objects). Let $SG' \xrightarrow{h'} H' \rightarrow J'$ be a short exact sequence with $H' \in \mathbf{GInj}\mathcal{A}$ and $J' \in \mathcal{W}^t$. The morphism $h: SG \rightarrow H$ is a (special) Gorenstein injective preenvelope of SG since it is monic and its cokernel $J \in \mathcal{W}^t$ satisfies $\mathrm{Ext}_{\mathcal{A}}^1(J, X) = 0$ for all $X \in \mathbf{GInj}(\mathcal{A})$; see [34, Prop. 2.1.4]. Thus, the morphism $h'Su: SG \rightarrow H' \in \mathbf{GInj}(\mathcal{A})$ lifts to a morphism $u_0: H \rightarrow H'$ such that $u_0h = h'Su$. This gives commutativity of the left wall in the following diagram:

$$(\#9) \quad \begin{array}{ccccccc} & & H & \xrightarrow{i_0} & E & \xrightarrow{p_0} & \tilde{H} \\ & \nearrow h & \downarrow & \nearrow e & \downarrow & \nearrow \tilde{h} & \parallel \\ SG & \xrightarrow{Si} & SP & \xrightarrow{Sp} & S\tilde{G} & & \\ \downarrow Su & \downarrow u_0 & \downarrow & \downarrow t_0 & \downarrow & \downarrow & \parallel \\ & \nearrow h' & H' & \xrightarrow{v_0} & H'' & \xrightarrow{w_0} & \tilde{H} \\ & \downarrow h' & \downarrow St & \downarrow h'' & \downarrow & \downarrow \tilde{h} & \parallel \\ SG' & \xrightarrow{Sv} & SG'' & \xrightarrow{Sw} & S\tilde{G} & & \end{array}$$

The top wall in (#9) is just the upper half of the commutative diagram (#8). The back wall is the (commutative) pushout diagram of the morphisms $H' \xleftarrow{u_0} H \xrightarrow{i_0} E$. The right wall is evidently commutative. The front wall in (#9) is obtained by applying the exact functor S to the diagram (#5). Since S is a left adjoint functor, it preserves colimits, so the front wall

in (#9) is (still) a pushout diagram. As $(v_0h')Su = v_0u_0h = t_0i_0h = (t_0e)Si$ and since SG'' is the pushout of $SG' \xleftarrow{Su} SG \xrightarrow{Si} SP$, there exists a (unique) morphism $h'' : SG'' \rightarrow H''$ such that $h''Sv = v_0h'$ and $h''St = t_0e$. The first of these identities show that the left square in the bottom wall in (#9) is commutative. It follows from the universal property of the pushout SG'' that the right square in the bottom wall is commutative as well. By applying the Snake Lemma to this bottom wall, we see that h'' is monic (as h' and \tilde{h} are so) and that the cokernel J'' of h'' sits in a short exact sequence $0 \rightarrow J' \rightarrow J'' \rightarrow \tilde{J} \rightarrow 0$. Since $J', \tilde{J} \in \mathcal{W}^t$ it follows that $J'' \in \mathcal{W}^t$. Since h, h', h'' , and \tilde{h} are (admissible) monomorphisms in \mathcal{U}^t whose cokernels belong to \mathcal{W}^t (which are the trivally cofibrant objects in \mathcal{U}^t), they are trivial cofibrations in the exact model structure on \mathcal{U}^t ; see [17, Def. 3.1]. In particular, h, h', h'' , and \tilde{h} are weak equivalences in \mathcal{U}^t and therefore isomorphisms in $\text{Ho}(\mathcal{U}^t)$. The commutative diagram (#9) now shows that in the homotopy category $\text{Ho}(\mathcal{U}^t)$, the diagram (#7) is isomorphic to

$$(10) \quad H \xrightarrow{u_0} H' \xrightarrow{v_0} H'' \xrightarrow{w_0} \tilde{H}.$$

By definition, and by commutativity of the back wall in (#9), the diagram (#10) is a standard triangle in $\underline{\text{GInj}}(\mathcal{A})$, and consequently, (#7) is a distinguished triangle in $\text{Ho}(\mathcal{U}^t)$. \square

Corollary 3.12. *If there exists a Sharp–Foxby adjunction (S, T) on \mathcal{A} , then there is an equivalence of triangulated categories, $\underline{\text{GProj}}(\mathcal{A}) \simeq \underline{\text{GInj}}(\mathcal{A})$.*

Proof. By Theorems 3.7, 3.11, and 3.9 there are the following equivalences of triangulated categories, $\underline{\text{GProj}}(\mathcal{A}) \simeq \text{Ho}(\mathcal{U}^t) \simeq \text{Ho}(\mathcal{U}^t) \simeq \underline{\text{GInj}}(\mathcal{A})$. \square

Remark 3.13. Before closing this section, we record a biproduct of Proposition 3.10 concerning virtually Gorenstein rings, which should be well known. We recall from [3, 4] that an Artin algebra A is called virtually Gorenstein if $(\text{GProj}(A))^\perp = {}^\perp(\text{GInj}(A))$. The same notion for commutative rings has also been studied in [36]. In what follows, assume that A is an Artin algebra or a commutative noetherian ring with finite Krull dimension. In both cases, it is well known [4, 19, 26] that there are Hovey triples

$$(\text{GProj}(A), (\text{GProj}(A))^\perp, \text{Mod}(A)) \quad \text{and} \quad (\text{Mod}(A), {}^\perp(\text{GInj}(A)), \text{GInj}(A)).$$

Applying Proposition 3.10 in the case where $F = G = I_{\text{Mod}(A)}$, we obtain that virtually Gorensteiness of A implies that the identity is a Quillen equivalence between the two model structures. Therefore the homotopy categories of these two models are, in fact, isomorphic. In case A is, in addition, commutative Gorenstein we recover the analogous statement for Gorenstein rings (see the comments after Theorem 8.6 in [25]).

4. THE CASE OF CHAIN COMPLEXES

Recall from the beginning of Section 2 that \mathcal{A} always denotes any bicomplete abelian category with enough projectives and enough injectives. In this section, we consider the abelian category $\text{Ch}(\mathcal{A})$ of unbounded chain complexes in \mathcal{A} and prove that, under suitable conditions, a Sharp–Foxby adjunction (S, T) on \mathcal{A} induces a Sharp–Foxby adjunction on $\text{Ch}(\mathcal{A})$ by degreewise application of the functors S and T . First we recall the following.

4.1. The *finitistic projective dimension*, $\text{FPD}(\mathcal{A})$, of \mathcal{A} is defined as

$$\text{FPD}(\mathcal{A}) = \sup\{\text{pd}_{\mathcal{A}}M \mid M \text{ is an object in } \mathcal{A} \text{ with finite projective dimension}\}.$$

Dually, the *finitistic injective dimension*, $\text{FID}(\mathcal{A})$, of \mathcal{A} is

$$\text{FID}(\mathcal{A}) = \sup\{\text{id}_{\mathcal{A}}M \mid M \text{ is an object in } \mathcal{A} \text{ with finite injective dimension}\}.$$

The *finitistic Gorenstein projective dimension*, $\text{FGPD}(\mathcal{A})$, and the *finitistic Gorenstein injective dimension*, $\text{FGID}(\mathcal{A})$, are defined similarly.

For most abelian categories that appear in applications, the finitistic dimensions defined above turn out to be finite. As in [23, (proofs of) Thms. 2.28 and 2.29] one easily proves:

Lemma 4.2. *There are equalities $\text{FGPD}(\mathcal{A}) = \text{FPD}(\mathcal{A})$ and $\text{FGID}(\mathcal{A}) = \text{FID}(\mathcal{A})$. Thus, if $\text{FPD}(\mathcal{A})$, respectively, $\text{FID}(\mathcal{A})$, is finite, then so is $\text{FGPD}(\mathcal{A})$, respectively, $\text{FGID}(\mathcal{A})$. \square*

In \mathcal{A} we have the subcategories $\mathcal{U}_\mathcal{A}^\pi$, $\mathcal{C}_\mathcal{A}^\pi$, $\mathcal{W}_\mathcal{A}^\pi$ and $\mathcal{F}_\mathcal{A}^\pi$ from Definition 3.1. Similarly, in $\mathcal{B} = \text{Ch}(\mathcal{A})$ we have the subcategories $\mathcal{U}_\mathcal{B}^\pi$, $\mathcal{C}_\mathcal{B}^\pi$, $\mathcal{W}_\mathcal{B}^\pi$ and $\mathcal{F}_\mathcal{B}^\pi$. The following result explains the relation between all these subcategories.

Proposition 4.3. *Assume that $\text{FPD}(\mathcal{A}) < \infty$ and let $X = \cdots \rightarrow X_{n+1} \rightarrow X_n \rightarrow X_{n-1} \rightarrow \cdots$ be an object in $\mathcal{B} := \text{Ch}(\mathcal{A})$. The following conclusions hold.*

- (i) X belongs to $\mathcal{U}_\mathcal{B}^\pi$ if and only if every X_n belongs to $\mathcal{U}_\mathcal{A}^\pi$.
- (ii) X belongs to $\mathcal{C}_\mathcal{B}^\pi$ if and only if every X_n belongs to $\mathcal{C}_\mathcal{A}^\pi$.
- (iii) X belongs to $\mathcal{W}_\mathcal{B}^\pi$ if and only if X is exact and every cycle $Z_n(X)$ belongs to $\mathcal{W}_\mathcal{A}^\pi$.
- (iv) X belongs to $\mathcal{F}_\mathcal{B}^\pi$ if and only if every X_n belongs to $\mathcal{F}_\mathcal{A}^\pi$.

Proof. Part (ii) is proved in [35, Thm. 2.2] in the case $\mathcal{A} = \text{Mod}(A)$ where A is any ring, but the proof works in any abelian category (with enough projectives).

In view of (ii), the “only if” part in (i) is clear. To prove the “if” part in (i), assume that every X_n is in $\mathcal{U}_\mathcal{A}^\pi$, that is, $\text{Gpd}_\mathcal{A}(X_n) < \infty$. By our assumption $\text{FPD}(\mathcal{A}) < \infty$ and by Lemma 4.2, it follows that $s = \sup\{\text{Gpd}_\mathcal{A}(X_n) \mid n \in \mathbb{Z}\}$ belongs to \mathbb{N}_0 . The proof is now by induction on s . If $s = 0$, then X is even in $\mathcal{C}_\mathcal{B}^\pi \subseteq \mathcal{U}_\mathcal{B}^\pi$ by part (ii). Now assume that $s > 0$. Choose any exact sequence

$$0 \longrightarrow K \longrightarrow P^{s-1} \longrightarrow \cdots \longrightarrow P^1 \longrightarrow P^0 \longrightarrow X \longrightarrow 0$$

in $\mathcal{B} = \text{Ch}(\mathcal{A})$ where P^0, \dots, P^{s-1} are complexes consisting of projective objects in \mathcal{A} . For each $n \in \mathbb{Z}$ we have an exact sequence $0 \rightarrow K_n \rightarrow P_n^{s-1} \rightarrow P_n^1 \rightarrow P_n^0 \rightarrow X_n \rightarrow 0$ in \mathcal{A} , and since P_n^0, \dots, P_n^{s-1} are projectives and $\text{Gpd}_\mathcal{A}(X_n) \leq s$, it follows that K_n is Gorenstein projective; cf. [23, (proof of) Prop. 2.7]. Thus, K is a complex of Gorenstein projective objects in \mathcal{A} , which by (ii) means that K is a Gorenstein projective object in $\mathcal{B} = \text{Ch}(\mathcal{A})$. So the exact sequence displayed above shows that $\text{Gpd}_\mathcal{B}(X) \leq s < \infty$, that is, $X \in \mathcal{U}_\mathcal{B}^\pi$.

To prove (iii), let $X \in \mathcal{W}_\mathcal{B}^\pi$, which means that we have an exact sequence

$$(\#11) \quad 0 \longrightarrow P^m \longrightarrow \cdots \longrightarrow P^1 \longrightarrow P^0 \longrightarrow X \longrightarrow 0$$

in $\mathcal{B} = \text{Ch}(\mathcal{A})$ where P^0, \dots, P^m are projective objects; i.e. each P^i is a split exact complex of projective objects in \mathcal{A} , and thus each cycle $Z_n(P^i)$ is also projective in \mathcal{A} . As the complexes P^0, \dots, P^m are, in particular, exact, so is X (and the same are all the kernel and cokernel complexes of the chain maps that appear in (#11)). This implies that the functor $Z_n(-)$ leaves the sequence (#11) exact, and the hereby obtained exact sequence

$$0 \longrightarrow Z_n(P^m) \longrightarrow \cdots \longrightarrow Z_n(P^1) \longrightarrow Z_n(P^0) \longrightarrow Z_n(X) \longrightarrow 0$$

shows that $Z_n(X)$ has finite projective dimension in \mathcal{A} , that is, $Z_n(X)$ belongs to $\mathcal{W}_\mathcal{A}^\pi$.

The proof of the “if” part in (iii) is based on a standard construction; see (the dual of) [16, Thm. 3.1.3] (for this argument to work we make use the hypothesis $\text{FPD}(\mathcal{A}) < \infty$).

Part (iv) is just a repetition of part (i) since $\mathcal{F}_\mathcal{B}^\pi = \mathcal{U}_\mathcal{B}^\pi$ and $\mathcal{F}_\mathcal{A}^\pi = \mathcal{U}_\mathcal{A}^\pi$. \square

In \mathcal{A} we also have the subcategories $\mathcal{U}_\mathcal{A}^l$, $\mathcal{C}_\mathcal{A}^l$, $\mathcal{W}_\mathcal{A}^l$ and $\mathcal{F}_\mathcal{A}^l$ from Definition 3.2. Similarly, in $\mathcal{B} = \text{Ch}(\mathcal{A})$ we have the subcategories $\mathcal{U}_\mathcal{B}^l$, $\mathcal{C}_\mathcal{B}^l$, $\mathcal{W}_\mathcal{B}^l$ and $\mathcal{F}_\mathcal{B}^l$. By an argument dual to the proof of Proposition 4.3, one shows the following result.

Proposition 4.4. *Assume that $\text{FID}(\mathcal{A}) < \infty$ and let $Y = \cdots \rightarrow Y_{n+1} \rightarrow Y_n \rightarrow Y_{n-1} \rightarrow \cdots$ be an object in $\mathcal{B} := \text{Ch}(\mathcal{A})$. The following conclusions hold.*

- (i) *Y belongs to $\mathcal{U}_{\mathcal{B}}^l$ if and only if every Y_n belongs to $\mathcal{U}_{\mathcal{A}}^l$.*
- (ii) *Y belongs to $\mathcal{C}_{\mathcal{B}}^l$ if and only if every Y_n belongs to $\mathcal{C}_{\mathcal{A}}^l$.*
- (iii) *Y belongs to $\mathcal{W}_{\mathcal{B}}^l$ if and only if Y is exact and every cycle $Z_n(Y)$ belongs to $\mathcal{W}_{\mathcal{A}}^l$.*
- (iv) *Y belongs to $\mathcal{F}_{\mathcal{B}}^l$ if and only if every Y_n belongs to $\mathcal{F}_{\mathcal{A}}^l$. \square*

We can now prove the main result of this section.

Theorem 4.5. *Let (S, T) be a Sharp–Foxby adjunction on \mathcal{A} , in particular, $\underline{\text{GProj}}(\mathcal{A})$ and $\underline{\text{GInj}}(\mathcal{A})$ are equivalent as triangulated categories by Corollary 3.12. If $\text{FPD}(\mathcal{A}) < \infty$ and $\text{FID}(\mathcal{A}) < \infty$, then degreewise application of S and T yields a Sharp–Foxby adjunction on $\mathcal{B} = \text{Ch}(\mathcal{A})$, and hence $\underline{\text{GProj}}(\mathcal{B})$ and $\underline{\text{GInj}}(\mathcal{B})$ are equivalent as triangulated categories.*

Proof. Write \bar{S} and \bar{T} for the endofunctors on $\mathcal{B} = \text{Ch}(\mathcal{A})$ that are given by degreewise application of S and T , and let η and ε be the unit and counit of the adjunction (S, T) on \mathcal{A} . It is straightforward to verify that (\bar{S}, \bar{T}) is an adjunction on \mathcal{B} with unit $\bar{\eta}$ and counit $\bar{\varepsilon}$ given by $(\bar{\eta}_X)_n = \eta_{X_n}$ and $(\bar{\varepsilon}_X)_n = \varepsilon_{X_n}$, where X is a chain complex and n is an integer.

By assumption, S restricts to an exact functor $S : \mathcal{U}_{\mathcal{A}}^{\pi} \rightarrow \mathcal{U}_{\mathcal{A}}^l$ which maps $\mathcal{W}_{\mathcal{A}}^{\pi}$ to $\mathcal{W}_{\mathcal{A}}^l$; see (SF1) and (SF2) in Definition 3.4. It therefore follows from Propositions 4.3 and 4.4 that \bar{S} restricts to an exact functor $\bar{S} : \mathcal{U}_{\mathcal{B}}^{\pi} \rightarrow \mathcal{U}_{\mathcal{B}}^l$ which maps $\mathcal{W}_{\mathcal{B}}^{\pi}$ to $\mathcal{W}_{\mathcal{B}}^l$, that is, the adjunction (\bar{S}, \bar{T}) also satisfies conditions (SF1) and (SF2). A similar argument shows that this adjunction satisfies (SF3) and (SF4) as well. By (SF5) in Definition 3.4 we know that the unit $\eta_A : A \rightarrow TSA$ of (S, T) is an isomorphism for $A \in \mathcal{U}_{\mathcal{A}}^{\pi}$. From the definition of $\bar{\eta}$ and from Proposition 4.3 it now follows that $\bar{\eta}_X : X \rightarrow \bar{T}\bar{S}X$ is an isomorphism for $X \in \mathcal{U}_{\mathcal{B}}^{\pi}$, that is, (\bar{S}, \bar{T}) satisfies (SF5). Similarly, (\bar{S}, \bar{T}) also satisfies condition (SF6). \square

Corollary 4.6. *Let (S, T) be a Sharp–Foxby adjunction on \mathcal{A} for which $\text{FPD}(\mathcal{A}) < \infty$ and $\text{FID}(\mathcal{A}) < \infty$. Then degreewise application of S and T yields a Sharp–Foxby adjunction on the category $\text{Ch}^2(\mathcal{A})$ of double complexes (also called bicomplexes) in \mathcal{A} .*

Proof. The category $\text{Ch}^2(\mathcal{A})$ of double complexes in \mathcal{A} is naturally identified with the category $\text{Ch}(\text{Ch}(\mathcal{A}))$. Thus, the desired conclusion follows by applying Theorem 4.5 to the category $\text{Ch}(\mathcal{A})$ (in place of \mathcal{A}). However, to do this we must first argue that the theorem’s hypothesis is satisfied, i.e. that the numbers $\text{FPD}(\text{Ch}(\mathcal{A}))$ and $\text{FID}(\text{Ch}(\mathcal{A}))$ are finite. But is immediate from (the proofs of) Propositions 4.3(iii) and 4.4(iii) that these numbers agree with $\text{FPD}(\mathcal{A})$ and $\text{FID}(\mathcal{A})$, which are finite by assumption. \square

Example 4.7. Let A be a commutative noetherian ring with a dualizing module. By Example 3.6 there exists a Sharp–Foxby adjunction on $\text{Mod}(A)$. The finitistic projective/injective dimensions of $\text{Mod}(A)$ are usually referred to as the finitistic projective/injective dimensions of the ring A , and they are denoted by $\text{FPD}(A)$ and $\text{FID}(A)$. These numbers are finite, indeed, one has $\text{FPD}(A) = \dim A \geq \text{FID}(A)$ by [30, Thm. II.(3.2.6) p. 84] and [2, Cor. 5.5], and $\dim A$ is finite by [22, Cor. V.7.2].

Theorem 4.5 and Corollary 4.6 now imply that the category $\text{Ch}(A)$ of chain complexes and the category $\text{Ch}^2(A)$ of double complexes of A -modules both have Sharp–Foxby adjunctions. In particular, there are by Corollary 3.12 equivalences of triangulated categories,

$$\underline{\text{GProj}}(\text{Ch}(A)) \simeq \underline{\text{GInj}}(\text{Ch}(A)) \quad \text{and} \quad \underline{\text{GProj}}(\text{Ch}^2(A)) \simeq \underline{\text{GInj}}(\text{Ch}^2(A)).$$

The key ingredient in the proof of Theorem 4.5 is that in the category $\mathcal{B} = \text{Ch}(\mathcal{A})$ the Gorenstein projective/injective objects can be suitably described in terms of the Gorenstein

projective/injective objects in \mathcal{A} (as recorded in Propositions 4.3 and 4.4). This is also the case for the category $\mathcal{B} = \text{Rep}(Q, \mathcal{A})$ of \mathcal{A} -valued representations of a left and right rooted quiver Q ; see [13, Thm. 3.5.1]; thus by using the same methods as above one can prove:

Theorem 4.8. *Let (S, T) be a Sharp–Foxby adjunction on \mathcal{A} . If one has $\text{FPD}(\mathcal{A}) < \infty$ and $\text{FID}(\mathcal{A}) < \infty$, then vertexwise application of S and T yields a Sharp–Foxby adjunction on $\mathcal{B} = \text{Rep}(Q, \mathcal{A})$, so $\underline{\text{GProj}}(\mathcal{B})$ and $\underline{\text{GInj}}(\mathcal{B})$ are equivalent as triangulated categories. \square*

ACKNOWLEDGEMENT

We thank the referee for comments that led to a clearer and more focused exposition.

REFERENCES

1. Luchezar L. Avramov and Hans-Bjørn Foxby, *Ring homomorphisms and finite Gorenstein dimension*, Proc. London Math. Soc. (3) **75** (1997), no. 2, 241–270. MR1455856
2. Hyman Bass, *Injective dimension in Noetherian rings*, Trans. Amer. Math. Soc. **102** (1962), 18–29. MR0138644
3. Apostolos Beligiannis and Henning Krause, *Thick subcategories and virtually Gorenstein algebras*, Illinois J. Math. **52** (2008), no. 2, 551–562. MR2524651
4. Apostolos Beligiannis and Idun Reiten, *Homological and homotopical aspects of torsion theories*, Mem. Amer. Math. Soc. **188** (2007), no. 883, viii+207. MR2327478
5. David J. Benson, Jon F. Carlson, and Jeremy Rickard, *Thick subcategories of the stable module category*, Fund. Math. **153** (1997), no. 1, 59–80. MR1450996
6. David J. Benson, Srikanth B. Iyengar, and Henning Krause, *Stratifying modular representations of finite groups*, Ann. of Math. (2) **174** (2011), no. 3, 1643–1684. MR2846489
7. David J. Benson and Henning Krause, *Complexes of injective kG -modules*, Algebra Number Theory **2** (2008), no. 1, 1–30. MR2377361
8. Ragnar-Olaf Buchweitz, *Maximal Cohen–Macaulay modules and Tate-cohomology over Gorenstein rings*, University of Hannover, 1986, available at <http://hdl.handle.net/1807/16682>.
9. Lars W. Christensen, Anders Frankild, and Henrik Holm, *On Gorenstein projective, injective and flat dimensions—A functorial description with applications*, J. Algebra **302** (2006), no. 1, 231–279. MR2236602
10. Edgar E. Enochs and Overtoun M. G. Jenda, *Gorenstein injective and projective modules*, Math. Z. **220** (1995), no. 4, 611–633. MR1363858
11. ———, *Relative homological algebra*, de Gruyter Expositions in Mathematics, vol. 30, Walter de Gruyter & Co., Berlin, 2000. MR1753146
12. Edgar E. Enochs, Overtoun M. G. Jenda, and Jinzhong Xu, *Foxby duality and Gorenstein injective and projective modules*, Trans. Amer. Math. Soc. **348** (1996), no. 8, 3223–3234. MR1355071
13. Hossein Eshraghi, Rasool Hafezi, and Shokrollah Salarian, *Total acyclicity for complexes of representations of quivers*, Comm. Algebra **41** (2013), no. 12, 4425–4441. MR3169529
14. Hans-Bjørn Foxby, *Gorenstein modules and related modules*, Math. Scand. **31** (1972), 267–284 (1973). MR0327752
15. Peter Freyd, *Abelian categories. An introduction to the theory of functors*, Harper’s Series in Modern Mathematics, Harper & Row Publishers, New York, 1964. MR0166240
16. Juan Ramon García Rozas, *Covers and envelopes in the category of complexes of modules*, Chapman & Hall/CRC Research Notes in Mathematics, vol. 407, Chapman & Hall/CRC, Boca Raton, FL, 1999. MR1693036
17. James Gillespie, *Model structures on exact categories*, J. Pure Appl. Algebra **215** (2011), no. 12, 2892–2902. MR2811572
18. ———, *Exact model structures and recollements*, J. Algebra **458** (2016), 265–306. MR3500779
19. ———, *Gorenstein complexes and recollements from cotorsion pairs*, Adv. Math. **291** (2016), 859–911. MR3459032
20. Rüdiger Göbel and Jan Trlifaj, *Approximations and endomorphism algebras of modules*, de Gruyter Expositions in Mathematics, vol. 41, Walter de Gruyter GmbH & Co. KG, Berlin, 2006. MR2251271
21. Dieter Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, vol. 119, Cambridge University Press, Cambridge, 1988. MR935124

22. Robin Hartshorne, *Residues and duality*, Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, vol. 20, Springer-Verlag, Berlin, 1966. MR0222093
23. Henrik Holm, *Gorenstein homological dimensions*, J. Pure Appl. Algebra **189** (2004), no. 1-3, 167–193. MR2038564
24. Mark Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999. MR1650134
25. ———, *Cotorsion pairs, model category structures, and representation theory*, Math. Z. **241** (2002), no. 3, 553–592. MR1938704
26. Henning Krause, *The stable derived category of a Noetherian scheme*, Compos. Math. **141** (2005), no. 5, 1128–1162. MR2157133
27. Henning Krause and Greg Stevenson, *A note on thick subcategories of stable derived categories*, Nagoya Math. J. **212** (2013), 87–96. MR3290681
28. Amnon Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001. MR1812507
29. Dmitri O. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models* (in Russian), Tr. Mat. Inst. Steklova **246** (2004), Algebr. Geom. Metody, Svyazi i Prilozh., 240–262, English translation in Proc. Steklov Inst. Math. 2004, no. 3 (246), 227–248. MR2101296
30. Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89. MR0308104
31. Rodney Y. Sharp, *Finitely generated modules of finite injective dimension over certain Cohen–Macaulay rings*, Proc. London Math. Soc. (3) **25** (1972), 303–328. MR0306188
32. Jan Štíviček, *Exact model categories, approximation theory, and cohomology of quasi-coherent sheaves*, Advances in representation theory of algebras, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2013, pp. 297–367. MR3220541
33. Ryo Takahashi, *Classifying thick subcategories of the stable category of Cohen–Macaulay modules*, Adv. Math. **225** (2010), no. 4, 2076–2116. MR2680200
34. Jinzhong Xu, *Flat covers of modules*, Lecture Notes in Mathematics, vol. 1634, Springer-Verlag, Berlin, 1996. MR1438789
35. Xiaoyan Yang and Zhongkui Liu, *Gorenstein projective, injective, and flat complexes*, Comm. Algebra **39** (2011), no. 5, 1705–1721. MR2821502
36. Fatemeh Zareh-Khoshchereh, Mohsen Asgharzadeh, and Kamran Divaani-Aazar, *Gorenstein homology, relative pure homology and virtually Gorenstein rings*, J. Pure Appl. Algebra **218** (2014), no. 12, 2356–2366. MR3227307
37. Yuefei Zheng and Zhaoyong Huang, *Triangulated equivalences involving Gorenstein projective modules*, Canad. Math. Bull. **60** (2017), no. 4, 879–890. MR3710670

(G.D.) DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30100 MURCIA, SPAIN
E-mail address: georgios.dalezios@um.es

(S.E.) DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30100 MURCIA, SPAIN
E-mail address: sestrada@um.es

(H.H.) DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETS-
 PARKEN 5, 2100 COPENHAGEN Ø, DENMARK
E-mail address: holm@math.ku.dk
URL: <http://www.math.ku.dk/~holm/>