

**The tumor necrosis factor  $\alpha$  of the bony fish seabream exhibits the *in vivo* proinflammatory and proliferative activities of its mammalian counterparts, yet it functions in a species-specific manner**

J. García-Castillo, E. Chaves-Pozo, P. Olivares, P. Pelegrín, J. Meseguer and V. Mulero\*

Department of Cell Biology, Faculty of Biology, University of Murcia, 30100 Murcia (Spain)

\*Corresponding autor: Fax: +34 968 363963, e-mail: vmulero@um.es

## **Abstract**

Information on the bioactivities of non-mammalian cytokines is scant due to the lack of the recombinant molecules and specific antibodies. We produced the mature predicted peptide of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) from the bony fish gilthead seabream (*Sparus aurata* L.) (sbTNF $\alpha$ ), and its biological role was determined *in vitro* and *in vivo*. We first demonstrated by analytical size-exclusion chromatography that sbTNF $\alpha$  is an oligomeric protein but the dimer appears to predominate over the trimeric form, in contrast to mammalian TNF $\alpha$ . Intraperitoneal injection of native sbTNF $\alpha$  resulted in (i) priming of the respiratory burst of the peritoneal exudate and head-kidney (HK) leukocytes, the latter being the bone marrow equivalent in fish; (ii) rapid recruitment of phagocytic granulocytes to the injection site, and (iii) induction of granulopoiesis in the HK. Interestingly, sbTNF $\alpha$  was able to induce a strong proliferation of HK cells *in vitro*, whereas human TNF $\alpha$  did not. Conversely, sbTNF $\alpha$  was not cytotoxic for murine L929 fibroblasts.

**Key words:** Cytokine; TNF $\alpha$ ; inflammation; evolution; fish.

## Introduction

Tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) is a pleiotropic cytokine that plays pivotal roles in the organization and functions of the immune system. TNF $\alpha$  exerts its functions by interacting with two distinct receptors, TNFR1 (p60TNFR or CD120a) [1] and TNFR2 (p80TNFR or CD120b) [2]. TNFR1 is expressed in virtually all mammalian cell types, whereas TNFR2 is expressed only in cells of the immune system and in endothelial cells [3, 4]. The TNFR1 intracellular sequence contains a death domain motif, which is also found in some receptors of the TNF superfamily and which allows the recruitment of death domain-containing adapter proteins involved in cell death signaling in many cell types [5]. Unlike TNFR1, TNFR2 does not contain a death domain motif but still recruits adapter proteins to perform its intracellular signaling activities. As well as being proliferative in some cell types, engagement of TNFR2 has been found to have proapoptotic effects of varying magnitude [6]. On the other hand, the binding of TNF to TNFRs, especially to TNFR1, can also activate the NF- $\kappa$ B transcription factor [7] that serves as a critical regulator of inducible expression of many genes [8]. Functional analogues of mammalian TNF $\alpha$  and TNFR have been described in non-mammalian vertebrates, such as fish, and even in invertebrates [9]. The analogy is based mainly on the cross-reactivity of antibodies elicited against mammalian TNF $\alpha$ , the sensitivity of non-mammalian immune cells to the action of mammalian recombinant TNF $\alpha$ , and the responsiveness of mammalian immune cells to non-mammalian TNF-like factors. Thus, the presence of TNF and TNFR-like molecules in fish was suggested from early studies, where human recombinant TNF $\alpha$  (hTNF $\alpha$ ) was shown to enhance rainbow trout *Oncorhynchus mykiss* neutrophil migration and respiratory burst activity, effects which were inhibited by monoclonal antibodies (mAbs) against human TNFR1 [10]. However, without knowing the amino acid or gene sequences of putative non-mammalian cytokine analogues, unequivocal demonstration of a phylogenetic relationship between mammalian cytokines and their

invertebrate and lower vertebrate functional analogues has not been possible. The existence of TNF in fish has been directly confirmed by the identification of a TNF $\alpha$  gene orthologue in several fish species [11 – 14], while two TNFR-like genes have been identified in the Japanese flounder *Paralichthys olivaceus* [15]. Surprisingly, however, no work has addressed the question of whether TNF $\alpha$  shows conserved biological activities in fish and whether or not it displays species specificity, as seems to be indicated by the above-mentioned studies. Therefore, we produced the mature predicted peptide of TNF $\alpha$  from the bony fish, gilthead seabream (*Sparus aurata* L.), and determined its biological role *in vitro* and *in vivo* from its effects on the respiratory burst, mobilization and proliferation of phagocytes. Overall, our results indicate that vertebrate TNF $\alpha$  displays selective species specificity yet evolutionarily conserved biological activities.

## **Materials and methods**

### **Animals**

Healthy specimens (100 g mean weight) of the hermaphroditic protandrous marine teleost gilthead seabream (*S. aurata* L.) were obtained from Culmarex S.A. (Murcia, Spain). They were kept in 260-l running-seawater aquaria (flow rate 1500 l/h) at 20°C with a 12-h light/dark cycle, and were fed with a commercial pellet diet (Trouvit) at a feeding rate of 15 g dry diet/kg biomass of fish/day. The studies presented in this manuscript were approved by the Bioethical Committee of the University of Murcia.

### **Cloning of gilthead seabream TNF $\alpha$ into expression vector**

Gilthead seabream mature TNF $\alpha$  (sbTNF $\alpha$ ) (GenEMBL accession no. AJ413189) was obtained by polymerase chain reaction (PCR) amplification using a proof-reading DNA polymerase (Ecotaq Plus; Ecogen, Barcelona, Spain). The PCR primers were FE4 (5'-AAGGATCCGCTGAAGCGCATCAGCAGC-3') and RE5 (5'-AAGGATCCTTAAAGTGCAAACACACCAAA-3'), and *Escherichia coli* lipopolysaccharide (LPS; Sigma, Madrid, Spain) stimulated head-kidney (HK; bone marrow equivalent) cDNA was used as template. The PCR amplified fragment was cloned into the SmaI site of the pBluescript SK+ vector and then subcloned into the BamHI site of the pET15b vector for bacterial expression of the His6-tagged protein.

### **Overexpression and purification of sbTNF $\alpha$**

*E. coli* BL21(DE3) cells freshly transformed with the pET15b-sbTNF $\alpha$  construct were cultured overnight in LB ampicillin medium. After dilution into fresh LBampicillin, the cultures were grown at 37 °C to A600 of 0.8 and induced with 1 mM isopropyl-D-thiogalactoside (IPTG; Applichem, Darmstadt, Germany) for 0.25 – 4 h at either 25 or 37°C. Protein expression in

whole-cell extracts was checked by centrifuging 0.1 ml of induced culture (14,000 rpm in a Microfuge), and the cell pellet was lysed by boiling in SDS-loading buffer for analysis by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) [16] and Western blotting using an anti-polyhistidine mAb (Sigma). To check the solubility of the expressed protein, the cell pellet (obtained as above) was suspended in buffer A (50 mM Tris, 2 mM EDTA, 200 mM NaCl, 4 mM 2-mercaptoethanol, pH 7.5), sonicated and centrifuged, and the supernatant and pellet were separately analyzed by Western blotting.

Pelleted cells from 0.25-l cultures induced for 2 h at 25 °C were suspended in 5 ml of ice-cold extraction buffer (50 mM Tris-HCl, 150 mM NaCl, pH 7.8) containing protease inhibitors (Cocktail P8340; Sigma) and sonicated as above. Recombinant His6-tagged sbTNF $\alpha$  protein from clarified supernatants was purified using TALON metal affinity resin and the accompanying purification protocol (BD Biosciences, Madrid, Spain). After purification, the His6 tag was removed using thrombin agarose (Recom-T; Sigma) and aliquots were stored at -80 °C in the presence of 10 % glycerol.

### **Polyclonal anti-sbTNF $\alpha$ antibody**

Anti-sbTNF $\alpha$  rabbit polyclonal antibody was obtained by standard procedures using native sbTNF $\alpha$  for immunization [17]. The reactivity of the antiserum to sbTNF $\alpha$  was checked by Western blotting (see above).

### **Analytical size-exclusion chromatography**

Analytical high-performance liquid chromatography (HPLC) size-exclusion data were obtained at room temperature using a Superdex-200 column equilibrated with buffer A with 150 mM NaCl, which is sufficient to minimize non-specific interactions with the column matrix. Column calibration was done using vitamin B12 (1.355 kDa), cytochrome c (12.4 kDa),

carbonic anhydrase (29 kDa), ovalbumin (43 kDa), bovine serum albumin (BSA) (66 kDa), yeast alcohol dehydrogenase (150 kDa), and  $\beta$ -amylase (200 kDa) (all from Sigma). Samples (150  $\mu$ l) of sbTNF $\alpha$  at 10 – 100  $\mu$ M were injected at 0.4 ml/min, and the elution was tracked by absorbance at 280, 235, and 220 nm. Void ( $V_o$ ) and total ( $V_t$ ) bed volumes were determined using blue dextran (2000 kDa; Sigma) and vitamin B12, respectively. Elution volumes,  $V_e$ , were assigned for sbTNF $\alpha$  by verifying peak identities by Coomassie-stained SDS-PAGE and Western blotting. Stokes radii,  $R_S$  (in nm) for the standards were obtained from Potschka [18]. The following calibration curves were generated from the data for the standards employing SigmaPlot (Jandel Scientific, San Rafael, Calif.) with correlation coefficients  $\geq 0.99$  in each case:  $\log M_r = 7.56 - 0.206 V_e$ . This was then used to estimate the apparent  $M_r$  for sbTNF $\alpha$  [19].

### **Treatments and cell culture**

For the *in vivo* studies, fish were injected intraperitoneally (i. p.) with 1 ml phosphate-buffered saline (PBS) containing either 10  $\mu$ g of recombinant sbTNF $\alpha$  or buffer alone (negative control). At 4, 16 and 40 h post-injection, HK, peripheral blood and peritoneal exudate (PE) leukocytes were obtained as described elsewhere [20] and processed for subsequent chemiluminescence and flow cytometry analysis (see below). In parallel, groups of three fish (treated as above) were injected i. p. with 50  $\mu$ g/kg body weight of 5-bromo-2'-deoxyuridine (BrdU) and, after 2 h, the HK was removed and processed for light microscopy.

For the *in vitro* studies, seabream HK leukocytes from non-injected fish were incubated at 25  $^{\circ}$ C for different periods of time with 0.1 – 100  $\mu$ g/ml sbTNF $\alpha$  in sRPMI [RPMI-1640 culture medium (Gibco, Madrid, Spain) adjusted to gilthead seabream serum osmolarity (353.33 mOs) with 0.35 % NaCl] supplemented with 5 % fetal calf serum (FCS) (Gibco) and 100 IU/ml penicillin and 100  $\mu$ g/ml streptomycin (Biochrom, Berlin, Germany).

Murine L929 fibrosarcoma cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% FCS and penicillin/streptomycin at 37°C.

### **Chemiluminescence assay**

Respiratory burst activity was measured as the luminol-dependent chemiluminescence produced by HK and PE leukocytes [21]. This was brought about by adding 100 µM luminol (Sigma) and 1 µg/ml phorbol myristate acetate (PMA; Sigma), while the chemiluminescence was recorded every 117 s for 1 h in a FLUOstart luminometer (BGM LabTech, Offenburg, Germany). The values reported are the average of octuple readings, expressed as the slope of the reaction curve from 117 to 1170 s, from which the apparatus background was subtracted.

### **Immunofluorescence staining and flow cytometry analysis**

Aliquots of 10<sup>5</sup> cells were washed in PBS containing 2 % FCS and 0.05% sodium azide, incubated for 20 min on ice with 100 µl of a 1/100 dilution of a mAb specific to gilthead seabream acidophilic granulocytes (G7) [20], washed, and incubated with 50 µl of a 1/100 dilution of FITC-labeled anti-mouse IgG F(ab')<sub>2</sub> fragment of goat antibody (Sigma) for 20 min on ice. Cells were then washed twice and data were collected in the form of twoparameter forward scatter (FSC) and side scatter (SSC) dot plot and green fluorescence (FL1) or red fluorescence (FL2) histograms using a FACS (BD Biosciences). The specificity of staining was checked using a mAb isotype standard.

### **Light microscopy and immunohistochemical staining**

The HKs were fixed in Bouin-Hollande fluid solution, embedded in Paraplast Plus (Sherwood Medical, Norfolk, Nebr.) and sectioned at 5 µm. After dewaxing and rehydration, the sections were subjected to an indirect immunocytochemical method [22] using the G7 mAb



[20] or an anti-BrdU mAb (BD Biosciences), at the optimal dilution of 1:10 or 1:5, respectively. The specificity of the reactions was determined by omitting the first antiserum.

### **Cytotoxicity and proliferation assays**

Murine L929 and seabream HK cells were seeded in 96well microtiter plates at  $10^4$  and  $5 \times 10^5$  cells/well, respectively. The next day, a serial dilution of sbTNF $\alpha$  or hTNF $\alpha$  (Sigma) in culture medium supplemented with FCS was added to the cells in the presence or absence of actinomycin D (5 mg/ml; Sigma). After 24 h incubation, surviving mouse L929 cells were quantified by the 2,3bis(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide (XTT; Sigma) colorimetric method [23]. Similarly, the proliferation of seabream primary HK cells was determined after 48 and 72 h of incubation by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT; Sigma) colorimetric method [24] and further confirmed by hemocytometer counts.

### **Protein determination**

The protein concentrations of recombinant sbTNF $\alpha$  samples and cell lysates were estimated by the BCA protein assay reagent (Pierce, Rockford, Ill.) using BSA as a standard.

### **Statistical analysis**

Data were analyzed by one- or two-way analysis of variance and an unpaired Student's t test to determine difference between groups. A quantitative study of the FC results was made using the statistical option of the Lysis Software Package (BD Biosciences).

## Results

### Recombinant mature sbTNF $\alpha$ shows an oligomeric nature

Western blot using an anti-polyhistidine mAb revealed that recombinant His6-tagged sbTNF $\alpha$  was induced as early as 15 min after the addition of IPTG at both 25 and 37 °C, although the amount of soluble protein rapidly decreased at 37°C, probably due to the formation of inclusion bodies, while it remained soluble for up to 4 h at 25 °C (fig. 1 A). Interestingly, an additional polypeptide of about 40 kDa was also detected by the anti-polyhistidine mAb, which might represent a dimeric form of the sbTNF $\alpha$ .

The above results, together with the trimeric nature of bioactive mammalian TNF $\alpha$  [25, 26], prompted us to assess the oligomeric nature of recombinant sbTNF $\alpha$  by analytical gel filtration HPLC [19]. Native His6-tagged sbTNF $\alpha$  was affinity-purified from cultures induced at 25 °C for 2 h and the tag subsequently cleaved off by thrombin (fig. 1 B). sbTNF $\alpha$  eluted in a main peak with a molecular mass of 41 kDa and a secondary peak with a molecular mass of 60 kDa (fig. 2), which may correspond to sbTNF $\alpha$  dimers and trimers, respectively. No additional peaks were detected, indicating that recombinant sbTNF $\alpha$  shows an oligomeric nature and that dimers predominate over trimers.

sbTNF $\alpha$  i.p. injection results in phagocyte activation, mobilization, and proliferation

The biological activities of recombinant sbTNF $\alpha$  were evaluated by injecting fish i. p. with 10 mg of the purified protein. Injection resulted in the priming of the respiratory burst activity of PE (fig. 3A) and HK (fig. 3B) cells triggered by PMA. The maximum enhancement of PE cell respiratory burst activity was observed 16 h post-injection, although a significant priming effect was seen as early as 4 h, lasting to about 40 h post-injection. In HK cells, however, the respiratory burst was only primed 40 h post-injection but remained at basal levels at 4 and 16 h following the i. p. injection of sbTNF $\alpha$ . A weak, but significant ( $p < 0.05$ ), time-dependent

increase in the respiratory burst of both HK and PE cells was observed after the i. p. injection of PBS containing the protein purification buffer alone.

Although gilthead seabream leukocytes are unable to respond to concentrations of LPS below 5 mg/ml [27], we checked whether the observed priming effects on the respiratory burst of PE and HK cells after sbTNF $\alpha$  i.p. injection was solely due to the recombinant cytokine rather than to possible contamination with LPS from *E. coli*. For this reason and taking into account that LPS is heat resistant, we evaluated the activity of sbTNF $\alpha$  samples that had been pre-heated at 80°C for 30 min. Figure 4 shows that pre-heated sbTNF $\alpha$  was unable to prime the respiratory burst of PE cells, confirming that sbTNF $\alpha$  was actually responsible for the observed *in vivo* priming effects. We next evaluated the proinflammatory activity of sbTNF $\alpha$  by FACS using a mAb specific to gilthead seabream acidophilic granulocytes, since this cell type represents the main phagocytic cell of this species and has been considered functionally equivalent to the neutrophil of higher vertebrates [20]. Injection of fish i.p. with sbTNF $\alpha$  resulted in a dramatic mobilization of leukocytes 16 h post-injection (fig. 5A) concomitant with a rapid (4 h post-injection) increase in the percentage of acidophilic granulocytes (fig. 5B), while a much slower and weaker increase in both the total amount of PE cells and the percentage of acidophilic granulocytes was also observed in buffer-injected fish (fig. 5A and data not shown). These changes were matched by those occurring in the peripheral blood, where the percentage of acidophilic granulocytes significantly increased at 16 h and slightly decreased at 40 h following the injection of sbTNF $\alpha$ , but remained unaltered in buffer-injected fish (fig. 5B and data not shown). The amount of peripheral blood leukocytes (PBL), however, was similar in control and sbTNF $\alpha$ -injected groups (data not shown). Interestingly, the injection of sbTNF $\alpha$  also resulted in a marked increase at 16 h post-injection of the percentage of acidophilic granulocytes in the HK, the bone-marrow equivalent of fish (fig. 5B), while

injection of buffer alone resulted in a slower, but significant ( $p < 0.05$ ), increased percentage of these immune cells (data not shown).

The above results prompted us to investigate whether sbTNF $\alpha$  injection was able to enhance the proliferation rate of HK granulopoietic cells (fig. 6A). In fact, while proliferating cells (i.e. BrdU $^{+}$  cells) were mainly restricted to the periphery of HK granulopoietic areas in PBS-injected fish (fig. 6B), an abundant number of proliferating cells were distributed throughout the entire organ 16 h (fig. 6C) and, to some extent, 40 h (fig. 6D) following injection of sbTNF $\alpha$ . However, the HK from control and TNF $\alpha$ -injected groups showed bigger granulopoietic areas 40 h post-injection when compared to 4 and 16-h post-injected fish, confirming the increased percentage of HK acidophilic granulocytes observed by FACS at the longest post-injection time.

### **sbTNF $\alpha$ fails to prime the *in vitro* phagocyte respiratory burst, but acts as a growth-promoting factor**

We next examined the *in vitro* activity of sbTNF $\alpha$ . Figure 7 shows that sbTNF $\alpha$  failed to prime the respiratory burst of HK leukocytes triggered by PMA. Similar results were obtained with PE leukocytes and using 0, 10, 100, or 1000 ng/ml PMA, 250 mg/ml concanavalin A (ConA) or 1 mg/ml opsonized zymosan to trigger the respiratory burst (data not shown). As TNF $\alpha$  is involved in cell death signaling in many mammalian cell types, we therefore assessed cell viability by means of the commonly used MTT colorimetric assay [24]. Of note is the fact that the addition of sbTNF $\alpha$  did not have a significant impact on the viability of HK cells after 24 h of incubation (data not shown), but markedly increased (in a dose-dependent manner) the proliferation of HK cells after 48 and 72 h incubation (fig. 8 A). As expected, the addition of 5 mg/ml ConA induced a strong proliferation of HK cells. Notably, the growth-promoting

effect of the sbTNF $\alpha$  was thermolabile (fig. 8B) and could be partially abrogated with neutralizing anti-sbTNF $\alpha$  antibodies (fig. 8 C).

### **Mammalian and fish TNF $\alpha$ show selective species specificity**

Earlier studies seemed to indicate that fish, and even invertebrate, cells are able to respond to recombinant hTNF $\alpha$  [9, 10]. We therefore studied whether seabream HK leukocytes would respond to hTNF $\alpha$  and whether mouse cells would respond to sbTNF $\alpha$ . For the latter, we used the well-established L929 cytotoxic assay. Interestingly, hTNF $\alpha$  was unable to affect the proliferation of HK cells (fig. 9), in sharp contrast to the strong growth-promoting effect of sbTNF $\alpha$  on these cells (fig. 8 A). Conversely, sbTNF $\alpha$  had no cytotoxic effect on L929 cells, whereas hTNF $\alpha$  killed these cells in both the absence and presence of actinomycin D (IC<sub>50</sub> = ~1 ng/ml and < 0.1 ng/ml, respectively) (fig. 10). These results suggest that sbTNF $\alpha$  failed to interact and/or activate either murine TNFR1 or TNFR2.

## Discussion

Our current knowledge of invertebrate cytokines and, to some extent, of lower vertebrate cytokines is based mainly on immunocytochemical methods and functional assays. Cross-reactivity based on mammalian antibodies to mammalian cytokines or receptors does not unequivocally prove that the antigen in question is present in invertebrates or in lower vertebrates. In the absence of a further molecular characterization of the cross-reacting molecules, any conclusions regarding homology or common evolutionary origin remain presumptive [9]. The molecular and functional characterization of fish cytokines is now possible due to the availability of the sequences of various cytokine genes from different species as well as the full genome sequence of the fugu and zebrafish. Nevertheless, bony fish are the vertebrate group that contain the highest number of species as well as the highest phylogenetic diversity. Therefore, findings referring to other classes of vertebrate, namely mammals and birds, do not necessarily apply to all bony fish species. In this report, we produced recombinant TNF $\alpha$  from the marine fish gilthead seabream, and studied its oligomeric nature as well as its biological activities in fish (seabream) and mammalian (mouse) cells. First, we demonstrated by HPLC that recombinant mature sbTNF $\alpha$  has an oligomeric nature, since the monomer is completely absent under native conditions, while the dimer is the main species and is accompanied by a much less abundant trimer. Although monomeric, dimeric and trimeric species of trout recombinant His6-tagged TNF $\alpha$  have recently been described by means of SDS-PAGE analysis [28], the reducing conditions used during electrophoresis may induce the transition between native oligomers and denatured monomers, as occurs with mammalian TNF $\alpha$  after treatment with particular agents [29, 30]. In any case, the predominant dimeric nature of fish TNF $\alpha$  is surprising in view of the fact that bioactive mammalian TNF $\alpha$  is a trimer and that the monomer lacks biological activity [25, 26, 30]. Nevertheless, the demonstration that monomeric sbTNF $\alpha$  does not exist in native conditions

suggests that oligomerization is also required for the bioactivity of fish TNF $\alpha$ . Whether both dimers and trimers of fish TNF $\alpha$  are bioactive must await further studies.

We demonstrate the *in vivo* proinflammatory activities of fish TNF $\alpha$  for the first time in a lower vertebrate. When injected i.p., sbTNF $\alpha$  is biologically active and able to regulate the main activities of innate immune cells at local and systemic levels. These activities include: (i) the priming of the respiratory burst of PE and HK cells; (ii) the recruitment of local, and likely HK, phagocytic cells to the site of injection, and (iii) the induction of granulopoiesis. In contrast, sbTNF $\alpha$  is unable to prime *in vitro* the respiratory burst of HK cells triggered by different agents, including PMA, ConA, and opsonized zymosan (fig. 7 and data not shown). In mammalian neutrophils and macrophages, some investigators have also noted that TNF $\alpha$  lacks a priming effect on PMA-triggered superoxide formation [31, 32], but others have found that TNF $\alpha$  primes the oxidative burst activity triggered by PMA [33, 34] or parasites [35, 36]. In the trout, a recent *in vitro* study showed that recombinant trout TNF $\alpha$  enhances leukocyte migration and phagocytosis, but the respiratory burst of leukocytes treated with trout TNF $\alpha$  was not evaluated [28]. Our apparently contradictory *in vitro* and *in vivo* data on the effects of sbTNF $\alpha$  on the phagocyte respiratory burst might be explained by taking into account that the activation of the respiratory burst and the recruitment of acidophilic granulocytes peak simultaneously in the peritoneum (16 h post-injection). Therefore, the observed *in vivo* effect of sbTNF $\alpha$  on the respiratory burst may be the result of an increased percentage of acidophilic granulocytes in this immune tissue rather than a true priming effect of resident cells. In the HK, however, the priming of the respiratory burst was observed 40 h post-injection of sbTNF $\alpha$ , whereas the increased percentage of acidophilic granulocytes was observed earlier (i.e., 16 h post-injection). Therefore, this will need further investigation with immunopurified acidophilic granulocytes and/or single-cell analysis of the respiratory burst by FACS.

The ability of sbTNF $\alpha$  to regulate cellular proliferation *in vitro* and *in vivo* was also demonstrated in this study. This is particularly important in view of the fact that a ‘doubleedged sword’ role has been assigned to mammalian TNF $\alpha$ , since it either induces cellular proliferation, survival, differentiation, or apoptosis [37]. We observed no apoptotic or cytotoxic effect of sbTNF $\alpha$  on HK leukocytes, but rather a strong growth-promoting effect both *in vitro* and *in vivo*. Notably, while proliferation is mainly restricted to the periphery of granulopoietic areas in the HK of control fish, the injection of sbTNF $\alpha$  results in a marked increase in the number of proliferating cells that appear scattered throughout the hematopoietic organ. In contrast, a large number of studies have suggested that TNF $\alpha$  negatively regulates the growth of murine and human hematopoietic stem cell populations [38 – 40]. Nevertheless, in a more recent demonstration TNF $\alpha$  did not appear to enhance apoptosis or affect the cell cycle distribution of cultured progenitors but, rather, promoted myeloid differentiation [41]. In addition, TNF $\alpha$  is able to skew monocyte differentiation from macrophages to dendritic cells and this facilitates the induction of adaptive immunity [42]. Studies are in progress in our laboratory to ascertain whether similar mechanisms operate in fish, that is whether TNF $\alpha$  is able to regulate the differentiation of fish hematopoietic cells and/or whether it might facilitate the induction of fish adaptive immunity. This will be interesting not only from a phylogenetic point of view, but also for its potential use as vaccine adjuvant in aquaculture.

The most important finding of this study is that mammalian and fish TNF $\alpha$  show species specificity. Thus, hTNF $\alpha$  is able to kill murine L929 cells whereas sbTNF $\alpha$  cannot. Conversely, hTNF $\alpha$  is unable to affect the proliferation of seabream HK cells, while sbTNF $\alpha$  is a strong growth-promoting factor for such cells. In view of the results reported in this study one is tempted to speculate that the species specificity of TNF $\alpha$  may be related to the different oligomeric natures of hTNF $\alpha$  (trimeric) and sbTNF $\alpha$  (predominantly dimeric) and/or to the



biological activities of soluble versus membrane-bound TNF $\alpha$ . In fact, mammalian TNFR2, unlike TNFR1, is poorly activated by soluble TNF $\alpha$ , and efficient activation of TNFR2 requires membrane-bound TNF $\alpha$  [43]. Alternatively, the homology between fish and mammalian TNF $\alpha$ s, which show about 35% identity and 55% similarity, may not be sufficient to allow cross-receptor engagement. This last hypothesis must be taken into account, since hTNF $\alpha$  binds to murine TNFR1 but not to murine TNFR2 [44], despite the high homology between mTNF $\alpha$  and hTNF $\alpha$  (78 % identity and 90 % similarity). Whatever the case, the molecular and functional characterization of fish TNFRs will be needed before the reasons for the species specificity of TNF $\alpha$  in vertebrates can be ascertained.

In conclusion, we demonstrated for the first time an *in vivo* physiological role for TNF $\alpha$  in fish. sbTNF $\alpha$  is mainly found as a dimer and is able to regulate the respiratory burst, the mobilization and the proliferation of phagocytes. In addition, sbTNF $\alpha$  is not bioactive for mammalian cells and hTNF $\alpha$  is inactive in fish cells. Our data point to the need for further research from a phylogenetic perspective to complete our knowledge about the complex evolution of TNF $\alpha$  in vertebrates.

**Acknowledgements.** This work was supported by the Spanish Ministry of Science and Technology (grants BIO2001-2324-C02-02, AGL2002-03529 and AGL2002-04306-C03-01, and fellowship to J. García-Castillo), Spanish Ministry of Education, Culture and Sport (fellowship to E. Chaves-Pozo) and Fundación Séneca, Coordination Centre for Research (grant PI-51/00782/FS/01, and fellowship to P. Pelegrín). We thank Dr. S Padmanabhan for his valuable help with the HPLC study, and the Tissue Culture Facility (SACE, University of Murcia) for maintaining the cell lines.

## References

- 1 Loetscher H., Pan Y. C., Lahm H. W., Gentz R., Brockhaus M., Tabuchi H. et al. (1990) Molecular cloning and expression of the human 55 kd tumor necrosis factor receptor. *Cell* 61: 351–359
- 2 Smith C. A., Davis T., Anderson D., Solam L., Beckmann M. P., Jerzy R. et al. (1990) A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. *Science* 248: 1019–1023
- 3 Brockhaus M., Schoenfeld H. J., Schlaeger E. J., Hunziker W., Lesslauer W. and Loetscher H. (1990) Identification of two types of tumor necrosis factor receptors on human cell lines by monoclonal antibodies. *Proc. Natl. Acad. Sci. USA* 87: 3127 – 3131
- 4 Hohmann H. P., Remy R., Brockhaus M. and van Loon A. P. (1989) Two different cell types have different major receptors for human tumor necrosis factor (TNF $\alpha$ ). *J. Biol. Chem.* 264: 14927 – 14934
- 5 Wajant H, Henkler F. and Scheurich P. (2001) The TNF-receptor associated factor family: scaffold molecules for cytokine receptors, kinases and their regulators. *Cell Signal.* 13: 389–400
- 6 Grell M., Becke F. M., Wajant H., Mannel D. N. and Scheurich P. (1998) TNF receptor type 2 mediates thymocyte proliferation independently of TNF receptor type 1. *Eur. J. Immunol.* 28: 257 – 263
- 7 McFarlane S. M., Pashmi G., Connell M. C., Littlejohn A. F., Tucker S. J., Vandenabeele P. et al. (2002) Differential activation of nuclear factor-kB by tumour necrosis factor receptor subtypes: TNFR1 predominates whereas TNFR2 activates transcription poorly. *FEBS Lett.* 515: 119–126
- 8 Karin M. (1999) How NF-kB is activated: the role of the I $\kappa$ B kinase (IKK) complex. *Oncogene* 18: 6867 – 6874

- 9 Beschin A., Bilej M., Torreele E. and De Baetselier P. (2001) On the existence of cytokines in invertebrates. *Cell. Mol. Life Sci.* 58: 801–814
- 10 Jang S.I., Mulero V., Hardie L.J. and Secombes C.J. (1995) Inhibition of rainbow trout phagocyte responsiveness to human tumor necrosis factor  $\alpha$  (hTNF $\alpha$ ) with monoclonal antibodies to the hTNF $\alpha$  55 kDa receptor. *Fish Shellfish Immunol.* 5: 61–69
- 11 Hirono I., Nam B. H., Kurobe T. and Aoki T. (2000) Molecular cloning, characterization, and expression of TNF cDNA and gene from Japanese flounder *Paralichthys olivaceus*. *J. Immunol.* 165: 4423–4427
- 12 Laing K. J., Wang T., Zou J., Holland J., Hong S., Bols N. et al. (2001) Cloning and expression analysis of rainbow trout *Oncorhynchus mykiss* tumour necrosis factor  $\alpha$ . *Eur. J. Biochem.* 268: 1315–1322
- 13 García-Castillo J., Pelegrín P., Mulero V. and Meseguer J. (2002) Molecular cloning and expression analysis of tumor necrosis factor  $\alpha$  from a marine fish reveal its constitutive expression and ubiquitous nature. *Immunogenetics* 54: 200 – 207
- 14 Saeij J. P., Stet R. J., Vries B. J. de, Muiswinkel W. B. van and Wiegertjes G. F. (2003) Molecular and functional characterization of carp TNF: a link between TNF polymorphism and trypanotolerance? *Dev. Comp. Immunol.* 27: 29–41
- 15 Park C. I., Kurobe T., Hirono I. and Aoki T. (2003) Cloning and characterization of cDNAs for two distinct tumor necrosis factor receptor superfamily genes from Japanese flounder *Paralichthys olivaceus*. *Dev. Comp. Immunol.* 27: 365–375
- 16 Laemmli U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680 – 685
- 17 Ausubel F. M., Brent R., Kingston R., Seidman J. G., Smith J. A. and Struhl K. (1995) *Current Protocols in Molecular Biology*, Supplement 13, pp. 2.4.1 – 2.4.9, Wiley, New York

- 18 Potschka M. (1987) Universal calibration of gel permeation chromatography and determination of molecular shape in solution. *Anal. Biochem.* 162: 47 – 64
- 19 Ackers G. K. (1970) Analytical gel chromatography of proteins. *Adv. Protein Chem.* 24: 343 – 446
- 20 Sepulcre M. P., Pelegrín P., Mulero V. and Meseguer J. (2002) Characterisation of gilthead seabream acidophilic granulocytes by a monoclonal antibody unequivocally points to their involvement in fish phagocytic response. *Cell Tissue Res.* 308: 97–102
- 21 Mulero V., Pelegrín P., Sepulcre M. P., Munoz J. and Meseguer J. (2001) A fish cell surface receptor defined by a mAb mediates leukocyte aggregation and deactivation. *Dev. Comp. Immunol.* 25: 619 – 627
- 22 Sternberger L. A. (1986) *Immunocytochemistry*, 2nd edn Wiley, New York
- 23 Roehm N. W., Rodgers G. H., Hatfield S. M. and Glasebrook A. L. (1991) An improved colorimetric assay for cell proliferation and viability utilizing the tetrazolium salt XTT. *J. Immunol. Methods* 142: 257 – 265
- 24 Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65: 55 – 63
- 25 Corti A., Fassina G., Marcucci F., Barbanti E. and Cassani G. (1992) Oligomeric tumour necrosis factor a slowly converts into inactive forms at bioactive levels. *Biochem. J.* 284: 905– 910
- 26 Ameloot P., Declercq W., Fiers W., Vandenabeele P. and Brouckaert P. (2001) Heterotrimers formed by tumor necrosis factors of different species or muteins. *J. Biol. Chem.* 276: 27098 – 27103
- 27 Pelegrín P., Chaves-Pozo E., Mulero V. and Meseguer J. (2004) Production and mechanism of secretion of interleukin-1b from the marine fish gilthead seabream. *Dev. Comp. Immunol.* 28: 229 – 237

- 28 Zou J., Peddie S., Scapigliati G., Zhang Y., Bols N. C., Ellis A. E. et al. (2003) Functional characterisation of the recombinant tumor necrosis factors in rainbow trout, *Oncorhynchus mykiss*. *Dev. Comp. Immunol.* 27: 813–822
- 29 Alzani R., Cozzi E., Corti A., Temponi M., Trizio D., Gigli M. et al. (1995) Mechanism of suramin-induced deoligomerization of tumor necrosis factor  $\alpha$ . *Biochemistry* 34: 6344 – 6350
- 30 Hlodan R. and Pain R. H. (1995) The folding and assembly pathway of tumour necrosis factor TNF  $\alpha$ , a globular trimeric protein. *Eur. J. Biochem.* 231: 381 – 387
- 31 Tennenberg S. D. and Solomkin J. S. (1990) Activation of neutrophils by cachectin/tumor necrosis factor: priming of Nformyl-methionyl-leucyl-phenylalanine-induced oxidative responsiveness via receptor mobilization without degranulation. *J. Leukoc. Biol.* 47: 217 – 223
- 32 Utsumi T., Klostergaard J., Akimaru K., Edashige K., Sato E. F. and Utsumi K. (1992). Modulation of TNF $\alpha$ -priming and stimulation-dependent superoxide generation in human neutrophils by protein kinase inhibitors. *Arch. Biochem. Biophys.* 294: 271 – 278
- 33 Berkow R. L., Wang D., Larrick J. W., Dodson R. W. and Howard T. H. (1987) Enhancement of neutrophil superoxide production by preincubation with recombinant human tumor necrosis factor. *J. Immunol.* 139: 3783 – 3791
- 34 Phillips W. A. and Hamilton J. A. (1989) Phorbol ester-stimulated superoxide production by murine bone marrow-derived macrophages requires preexposure to cytokines. *J. Immunol.* 142: 2445 – 2449
- 35 Ferrante A. (1989) Tumor necrosis factor  $\alpha$  potentiates neutrophil antimicrobial activity: increased fungicidal activity against *Torulopsis glabrata* and *Candida albicans* and associated increases in oxygen radical production and lysosomal enzyme release. *Infect. Immun.* 57: 2115 – 2122

- 36 Denis M. and Chadee K. (1989). Human neutrophils activated by interferon- $\gamma$  and tumour necrosis factor–a kill *Entamoeba histolytica* trophozoites *in vitro*. *J. Leukoc. Biol.* 46: 270 – 274
- 37 Aggarwal B. B. (2003) Signalling pathways of the TNF superfamily: a double-edged sword. *Nat. Rev. Immunol.* 3: 745 – 756
- 38 Jacobsen F. W., Rothe M., Rusten L., Goeddel D. V., Smeland E. B., Veiby O. P. et al. (1994) Role of the 75-kDa tumor necrosis factor receptor: inhibition of early hematopoiesis. *Proc. Natl. Acad. Sci. USA* 91: 10695 – 10699
- 39 Rusten L. S., Jacobsen F. W., Lesslauer W., Loetscher H., Smeland E. B. and Jacobsen S. E. (1994) Bifunctional effects of tumor necrosis factor  $\alpha$  (TNF  $\alpha$ ) on the growth of mature and primitive human hematopoietic progenitor cells: involvement of p55 and p75 TNF receptors. *Blood* 83: 3152 – 3159
- 40 Zhang Y., Harada A., Bluethmann H., Wang J. B., Nakao S., Mukaida N. et al. (1995) Tumor necrosis factor (TNF) is a physiologic regulator of hematopoietic progenitor cells: increase of early hematopoietic progenitor cells in TNF receptor p55-deficient mice *in vivo* and potent inhibition of progenitor cell proliferation by TNF  $\alpha$  *in vitro*. *Blood* 86: 2930 – 2937
- 41 Dybedal I., Bryder D., Fossum A., Rusten L. S. and Jacobsen S. E. (2001) Tumor necrosis factor (TNF)-mediated activation of the p55 TNF receptor negatively regulates maintenance of cycling reconstituting human hematopoietic stem cells. *Blood* 98: 1782 – 1791
- 42 Chomarat P., Dantin C., Bennett L., Banchereau J. and Palucka A. K. (2003) TNF skews monocyte differentiation from macrophages to dendritic cells. *J. Immunol.* 171: 2262 – 2269

- 43 Grell M., Douni E., Wajant H., Lohden M., Clauss M., Maxeiner B. et al. (1995) The transmembrane form of tumor necrosis factor is the prime activating ligand of the 80 kDa tumor necrosis factor receptor. *Cell* 83: 793 – 802
- 44 Lewis M., Tartaglia L. A., Lee A., Bennett G. L., Rice G. C., Wong G. H. et al. (1991) Cloning and expression of cDNAs for two distinct murine tumor necrosis factor receptors demonstrate one receptor is species specific. *Proc. Natl. Acad. Sci. USA* 88: 2830 – 2834



### **Figure Legends:**

**Figure 1. Production of recombinant sbTNF $\alpha$ .** (A) Western blot analysis of native His6-tagged sbTNF $\alpha$  in soluble fractions from *E. coli* cultures induced with 1 mM IPTG for the indicated times at either 25 or 37°C. The filters were hybridized with anti-polyhistidine mAb. (B) SDS-PAGE analysis of metal affinity-purified sbTNF $\alpha$  before (lane 1) and after (lane 2) removing the His6 tag using thrombinagarose. Proteins were stained with Coomassie Brilliant Blue R-250.

**Figure 2. Oligomeric nature of sbTNF $\alpha$  assayed by size exclusion chromatography.** Samples of sbTNF $\alpha$  at 20 $\mu$ M were applied to a Superdex-200 column. Protein elution was followed by measuring the absorption at 220 nm and 1-min fractions were collected. The column was calibrated with the following standards (arrows): BSA (66 kDa), ovalbumin (43 kDa), carbonic anhydrase (29 kDa) and cytochrome C (12.4 kDa).

**Figure 3. sbTNF $\alpha$  i.p. injection results in the priming of the phagocyte respiratory burst.** Fish were injected i.p. with 1 ml PBS containing either buffer alone or 10  $\mu$ g sbTNF $\alpha$ . At 4, 16, and 40 h post-injection, the respiratory burst activity of PE (A) and HK (B) cells was measured as the luminol-dependent chemiluminiscence triggered by PMA (1  $\mu$ g/ml). Data are presented as mean  $\pm$  SE of five fish. Versus PBS-injected fish: <sup>a</sup>p < 0.05, <sup>b</sup>p < 0.01, <sup>c</sup>p < 0.0001.

**Figure 4. The bioactivity of recombinant sbTNF $\alpha$  samples is heat labile.** Fish were injected i.p. with 1 ml PBS containing 10  $\mu$ g sbTNF $\alpha$  that had been pre-heated or not at 80°C for 30 min. At 16 h post-injection, the respiratory burst activity of PE cells was measured as indicated in the legend to figure 3. Data are presented as mean  $\pm$  SE of three fish. Versus pre-heated sbTNF $\alpha$ -injected fish: \*p < 0.0001.

**Figure 5. sbTNF $\alpha$  i. p. injection results in phagocyte mobilization.** Fish were injected i.p. with 1 ml of PBS containing either buffer alone or 10  $\mu$ g sbTNF $\alpha$ . At 4, 16 and 40 h post-injection, the number of PE cells was determined by hemocytometer counts (A) while the percentage of acidophilic granulocytes in PE, PBL and HK was evaluated by FACS after staining the cells with a mAb specific to these cells (G7) (B). Data are presented as mean  $\pm$  SE of three fish. Horizontal line represents the control value (PBS-injected fish). Versus PBS-injected fish: <sup>a</sup>p < 0.05, <sup>b</sup>p < 0.01, <sup>c</sup>p < 0.001.

**Figure 6. sbTNF $\alpha$  i.p. injection enhances granulopoiesis.** Fish injected i. p. with 1 ml of PBS containing either buffer alone (A, B) or 10  $\mu$ g sbTNF $\alpha$  (C, D) were pulsed for 2 h with 50 mg/kg body weight BrdU. HK sections were immunostained with the G7 mAb (A) or with an anti-BrdU mAb (B–D). Note that proliferating cells (arrow) were mainly restricted to the periphery of granulopoietic areas (AG) in control fish at 16 h post-injection (B), whereas they were abundant and distributed scattered in the HK of sbTNF $\alpha$ -injected fish at 16 h (C) and 40 h post-injection (D). V, blood vessel; \*, melanomacrophage center. Magnification, x10.

**Figure 7. sbTNF $\alpha$  fails to prime the *in vitro* phagocyte respiratory burst.** HK cells were incubated for the indicated times in the presence of 0 – 100 ng/ml sbTNF $\alpha$ . The respiratory burst was assayed as indicated in the legend to figure 3. Data are presented as mean  $\pm$  SE of three fish.


**Figure 8. sbTNF $\alpha$  stimulates *in vitro* the proliferation of HK cells.** Aliquots of  $5 \times 10^5$  HK cells were incubated for 48 or 72 h with 0–100 ng/ml sbTNF $\alpha$  or 5  $\mu$ g/ml ConA (A), or for 72 h with 0–100 ng/ml sbTNF $\alpha$  that had been pre-heated or not at 80°C for 30 min (B), or in the

absence or presence of 0.1  $\mu$ l of either pre-immune or anti-sbTNF $\alpha$  serum (C). After the indicated incubation times, cell proliferation was determined as the reduction of MTT. Data are presented as mean  $\pm$  SE of quadruplicate readings and are representative of three independent experiments. Versus PBS-injected fish: <sup>a</sup>p < 0.05, <sup>b</sup>p < 0.01, <sup>c</sup>p < 0.001; versus pre-heated sbTNF $\alpha$ -injected fish: \*p < 0.01; versus pre-immune serum: #p < 0.05.


**Figure 9. hTNF $\alpha$  was unable to stimulate the proliferation of HK cells.** HK cells were incubated with 0–100 ng/ml hTNF $\alpha$  and cell proliferation assayed 48 and 72 h later, as indicated in the legend to figure 8. Data are presented as mean  $\pm$  SE of quadruplicate readings and are representative of two independent experiments.

**Figure 10. sbTNF $\alpha$  is unable to kill murine L929 cells.** Aliquots of 10<sup>4</sup> L929 were incubated with 0–100 ng/ml of either sbTNF $\alpha$  or hTNF $\alpha$  in the presence or absence of 5  $\mu$ g/ml actinomycin D (ActD). After 24 h incubation, the number of surviving cells was determined by the XTT colorimetric assay. Data are presented as mean  $\pm$  SE of quadruplicate readings and are representative of two independent experiments. Versus control cells: <sup>a</sup>p < 0.01, <sup>b</sup>p < 0.001, <sup>c</sup>p < 0.0001.


**FIGURE 1**




**FIGURE 2**




**FIGURE 3**



**FIGURE 4**




**FIGURE 5**







**FIGURE 6**




**FIGURE 7**



**FIGURE 8**



**FIGURE 9**



**FIGURE 10**

