

2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

This document is the Submitted Manuscript version of a Published Work that appeared in final form in *Bioprocess and Biosystems Engineering*. To access the final edited and published work see <https://doi.org/10.1007/s00449-019-02243-1>.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13 **Development and economic evaluation of an eco-friendly biocatalytic**  
14 **synthesis of emollient esters.**  
15

16  
17  
18 **Authors:** Mar SERRANO-ARNALDOS\*, María Claudia MONTIEL, Salvadora  
19 ORTEGA-REQUENA, Fuensanta MÁXIMO, Josefa BASTIDA.  
20

21  
22 Department of Chemical Engineering, University of Murcia, Campus de Espinardo, 30071  
23 Murcia, Spain.  
24

25  
26 \*Corresponding author: [mar.serrano@um.es](mailto:mar.serrano@um.es), +34 868887924, [https://orcid.org/0000-0001-](https://orcid.org/0000-0001-5919-0951)  
27 [5919-0951](https://orcid.org/0000-0001-5919-0951)  
28  
29  
30  
31  
32

33 **Acknowledgements:** This work has been funded by the Spanish Ministry of Science,  
34 Innovation and Universities (CTQ2015-66723-R) and the European Commission  
35 (FEDER/ERDF). M. Serrano-Arnaldos and S. Ortega-Requena were beneficiaries of a FPI  
36 pre-doctoral scholarship from the Spanish Ministry of Economy and Competitiveness  
37 (MINECO) and a Torres Quevedo grant, respectively. We wish to acknowledge D. Ramiro  
38 Martínez Gutiérrez (Novozymes Spain S.A.) who kindly supplied the biocatalysts.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

During the last decades the understanding and prospects of enzyme-catalysed reactions have been massively widened and there are a number of implemented large-scale enzymatic processes mainly based in the use of commercial biocatalysts. As it might happen that the same process can be successfully carried out by different commercial lipases, the election of the biocatalyst must rely on productivity and economic considerations. This work presents productiveness and direct operation cost evaluation as a key tool for the selection between two commercial lipase catalysts, the versatile but expensive Novozym<sup>®</sup> 435 and a much more economical option, Lipozyme<sup>®</sup> TL IM, in the synthesis of spermaceti, a mixture of emollient esters with cosmetic applications. Proving that Novozym<sup>®</sup> 435 leads to minimum savings of 10% with respect to the cheapest immobilized derivative, biocatalyst cost does not appear to be the major contribution to the economics of the processes under study, due to their great capacity to be recovered and reused. At laboratory scale, the biggest economic investment is caused by substrates, which can be massively reduced at industrial scale by using bulk reagents. In such case, energy cost may be the major contribution to the process economy. This work proposes an optimized process ready to be scaled-up in order to accurately determine the energetic requirements of the possible industrial enzymatic synthesis.

**Key words:** *Candida antarctica*; *Thermomyces lanuginosus*; solvent-free; spermaceti; cost assessment.

## 1. Introduction

While the aim of basic scientific research is to contribute to expand the limits of knowledge by setting or revising fundamental principles, the main goal of applied research is to put into practice this elemental basis for the benefit of society. Nevertheless, looking over the wide number of scientific publications existing, it can be noticed that there are little systematic studies dealing with the feasibility of the production of new commodities, either establishing or substantially improving the industrial processes to obtain them. A clear paradigm of that is the assessment of the large-scale use of enzymatic catalysts, even if the great advantages of the enzymatic routes against chemical pathways are well known: lower energy consumption, increased product quality and/or cleaner and eco-friendly processes. In particular, the use of immobilized lipases for obtaining esters with application in food and cosmetic industries has been the subject of a large number of studies since the last years of 20<sup>th</sup> century [1–3]. However, lipases have not been exploited in its fullness at industrial scale yet, with the exception of a few enterprises, such as Evonik Industries, which commercializes five emollient esters, under the label “enzymatic”, produced by using the well-known *Candida antarctica* lipase B derivative Novozym<sup>®</sup> 435 [4, 5].

Despite lipases are considered one of the most valuable enzymes for synthetic purposes [6], the lack of a widespread biocatalysis-based industry might be chiefly due to the high price of enzymes, that is usually recoup through the immobilization of the lipase and its reuse in several batches. In fact, the investment in an immobilized derivative can be up to 90-95% of the process' cost [7] and so, a large number of viable uses of a biocatalyst are essential to turn the enzymatic synthesis economically profitable in comparison with conventional processes [8]. In this context, one of the objectives of recent research is the development of more active and stable enzymatic catalysts in order to maximize their reuse while reducing their cost [9]. Nonetheless, in view of the number of biocatalysts currently commercialized, it seems that these efforts are still ongoing.

One of the main producers of enzymatic derivatives, Novozymes [10], offers immobilized lipases from different origins adsorbed onto polymeric carriers. Novozym<sup>®</sup> 435 stands out among them owing to its high performance both in hydrolysis and synthesis reactions under solvent-free conditions or in organic solvent media [11–13], but also because of its expensive market price. On the other hand, Lipozyme<sup>®</sup> TL IM, which contains a fungal lipase from *Thermomyces lanuginosus*, may be less active but has a much lower cost [9].

Since the increasing number of publications proving the feasibility of the lipase-based esterification of fatty acids with cetyl alcohol gives an idea of the industrial importance of such wax esters [14–17], which are mainly used in cosmetic and pharmaceutical industries due to their emollient properties. As traces of organic solvents in the final product are highly restricted when used for such purposes, the solvent-free synthesis has been proposed to overcome this limitation [18].

1  
2  
3 In view of the above mentioned, this paper deals with the one-step solvent-free biosynthesis  
4 of the four cetyl esters of the spermaceti, a waxy substance formerly obtained from the head  
5 cavity of the sperm whale (*Physeter macrocephalus*). Spermaceti is composed of a mixture  
6 of cetyl esters and has been synthesised by following its natural proportions [19]. The  
7 enzymatic synthesis of such esters is a well-known process and can be developed by using  
8 lipases from different origin with high yields [20]. Consequently, the enzymatic production  
9 this specialty chemical is evaluated from a technical and an economic point of view in  
10 order to establish the possibility of implementing the process at industrial scale. For that  
11 purpose, some of the main factors contributing to the manufacturing expenses have been  
12 pondered, such as reactor's operation needs, biocatalyst cost and raw materials purchase.  
13 The reusability of the two enzymatic catalysts in successive synthesis batches has been  
14 taken into consideration [21].  
15  
16  
17  
18  
19  
20  
21

## 22 **2. Materials and methods**

### 23 **2.1 Materials**

24  
25  
26 Novozym<sup>®</sup> 435 is made of the isoenzyme B from *Candida antarctica* (CalB) immobilized  
27 by adsorption onto the macroporous DVB/methacrylate co-polymer resin Lewatit<sup>®</sup> VP OC  
28 1600. Its declared activity is 10000 PLU/g, where one unit corresponds to the synthesis of  
29 1  $\mu\text{mol}$  per minute propyl laureate from lauric acid and 1-propanol. Lipozyme<sup>®</sup> TL IM is an  
30 immobilized derivative of the fungal lipase *Thermomyces lanuginosus* (Tll) on a cationic  
31 silicate (250 UIN/g, interesterification units). Both were kindly donated by Novozymes  
32 Spain S.A.  
33  
34  
35

36  
37 Lauric acid (99%) was purchased at Acros Organics (Geel, Belgium); myristic acid (98%),  
38 palmitic acid (98%), stearic acid (95%) and cetyl alcohol at Sigma-Aldrich (Missouri,  
39 USA). The rest of chemicals used were all analytical reagent grade.  
40

### 41 **2.2 Methods**

#### 42 **2.2.1 Synthesis under vacuum**

43  
44  
45  
46 The solvent-free synthesis of the spermaceti analogue was optimized in a Parr 5100 series  
47 tank reactor (Parr Instrument Co., Illinois, USA), which is provided with a 100 mL vessel  
48 heated with a circulating water jacket and a turbine-type vertical stirrer, set at 350 rpm by  
49 the reactor's controller. All the assays were performed under an inert atmosphere composed  
50 of a 54 L/h flow of  $\text{N}_2$  from the autonomous generator 3848 NitroFlow Lab (Parker, Ohio,  
51 USA) and low pressure (21.3 kPa) conditions, which were optimized by the authors in  
52 previous work [22]. 20 g of substrates were used in all cases. Biocatalyst's concentrations  
53 (w/w, referred to substrates) ranging from 0.63% to 5% for Novozym<sup>®</sup> 435, or 10% for  
54  
55  
56  
57  
58  
59  
60

Lipozyme<sup>®</sup> TL IM, were tested during the synthesis of the cetyl esters separately (1:1 molar ratio, 70 °C). The influence of the temperature was studied in the one-step production of spermaceti, by using the best enzymatic catalyst concentration at temperatures comprised between 60 to 80 °C. Reaction mixture was composed of the substrates' amounts needed to obtain a final product with a 15.3% of cetyl laurate, 37.7% of cetyl myristate, 38.8% of cetyl palmitate and 8.2% of cetyl stearate [19]. The evaluation of the biocatalysts reuse was performed for the synthesis of spermaceti under best reaction conditions. After one hour of use, the immobilized derivative was recovered and rinsed with 3 x 10 mL of acetone, filtering and air drying before using them in another batch, with new fresh reagents.

### 2.2.2 Acid value determination

The reaction course can be followed by measuring the acid value, A.V., of samples withdrawn from reaction media, which is defined as the amount of KOH, in milligrams, necessary to neutralize the free carboxyl groups present in 1 g of sample [23].

Conversion can be stated as:

$$Conversion(\%) = \frac{A.V._0 - A.V._t}{A.V._0} \times 100$$

Where A.V.<sub>0</sub> and A.V.<sub>t</sub> are the acid values at reaction's beginning and at a specific sampling time, respectively. Results are the mean of two data and they are reported with their standard deviation.

## 3. Results and discussion

### 3.1 *Process technical optimization*

#### 3.1.1 Influence of biocatalyst concentration

As described in literature, the high cost of enzymes may be one of the limiting factors for the industrial application of biocatalytic systems [7, 8, 21]. This can be reduced by using the right quantity of an immobilized enzyme, which might compromise both technical (adequate space-time yield) and economic requirements.

Figures 1 and 2 represent the esterification process of cetyl alcohol with lauric, myristic, palmitic and stearic acid catalysed by different concentrations of Novozym<sup>®</sup> 435 and Lipozyme<sup>®</sup> TL IM. If both biocatalysts are studied independently, it can be observed that conversion evolves with time mostly in a nonlinear way and results are quite similar independently of the ester synthesized for each enzymatic product. This is in accordance with previous studies, which has proven that even if lower the reaction rates can be noticed

1  
2  
3 as the substrates' chain length increases, such differences were so little for a given amount  
4 of immobilized lipase that, for practical purposes, they do not lead to the preferential  
5 synthesis of some esters against the others and do not compromise their simultaneous  
6 production [20].  
7  
8

9 However, significant differences in reaction rate are evident if comparing both biocatalysts  
10 under same reaction conditions, being quite lower for the TII derivative. This must be  
11 mainly caused by the hydrophilic character of Lipozyme<sup>®</sup> TL IM's carrier, which is not so  
12 suitable for such hydrophobic reagents and products. Similar findings had been described in  
13 the literature, during the synthesis of flavouring esters in organic solvents [24] or oleic acid  
14 and methyl ricinoleate estolides [25], but in those cases, the results were explained by the  
15 1,3- positional selectivity of TII. On the contrary, Novozym<sup>®</sup> 435 displays a great synthetic  
16 activity, due to the hydrophobic nature of its acrylic matrix as well as to the origin of the  
17 lipase, since it has been reported that CalB is very active with a wide variety of esters [26].  
18  
19  
20  
21

22 Despite this, similar product yields (~98.5%) are achieved with this two immobilized  
23 lipases, if allowed the time needed according to the kinetics of the process. Thus, a  
24 concentration of biocatalyst of 2.5% (0.5 g) was selected as the most suitable for  
25 Novozym<sup>®</sup> 435, while a 5% (1 g) was chosen for Lipozyme<sup>®</sup> TL IM.  
26  
27

28 (Figure 1)  
29

30 (Figure 2)  
31

### 32 3.1.2 Effect of the temperature

33

34 Operation temperature has a key role in the synthesis of fatty acid esters in solvent-free  
35 systems, as the reduction of the viscosity of the reaction media cannot be attained through  
36 the use of organic solvents, but through the selection of a temperature that reduces mass  
37 transfer limitations while safeguarding the enzyme's functional conformation. In this sense,  
38 the one-step synthesis of the spermaceti analogue was carried out at 60, 70 and 80 °C.  
39 Lower temperatures were not tested due to the high melting points of the substances  
40 implied in the process, while values above 80 °C were not considered because of energy  
41 consumption.  
42  
43  
44  
45

46 Figure 3 a shows results for Lipozyme<sup>®</sup> TL IM, where it can be observed a loss of  
47 biocatalyst's activity between 70 and 80 °C, pointing out that this immobilized derivative  
48 may be less resistant to temperature. Even if TII is known as a thermostable lipase [9], these  
49 observations are in agreement with findings from other authors, which have reported the  
50 adverse consequences of temperature in Lipozyme<sup>®</sup> TL IM performance during  
51 esterification [24], alcoholysis [27] or transesterification [28] processes.  
52  
53  
54  
55  
56  
57  
58  
59  
60

In contrast, the above mentioned positive effect of temperature is reflected in the conversions reached during the simultaneous synthesis of a spermaceti analogue with Novozym<sup>®</sup> 435 (Figure 3 b), where an improvement of ~11 percentage units in the conversions is obtained at 30 min as temperature increases from 60 to 80 °C.

Hence, and taking into account that similar final conversions (>97.5%) were obtained independently of the temperature used, the optimum value was set at 70 °C, as this temperature reduces the time required to the reagents melting before starting the esterification process.

(Figure 3)

### 3.1.3 Reusability of the commercial biocatalysts

As previously stated, an operational stability that enables the biocatalyst reuse in the greatest number of consecutive batches is of paramount importance for a successful implementation of the enzymatic pathway at industrial scale. Consequently, Lipozyme<sup>®</sup> TL IM and Novozym<sup>®</sup> 435 have been tested in the synthesis of spermaceti under best reaction conditions up to 15 times, with the outcomes showed in Table 1. While losses of biocatalyst mass of 9.6% for Lipozyme<sup>®</sup> TL IM and 2.8% for Novozym<sup>®</sup> 435 happened between the first and the last batch, this has no noticeable effect in the enzymatic activity during the consecutive uses. Therefore, yields of ~95% can be achieved in only one hour, and both biocatalysts can be successfully reused, at least, in 14 additional cycles.

**Table 1** Amount of biocatalyst employed in the first use and recovered along 15 batches and conversions obtained under optimum reaction conditions (5% w/w of Lipozyme<sup>®</sup> TL IM - 2.5% w/w of Novozym<sup>®</sup> 435, 70 °C, 21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates)

Uses	Conversion (%)		Amount of biocatalyst (g)	
	Lipozyme <sup>®</sup> TL IM	Novozym <sup>®</sup> 435	Lipozyme <sup>®</sup> TL IM	Novozym <sup>®</sup> 435
1 <sup>st</sup>	93.6 ± 0.9	96.3 ± 1.0	1.0004 ± 1.10 <sup>-4</sup>	0.5001 ± 1×10 <sup>-4</sup>
5 <sup>th</sup>	94.4 ± 2.0	96.0 ± 1.4	0.9867 ± 2.10 <sup>-3</sup>	0.5081 ± 9×10 <sup>-3</sup>
10 <sup>th</sup>	94.2 ± 1.6	95.5 ± 1.3	0.9991 ± 7.10 <sup>-3</sup>	0.4919 ± 3×10 <sup>-3</sup>
15 <sup>th</sup>	93.4 ± 1.4	95.0 ± 0.6	0.9044 ± 8.10 <sup>-3</sup>	0.4863 ± 6×10 <sup>-3</sup>

These results are in agreement with the ones obtained by Dianóczy et al. [29], where Novozym<sup>®</sup> 435 was successfully reused up to 10 times in the synthesis of conjugated linoleic acid esters under similar conditions without significant loss of the biocatalyst mass



1  
2  
3 or activity. On the other hand, Martins et al. [24] has reported the loss of activity of both  
4 biocatalysts after the first (Lipozyme<sup>®</sup> TL IM) and the 7<sup>th</sup> (Novozym<sup>®</sup> 435) batch when  
5 used in a solvent-free process to obtain isobutyl propionate, putting into evidence that long  
6 chained reagents helps to preserve the lipase-based catalysts' activity.  
7  
8

#### 9 **3.1.4 Process productivity**

10  
11 Considering that the heterogeneous process under study is carried out under solvent-free  
12 conditions, process productivity has been calculated as the amount of product obtained per  
13 gram of biocatalyst and hour by using the conversions attained in the first use of the  
14 commercial derivatives during the reusability test under best reaction conditions (Table 1).  
15 According to this, the use of 1 g (5% w/w) of Lipozyme<sup>®</sup> TL IM leads to productivities of  
16 18.7 g<sub>product</sub>/(g<sub>biocatalyst</sub>·h) while process productivity when using 0.5 g (2.5% w/w) of  
17 Novozym<sup>®</sup> 435 is of 38.5 g<sub>product</sub>/(g<sub>biocatalyst</sub>·h), ~2 times higher than TII derivative. These  
18 results are in accordance with the ones observed in Sections 3.1.1 and 3.1.2, making  
19 evident again that the characteristics of Novozym<sup>®</sup> 435 are better to the esterification  
20 process of such hydrophobic reagents.  
21  
22  
23  
24  
25

#### 26 **3.2 Biocatalyst election based in cost estimation**

27  
28 In view of the foregoing it can be concluded that CalB biocatalyst is a more adequate  
29 candidate for spermaceti production from a technical point of view. Still, TII derivative  
30 shows an acceptable synthetic activity. Considering that the industrial viability of a  
31 biocatalytic pathway lies in its economic profitability, a cost assessment of the optimized  
32 production of spermaceti with Lipozyme<sup>®</sup> TL IM and Novozym<sup>®</sup> 435 has been also  
33 performed. With this aim, only direct operation costs have been taken into account, namely  
34 the raw materials (substrates and immobilized derivatives) and utilities (heating, nitrogen,  
35 vacuum and stirring) needs, as they will be the only ones affected by the biocatalyst  
36 selected. Capital investment, fixed and indirect operation costs have not been considered  
37 because they are essentially the same in these two processes.  
38  
39  
40  
41

42 In this respect, Table 2 gathers the prices of the raw materials and their supplier. It must be  
43 noted that none of the suppliers of the reagents used sells both of the biocatalysts, so the  
44 authors contacted privately Novozymes Spain S.A., which gently provided the guide prices  
45 for Lipozyme<sup>®</sup> TL IM and Novozym<sup>®</sup> 435 collected in Table 2. Besides, Table 3 shows the  
46 power consumption of the equipment employed during the reactions, as described in  
47 Materials and Methods: a thermostatic bath, reactor's controller and stirrer, a N<sub>2</sub> generator  
48 and a vacuum pump. In order to avoid calculations relying on consumption data furnished  
49 by the suppliers of the equipment and hypothetical cost determinations, real-time  
50 measurements of the current intensity were carried out by means of a clamp meter. The  
51 energy demanded for the start-up of the system and the maintenance of the operating  
52 conditions has been considered, assuming that the average tension at the terminals of the  
53  
54  
55  
56  
57  
58  
59  
60

equipments is 220 V. In the case of the thermostatic bath, 10 min are required in continuous operation to reach the temperature set, 70 °C, a time used to melt and homogenize the reaction mixture. After that, temperature's controller starts operating in an auto on/off mode, and as a result, its power consumption diminishes. Regarding the N<sub>2</sub> generator, the device requires 35 min to produce enough pressure (6701.3 kPa) to guarantee that an appropriate inert atmosphere is supplied for one hour; then, the generator reconnects for 15 min to raise the pressure, allowing, again, to one hour of autonomy. Given that electricity's price at the moment when measurements were made was 0.2086 €/kWh [30], a realistic power cost was estimated for the enzyme-based synthesis at laboratory scale (Table 3).

**Table 2** Costs and providers of the substrates and biocatalysts used

Materials	Price (€/kg)	Supplier
Novozym <sup>®</sup> 435	1300	Novozymes [10] (Gifts, prices communicated personally)
Lipozyme <sup>®</sup> TL IM	70	
Cetyl alcohol	41.7	Aldrich (H6800) [31]
Lauric acid	34.2	Acros Organics (10398103) [32]
Myristic acid	37.4	Sigma-Aldrich (70082) [33]
Palmitic acid	42	Aldrich (W283207) [34]
Stearic acid	42	Aldrich (W303518) [35]

**Table 3** Power consumption and cost of the manufacturing utilities

Equipment	Specifications	Current intensity (A)	Power (W)	Cost (€/min)
Reactor stirrer (Parr A1120HC)	Magnetic drive Stirrer motor 1/8 HP variable speed 4-Blade turbine type impeller, 35 mm			
Reactor controller (Parr 4875)	Power controller for handling heating, cooling, safety, and motor control devices.	0.3	66	$0.2 \times 10^{-3}$
Vacuum pump (KNF N816.3KN18)	Max. permissible operating pressure: 0.5 bar Delivery rate: 16 L/min (atm. pressure)			
N <sub>2</sub> generator (Parker 3848 NitroFlow Lab)	Max. flow rate: 32 L/min	<i>Initial generation and maintenance: 4.5</i>	990	$3.4 \times 10^{-3}$
Reactor's heated circulating bath (PolyScience)	Max. capacity: 6 L Flow rate: 7-15 L/min Heater wattage: 1000 W	<i>Initial: 8.5 Maintenance: 2.4</i>	1870 527	$6.5 \times 10^{-3}$ $1.8 \times 10^{-3}$

Once obtained the unit costs for raw materials and process equipment, direct operation cost has been weighted for the previously described optimized production of spermaceti with the Tll and CalB commercial derivatives. The following calculation bases have been established for this purpose: (1) 20 g of final product (in accordance with the scale used during the experimentation), (2) the minimum operation time needed to achieve a conversion of 98%. Putting into evidence, once again, the considerable differences in the activity of these biocatalysts, these processing times are quite unequal between them. Thereby, 103 min are required for Lipozyme<sup>®</sup> TL IM and 51 min for Novozym<sup>®</sup> 435, although the double amount of biocatalyst is used in the first case (5% and 2.5% w/w referred to substrates, respectively). Results are presented in Table 4, where the following cost items have been compiled:

1. Substrates cost, which is the same for both processes and is based on the percentage composition of natural spermaceti.
2. Biocatalyst cost, taking into account the optimum concentration of each one and their reusability: 15 batches (minimum) without an apparent loss of activity.
3. Power cost related to the thermostatic bath employed for heating the glass-jacketed reactor's vessel during the conditioning of the reagents (10 min) and the process time.
4. Power cost due to the reactor's controller and stirrer, as well as to the vacuum pump, devices that were connected together to the electrical outlet.
5. Power cost resulting from using the N<sub>2</sub> generator. It is noteworthy that after initial generation (35 min), a single extra run is necessary for the reaction catalysed by Lipozyme<sup>®</sup> TL IM, as it requires more than an hour (103 min) to reach a conversion of 98%. On the other hand, no further launching of this system is needed for completing the synthesis with Novozym<sup>®</sup> 435.

Even though it is obvious that the direct operation expenses proposed in Table 4 are far distant from the final cost of the whole process, in the last row these processing costs are given as percentage relatives, assigning 100% to the highest value: Lipozyme<sup>®</sup> TL IM catalysed synthesis. Surprisingly, direct operation costs for Novozym<sup>®</sup> 435 are a ~10% lower in spite of the fact that this biocatalyst is almost 20 times more expensive than Tll derivative. This might be explained by the great activity of the commercial immobilized CalB, which contributes to the economic profitability of the biosynthesis in two ways: first, lower concentrations of biocatalyst are necessary and, in addition, it entails shorter operation time and energy consumption.

**Table 4** Broken down cost of the synthesis of 20 g of spermaceti at laboratory scale

Cost item	Cost (€/20 g spermaceti)	
	Novozym <sup>®</sup> 435	Lipozyme TL IM
Substrates: 20 g		
Cetyl alcohol: 10.11 g	0.42	0.42
Lauric acid: 1.51 g	0.052	0.052
Myristic acid: 3.73 g	0.14	0.14
Palmitic acid: 3.83 g	0.16	0.16
Stearic acid: 0.81 g	0.034	0.034
Immobilized lipase Reuse (15 times)	(0.5 g) 0.043	(1 g) 0.0047
Reactor heated circulating bath (70 °C)	Initial (10 min): 0.065 Maintenance (51 min): 0.09	Initial (10 min): 0.065 Maintenance (103 min): 0.18
Reactor controller and stirrer (350 rpm) and vacuum pump (21.3 kPa)	(51 min) 0.01	(103 min) 0.02
N <sub>2</sub> generator (54 L/h)	Initial (35 min): 0.12 Maintenance: 0	Initial (35 min): 0.12 Maintenance (15 min): 0.051
<b>Total direct operation costs</b>	<b>1.13</b>	<b>1.25</b>
<b>Percent direct operation costs (%)</b>	<b>(90.4%)</b>	<b>(100%)</b>

Figure 4 represents all those elements grouped by categories, making patent that the contribution of the biocatalyst to the global cost is little: 0.4% for Lipozyme<sup>®</sup> TL IM and 3.8% for Novozym<sup>®</sup> 435. These values are clearly below the 5% proposed by Tufvesson et al. [21] as the maximum weight of biocatalyst cost for the enzymatic production of specialty chemicals, such as cosmetic ingredients. Needless to say that these results are a consequence of the good operational stability of these immobilized derivatives, which, as it has been proved experimentally, can be reused 14 times at least. As shown by this figure, direct operation costs are mainly dependent on the substrates cost, which represent more than a half of the process' economic requirements. This is primarily caused by the fact that these reagents have been acquired from companies that typically provides chemical products at small-scale, for analysis or research purposes.

(Figure 4)

If the scale-up of the process is desired, substrate's purchase must inevitably turn to bulk raw materials suppliers, with whom discounts might even be negotiated if these substances are bought in large quantities [36]. Some examples of these products' prices

and the minimum order quantity for large scale production are given in Table 5, where the purveyors have been selected taking into consideration that they are able to provide cosmetic grade reagents. Comparing these prices with the ones furnished in Table 2, it can be noticed that they can be up to 40 times less expensive when purchased at larger amounts. If calculations for 20 g of spermaceti are made using these new data, direct operation costs would be dramatically reduced: 0.35 € and 0.47 € for CalB and Tll derivatives respectively. This supposes a 25% of saving if Novozym<sup>®</sup> 435 is used instead Lipozyme<sup>®</sup> TL IM. In other words, for large-scale purposes the differences between the two processes would be enhanced by using bulk substrates, making even clearer that, still being more expensive, it is more profitable to choose a biocatalyst with better technical features (higher activity) if reused.

**Table 5** Bulk raw materials for the synthesis of spermaceti at large scale

Materials	Price (€/kg)	Minimum Order	Company
Cetyl alcohol (98%)	0.85-2.54	1000 kg	J.C. Enterprises. Maharashtra, India [37]
Lauric acid (99%)	1.69	100 kg	Guangzhou Kaoking Chemical Co. Ltd. Guangdong, China [38, 39]
Myristic acid (99.9%)	2.34	200 kg	
Palmitic acid (99%)	0.85-8.5	1 kg	Guangzhou Yiming Chemical Materials Co. Ltd. Guangdong, China [40]
Stearic acid (99.98%)	1.27	200 kg	Guangzhou Kaoking Chemical Co., Ltd. Guangdong, China [41]

In Figure 5 it can be observed that the main contribution to the direct operating costs is due to power consumption when bulk raw materials are used, which supposes now a 94% of the total for Lipozyme<sup>®</sup> TL IM and an 81% for Novozym<sup>®</sup> 435. Energy cost may be still lessen, as calculations has been made considering real-time energy consumption at laboratory scale, being all the equipment connected to a domestic electrical network with non-industrial market prices. In this sense, electricity charges for a medium size industry can suppose half of the household cost [42], but even in this case, power investment would be the major contribution to direct operation cost.

(Figure 5)

So, with a view to save energy at larger production scale, the percentage of the global power cost caused by each of the equipment used has been calculated (Figure 6). These results make patent that in both processes most of the energy is consumed to heat the circulating water going through the reactor's jacket and keeping operating temperature constant. Unfortunately, this parameter is largely restricted to the substrates' melting

1  
2  
3 point in a solvent-free system, but, as it has been pointed out in Section 3.1.2., it is still  
4 possible to carry out the synthesis of spermaceti at 60 °C, even if this entails a longer  
5 conditioning of reaction mixture and, possibly, only a slight benefit for such short  
6 reaction times. In spite of that, it is well known that that the economic inversion in  
7 operating utilities is reduced when increasing the size of a production plant [43, 44], and  
8 particularly in the case of the plant's heating energy requirements [45]. The scale-up of  
9 the system would be necessary for an accurate evaluation of the energy expenses of both  
10 biocatalytic processes.  
11  
12  
13

14 (Figure 6)  
15  
16  
17  
18

#### 19 4. Conclusions

20  
21 The production of a mixture of cetyl esters similar to natural spermaceti has been  
22 optimized by using the commercial biocatalysts Lipozyme® TL IM and Novozym® 435,  
23 During this study it has been put into evidence the great activity of Novozym® 435,  
24 which can double Lipozyme® TL IM's process productivity under optimal conditions.  
25 Nevertheless, the optimized process with both biocatalysts led to conversions higher  
26 than 97.5%, making patent that a preliminary economic assessment is needed in order to  
27 choose the best immobilized derivative. In this sense, cost analysis has pointed out that  
28 the better technical features of Novozym® 435 can compensate its high market price and  
29 entail direct operation costs lower than the 20 times cheaper Lipozyme® TL IM if  
30 reused. Although an appropriate evaluation of large scale energy consumptions is  
31 necessary, results by using bulk raw material prices lead to think that biocatalyst  
32 investment may not constitute the main economic requirement of the enzymatic process.  
33  
34  
35  
36  
37  
38  
39

40 **Acknowledgements:** This work has been funded by the Spanish Ministry of Science,  
41 Innovation and Universities (CTQ2015-66723-R) and the European Commission  
42 (FEDER/ERDF). M. Serrano-Arnaldos and S. Ortega-Requena were beneficiaries of a  
43 FPI pre-doctoral scholarship from the Spanish Ministry of Economy and  
44 Competitiveness (MINECO) and a Torres Quevedo grant, respectively. We wish to  
45 acknowledge D. Ramiro Martínez Gutiérrez (Novozymes Spain S.A.) who kindly  
46 supplied the biocatalysts.  
47  
48  
49  
50  
51

52 **Conflict of Interest:** The authors declare that they have no conflict of interest.  
53  
54  
55  
56  
57  
58  
59  
60



## References

1. Antonopoulou I, Varriale S, Topakas E, et al (2016) Enzymatic synthesis of bioactive compounds with high potential for cosmeceutical application. *Appl Microbiol Biotechnol* 100:6519–6543. <https://doi.org/10.1007/s00253-016-7647-9>
2. Ye R, Hayes DG, Burton R, et al (2016) Solvent-Free Lipase-Catalyzed Synthesis of Technical-Grade Sugar Esters and Evaluation of Their Physicochemical and Bioactive Properties. *Catalysts* 6:78. <https://doi.org/10.3390/catal6060078>
3. Yvergnaux F (2017) Lipases: particularly effective biocatalysts for cosmetic active ingredients. *Ocl-Oilseeds Fats Crops Lipids* 24:D408. <https://doi.org/10.1051/ocl/2017013>
4. Grunwald P (2014) *Industrial Biocatalysis*. CRC Press, Boca Raton (FL)
5. Hilterhaus L, Liese A, Kettling U, Antranikian G (2016) *Applied Biocatalysis: From Fundamental Science to Industrial Applications*. John Wiley & Sons
6. Daiha K de G, Angeli R, de Oliveira SD, Almeida RV (2015) Are Lipases Still Important Biocatalysts? A Study of Scientific Publications and Patents for Technological Forecasting. *Plos One* 10:e0131624. <https://doi.org/10.1371/journal.pone.0131624>
7. Borole AP, Davison BH (2007) Techno-economic analysis of biocatalytic processes for production of alkene epoxides. *Appl Biochem Biotechnol* 137–140:437–449. <https://doi.org/10.1007/s12010-007-9070-2>
8. Mustafa A, Karmali A, Abdelmoez W (2016) Optimisation and economic assessment of lipase-catalysed production of monoesters using *Rhizomucor miehei* lipase in a solvent-free system. *J Clean Prod* 137:953–964. <https://doi.org/10.1016/j.jclepro.2016.07.056>
9. Fernández-Lafuente R (2010) Lipase from *Thermomyces lanuginosus*: Uses and prospects as an industrial biocatalyst. *J Mol Catal B-Enzym* 62:197–212. <https://doi.org/10.1016/j.molcatb.2009.11.010>
10. Novozymes A/S Biocatalysis for a sustainable pharma future - Novozymes. <https://www.novozymes.com/en/advance-your-business/pharma>. Accessed 9 Jan 2019
11. Bansode SR, Hardikar MA, Rathod VK (2017) Evaluation of reaction parameters and kinetic modelling for Novozym 435 catalysed synthesis of isoamyl butyrate. *J Chem Technol Biotechnol* 92:1306–1314. <https://doi.org/10.1002/jctb.5125>
12. Koutinas M, Yiangou C, Osório NM, et al (2018) Application of commercial and non-commercial immobilized lipases for biocatalytic production of ethyl lactate in



- organic solvents. *Bioresour Technol* 247:496–503. <https://doi.org/10.1016/j.biortech.2017.09.130>
13. Madarasz J, Nemeth D, Bakos J, et al (2015) Solvent-free enzymatic process for biolubricant production in continuous microfluidic reactor. *J Clean Prod* 93:140–144. <https://doi.org/10.1016/j.jclepro.2015.01.028>
14. Cirillo NA, Quirrenbach CG, Corazza ML, Pedersen Voll FA (2018) Enzymatic kinetics of cetyl palmitate synthesis in a solvent-free system. *Biochem Eng J* 137:116–124. <https://doi.org/10.1016/j.bej.2018.05.021>
15. Lima LCD, Peres DGC, Mendes AA (2018) Kinetic and thermodynamic studies on the enzymatic synthesis of wax ester catalyzed by lipase immobilized on glutaraldehyde-activated rice husk particles. *Bioprocess Biosyst Eng* 41:991–1002. <https://doi.org/10.1007/s00449-018-1929-9>
16. Montiel MC, Serrano M, Máximo MF, et al (2015) Synthesis of cetyl ricinoleate catalyzed by immobilized Lipozyme® CalB lipase in a solvent-free system. *Catal Today* 255:49–53. <https://doi.org/10.1016/j.cattod.2014.09.015>
17. Serrano-Arnaldos M, Maximo-Martin MF, Montiel-Morte MC, et al (2016) Solvent-free enzymatic production of high quality cetyl esters. *Bioprocess Biosyst Eng* 39:641–9. <https://doi.org/10.1007/s00449-016-1545-5>
18. Veit T (2004) Biocatalysis for the production of cosmetic ingredients. *Eng Life Sci* 4:508–511. <https://doi.org/10.1002/elsc.200402148>
19. Wellendorf M (1963) Composition of Spermaceti. *Nature* 198:1086–1087. <https://doi.org/10.1038/1981086b0>
20. Serrano-Arnaldos M, Bastida J, Máximo F, et al (2018) One-Step Solvent-Free Production of a Spermaceti Analogue Using Commercial Immobilized Lipases. *ChemistrySelect* 3:748–752. <https://doi.org/10.1002/slct.201702332>
21. Tufvesson P, Lima-Ramos J, Nordblad M, Woodley JM (2011) Guidelines and Cost Analysis for Catalyst Production in Biocatalytic Processes. *Org Process Res Dev* 15:266–274. <https://doi.org/10.1021/op1002165>
22. Ortega-Requena S, Bódalo-Santoyo A, Bastida-Rodríguez J, et al (2014) Optimized enzymatic synthesis of the food additive polyglycerol polyricinoleate (PGPR) using Novozym® 435 in a solvent free system. *Biochem Eng J* 84:91–97. <https://doi.org/10.1016/j.bej.2014.01.003>
23. ASTM D974–02e1 (2002) Standard test method for acid and base number by color-indicator titration. ASTM International, West Conshohocken, PA
24. Martins AB, da Silva AM, Schein MF, et al (2014) Comparison of the performance of commercial immobilized lipases in the synthesis of different flavor

- 1  
2  
3 esters. J Mol Catal B-Enzym 105:18–25.  
4 <https://doi.org/10.1016/j.molcatb.2014.03.021>  
5  
6  
7 25. Aguiéiras ECG, Veloso CO, Bevilaqua JV, et al (2011) Estolides Synthesis  
8 Catalyzed by Immobilized Lipases. Enzyme Res 2011:1–7.  
9 <https://doi.org/10.4061/2011/432746>  
10  
11  
12 26. Anderson EM, Larsson KM, Kirk O (1998) One Biocatalyst–Many Applications:  
13 The Use of *Candida Antarctica* B-Lipase in Organic Synthesis. Biocatal  
14 Biotransformation 16:181–204. <https://doi.org/10.3109/10242429809003198>  
15  
16  
17 27. Basri M, Kassim MA, Mohamad R, Ariff AB (2013) Optimization and kinetic  
18 study on the synthesis of palm oil ester using Lipozyme TL IM. J Mol Catal B  
19 Enzym 85–86:214–219. <https://doi.org/10.1016/j.molcatb.2012.09.013>  
20  
21  
22 28. Choi N, Kim Y, Lee J-S, et al (2016) Synthesis of Fatty Acid Ethyl Ester from  
23 Acid Oil in a Continuous Reactor via an Enzymatic Transesterification. J Am Oil  
24 Chem Soc 93:311–318. <https://doi.org/10.1007/s11746-016-2786-9>  
25  
26  
27 29. Dianóczki C, Recseg K, Kővári K, Poppe L (2007) Convenient enzymatic  
28 preparation of conjugated linoleic acid alkyl esters with C6–C22 alcohols. J Mol  
29 Catal B Enzym 45:45–49. <https://doi.org/10.1016/j.molcatb.2006.11.005>  
30  
31  
32 30. Comisión Nacional de los Mercados y la Competencia (2016) Boletín de  
33 indicadores eléctricos 2016  
34  
35 31. Cetyl alcohol, Sigma-Aldrich.  
36 [https://www.sigmaaldrich.com/catalog/search?term=Cetyl+alcohol&interface=Pro](https://www.sigmaaldrich.com/catalog/search?term=Cetyl+alcohol&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
37 [duct%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&f](https://www.sigmaaldrich.com/catalog/search?term=Cetyl+alcohol&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
38 [ocus=productN=0%20220003048%20219853286%20219853112](https://www.sigmaaldrich.com/catalog/search?term=Cetyl+alcohol&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112). Accessed 3 Sep  
39 2018  
40  
41  
42 32. Lauric acid, 99%, ACROS Organics - Organic Building Blocks Chemicals.  
43 <https://www.fishersci.es/shop/products/lauric-acid-99-acros-organics-5/p-3736787>.  
44 Accessed 3 Sep 2018  
45  
46  
47 33. Myristic acid, Sigma-Aldrich.  
48 [https://www.sigmaaldrich.com/catalog/search?term=Myristic+acid&interface=Pro](https://www.sigmaaldrich.com/catalog/search?term=Myristic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
49 [duct%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&f](https://www.sigmaaldrich.com/catalog/search?term=Myristic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
50 [ocus=productN=0%20220003048%20219853286%20219853112](https://www.sigmaaldrich.com/catalog/search?term=Myristic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112). Accessed 3 Sep  
51 2018  
52  
53  
54 34. Palmitic acid, Sigma-Aldrich.  
55 [https://www.sigmaaldrich.com/catalog/search?term=Palmitic+acid&interface=Pro](https://www.sigmaaldrich.com/catalog/search?term=Palmitic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
56 [duct%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&f](https://www.sigmaaldrich.com/catalog/search?term=Palmitic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112)  
57 [ocus=productN=0%20220003048%20219853286%20219853112](https://www.sigmaaldrich.com/catalog/search?term=Palmitic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112). Accessed 3 Sep  
58 2018  
59  
60

- 1  
2  
3 35. Stearic acid, Sigma-Aldrich.  
4 <https://www.sigmaaldrich.com/catalog/search?term=Stearic+acid&interface=Product%20Name&N=0&mode=mode%20matchpartialmax&lang=es&region=ES&focus=productN=0%20220003048%20219853286%20219853112>. Accessed 3 Sep  
5  
6  
7  
8 2018  
9
- 10  
11 36. Brennan DJ (2004) Developing a Process Industry Culture. Morgan Printing
- 12  
13 37. Cetyl Alcohol Fatty Alcohol C1698 Supplier From India.  
14 [https://www.alibaba.com/product-detail/Cetyl-Alcohol-Fatty-Alcohol-C1698-](https://www.alibaba.com/product-detail/Cetyl-Alcohol-Fatty-Alcohol-C1698-Supplier_146547456.html)  
15 [Supplier\\_146547456.html](https://www.alibaba.com/product-detail/Cetyl-Alcohol-Fatty-Alcohol-C1698-Supplier_146547456.html). Accessed 6 Sep 2018  
16
- 17  
18 38. High quality cosmetic material Lauric acid 99%, Product Details from Guangzhou  
19 Kaoking Chemical Co., Ltd. [https://kao.en.alibaba.com/product/60399285976-](https://kao.en.alibaba.com/product/60399285976-802337594/high_quality_cosmetic_material_Lauric_acid_99_price.html?spm=a2700.8304367.prewdfa4cf.2.207346f37aIcYF)  
20 [802337594/high\\_quality\\_cosmetic\\_material\\_Lauric\\_acid\\_99\\_price.html?spm=a27](https://kao.en.alibaba.com/product/60399285976-802337594/high_quality_cosmetic_material_Lauric_acid_99_price.html?spm=a2700.8304367.prewdfa4cf.2.207346f37aIcYF)  
21 [00.8304367.prewdfa4cf.2.207346f37aIcYF](https://kao.en.alibaba.com/product/60399285976-802337594/high_quality_cosmetic_material_Lauric_acid_99_price.html?spm=a2700.8304367.prewdfa4cf.2.207346f37aIcYF). Accessed 6 Sep 2018  
22
- 23  
24 39. Myristic acid. [https://wholesaler.alibaba.com/product-detail/Soap-Detergent-](https://wholesaler.alibaba.com/product-detail/Soap-Detergent-Cosmetic-Surfactant-myristic-acid_60703624293.html)  
25 [Cosmetic-Surfactant-myristic-acid\\_60703624293.html](https://wholesaler.alibaba.com/product-detail/Soap-Detergent-Cosmetic-Surfactant-myristic-acid_60703624293.html). Accessed 7 Sep 2018  
26
- 27  
28 40. Chemical Raw Material Palmitic Acid For Cosmetics.  
29 [https://www.alibaba.com/product-detail/chemical-raw-material-palmitic-acid-](https://www.alibaba.com/product-detail/chemical-raw-material-palmitic-acid-for_60148565501.html?spm=a2700.7724838.2017115.109.4a105a41770i9)  
30 [for\\_60148565501.html?spm=a2700.7724838.2017115.109.4a105a41770i9](https://www.alibaba.com/product-detail/chemical-raw-material-palmitic-acid-for_60148565501.html?spm=a2700.7724838.2017115.109.4a105a41770i9).  
31 Accessed 6 Sep 2018  
32
- 33  
34 41. High quality cosmetic material Stearic acid, Product Details from Guangzhou  
35 Kaoking Chemical Co., Ltd. [https://kao.en.alibaba.com/product/60439480037-](https://kao.en.alibaba.com/product/60439480037-802394266/high_quality_cosemantic_material_Stearic_acid.html?spm=a2700.8304367.prewdfa4cf.2.28c79411Vzc7eM)  
36 [802394266/high\\_quality\\_cosemantic\\_material\\_Stearic\\_acid.html?spm=a2700.83043](https://kao.en.alibaba.com/product/60439480037-802394266/high_quality_cosemantic_material_Stearic_acid.html?spm=a2700.8304367.prewdfa4cf.2.28c79411Vzc7eM)  
37 [67.prewdfa4cf.2.28c79411Vzc7eM](https://kao.en.alibaba.com/product/60439480037-802394266/high_quality_cosemantic_material_Stearic_acid.html?spm=a2700.8304367.prewdfa4cf.2.28c79411Vzc7eM). Accessed 6 Sep 2018  
38
- 39  
40 42. Eurostat - Tables, Graphs and Maps Interface (TGM).  
41 [http://ec.europa.eu/eurostat/tgm/refreshTableAction.do?tab=table&plugin=1&pcod](http://ec.europa.eu/eurostat/tgm/refreshTableAction.do?tab=table&plugin=1&pcode=ten00117&language=en)  
42 [e=ten00117&language=en](http://ec.europa.eu/eurostat/tgm/refreshTableAction.do?tab=table&plugin=1&pcode=ten00117&language=en). Accessed 13 Dec 2018  
43
- 44  
45 43. Jiménez-González C, Constable DJC (2011) Green Chemistry and Engineering: A  
46 Practical Design Approach. John Wiley & Sons
- 47  
48 44. Vogel GH (2014) Production cost estimation. Ullmanns Encycl. Ind. Chem.
- 49  
50 45. Bermúdez JM, Beneroso D, Rey-Raap N, et al (2015) Energy consumption  
51 estimation in the scaling-up of microwave heating processes. Chem Eng Process  
52 Process Intensif 95:1–8. <https://doi.org/10.1016/j.cep.2015.05.001>  
53  
54  
55  
56  
57  
58  
59  
60

**Figure captions**

**Fig. 1** Cetyl laurate (a), myristate (b), palmitate (c) and stearate (d) solvent-free synthesis using different quantities of Lipozyme<sup>®</sup> TL IM (70 °C, 21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates, 1:1 molar ratio)

**Fig. 2** Cetyl laurate (a), myristate (b), palmitate (c) and stearate (d) solvent-free synthesis using different quantities of Novozym<sup>®</sup> 435 (70 °C, 21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates, 1:1 molar ratio)

**Fig. 3** Spermaceti solvent-free synthesis at different temperatures using 5% w/w of Lipozyme<sup>®</sup> TL IM (a) and 2.5% w/w of Novozym<sup>®</sup> 435 (b) (21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates)

**Fig. 4** Contribution of the biocatalyst, substrates and power to the direct operation costs of the synthesis of 20 g of spermaceti under optimal conditions at laboratory scale

**Fig. 5** Contribution of the biocatalyst, substrates and power to the direct operation costs of the synthesis of 20 g of spermaceti under optimal conditions when bulk raw material prices are considered

**Fig. 6** Contribution of the different equipment to the energy costs while synthesizing 20 g of spermaceti




Fig. 1 Cetyl laurate (a), myristate (b), palmitate (c) and stearate (d) solvent-free synthesis using different quantities of Lipozyme® TL IM (70 °C, 21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates, 1:1 molar ratio)

173x181mm (300 x 300 DPI)




Fig. 2 Cetyl laurate (a), myristate (b), palmitate (c) and stearate (d) solvent-free synthesis using different quantities of Novozym® 435 (70 °C, 21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates, 1:1 molar ratio)

172x181mm (300 x 300 DPI)




Fig. 3 Spermacti solvent-free synthesis at different temperatures using 5% w/w of Lipozyme® TL IM (a) and 2.5% w/w of Novozym® 435 (b) (21.3 kPa, 54 L/h N<sub>2</sub>, 350 rpm, 20 g of substrates)

173x106mm (300 x 300 DPI)




Fig. 4 Contribution of the biocatalyst, substrates and power to the direct operation costs of the synthesis of 20 g of spermaceti under optimal conditions at laboratory scale

163x85mm (300 x 300 DPI)






Fig. 5 Contribution of the biocatalyst, substrates and power to the direct operation costs of the synthesis of 20 g of spermaceti under optimal conditions when bulk raw material prices are considered

156x85mm (300 x 300 DPI)




Fig. 6 Contribution of the different equipment to the energy costs while synthesizing 20 g of spermaceti

144x112mm (300 x 300 DPI)