



RESEARCH COMMUNICATION

Helminths in a giraffe (*Giraffa camelopardalis giraffa*) from a zoo in Spain

M.M. GARIJO, J.M. ORTIZ and M.R. RUIZ DE IBÁÑEZ

Department of Parasitology, Faculty of Veterinary Science, University of Murcia
30100 Espinardo, Murcia, Spain

ABSTRACT

GARIJO, M.M., ORTIZ, J.M. & RUIZ DE YBÁÑEZ, M.R. 2004. Helminths in a giraffe (*Giraffa camelopardalis giraffa*) from a zoo in Spain. *Onderstepoort Journal of Veterinary Research*, 71:153–156

A pregnant female Cape giraffe (*Giraffa camelopardalis giraffa*) died from an unknown cause in the Aitana Zoo, Alicante, Spain. Neither clinical signs nor macroscopic lesions were observed at necropsy. The alimentary tract was removed and examined for parasites. A total of 2724 nematodes were found, including *Camelostromylus mentulatus*, *Trichostrongylus axei*, *Ostertagia ostertagi*, *Teladorsagia circumcincta*, *Teladorsagia trifurcata*, *Marshallagia marshalli*, *Trichostrongylus vitrinus*, *Trichostrongylus colubriformis*, *Spiculopteragia asymmetrica* and *Trichuris giraffae*. Only *T. giraffae* and *C. mentulatus* have been previously reported from giraffes. The other nematodes are common in mouflons, fallow and red deer, which can usually be found in the same paddock as the giraffes in Aitana Zoo. Although its occurrence is unusual in this host, *C. mentulatus* was the most abundant nematode in our giraffe. This parasite has been related to disease, and even death, in several wild ruminants.

Keywords: Gastrointestinal nematodes, *Giraffa camelopardalis giraffa*, helminths, Spain

INTRODUCTION

The interest in the parasitoses of wildlife must be related both to the increase in the number of game reserves and to the potential transmission of these parasites to domestic animals. Most of the available information is related to Artiodactyla, but little is known about that concerning giraffes (Round 1968; Pester & Laurence 1974; Boomker, Horak & De Vos 1986; Krecek, Boomker, Penzhorn & Scheepers 1990; Fukomoto, Uchida, Ohbayashi, Ikebe & Sasano 1996). The study of animals that have died in accidents or from other causes provides useful opportunities for advancing the knowledge of the parasites of artiodactylids. This is of particular impor-

tance in the case of the giraffe, a species which is considered to be significant in European zoos. This study provides data of parasites found in this species living in Spain and describes the presence of worms not usually found in giraffes.

MATERIALS AND METHODS

During November 2002, a 4-year-old pregnant (3rd month of gestation) female Cape giraffe (*Giraffa camelopardalis giraffa*) died from an unknown cause in the Aitana Zoo, Alicante, in the southeast of Spain. No clinical signs were observed. The feeding of the giraffes in the zoo is based on *Quercus rotundifolia*, *Daphne gnidium* and *Pinus halepensis* and lucerne is supplied in limited quantities. Both natural and artificial water resources are available to all animals.

The dead animal was sent to the Veterinary Science Faculty, University of Murcia, where it was necropsied. No macroscopic lesions were observed. The animal was in good condition as adipose tissue was present around the small and large intestines and the kidneys. The alimentary tract was removed and examined for parasites according to the method described by Ortiz, Ruiz de Ybáñez, Garijo, Goyena, Espeso, Abaigar & Cano 2001. In brief, the abomasum, small and large intestines were processed separately by scraping the mucosal surfaces and washing the material so gained as well as the gut contents through a series of mesh screens, the final one of which had apertures 0.3 mm in size. The material remaining on the sieves was collected separately and fixed in 10% formalin.

Each sample, divided into smaller portions, was examined under a stereoscopic microscope and the nematodes present were removed. Adult male parasites were cleared in lactophenol for 48 h, and identified with the aid of the keys of Skrjabin, Shikhobalova, Schulz, Popova, Boev & Delyaure (1961) and Durette-Desset (1983, 1989). Representative specimens of the nematodes recovered are deposited in the Faculty of Veterinary Science, University of Murcia's parasite collection.

RESULTS AND DISCUSSION

A total of 2724 nematodes were found (1158 males and 1566 females). Ten nematode species were recovered and neither trematodes nor cestodes were found. The abomasum yielded a total of 1109 helminths (593 females and 516 males). The study of the males gave the following distribution:

<i>Camelostrongylus mentulatus</i>	284
<i>Trichostrongylus axei</i>	101
<i>Ostertagia ostertagi</i>	84
<i>Teladorsagia circumcincta</i>	35
<i>Marshallagia marshalli</i>	4
<i>Trichostrongylus vitrinus</i>	3
<i>Spiculopteragia asymmetrica</i>	3
<i>Teladorsagia trifurcata</i>	2

The small intestine harboured 1601 nematodes, 967 females and 634 males, comprising 587 *T. vitrinus* and 47 *Trichostrongylus colubriformis*. Fourteen worms were found in the large intestine (six females and eight males). All those males were identified as *Trichuris giraffae*.

Little information is available in the literature on the parasites of giraffes, and all refer to those recovered from animals living in National Parks. Krecek

et al. (1990), in a study on six animals in the Etosha National Park, Namibia, recorded parasite loads ranging from 18 to 531. These numbers are much lower than those obtained in the present work. It should be mentioned that keeping the animals in paddocks, as the giraffe in this study was, makes re-infection easier. On the other hand, Boomker *et al.* (1986) recorded 2621 and 19157 helminths, respectively, in two giraffes in the Kruger National Park, South Africa, a much larger number than recorded for the animals from Etosha. Krecek *et al.* (1990) attributed these differences to the low rainfall at Etosha National Park.

Of the helminths recovered in our study only *T. giraffae* and *C. mentulatus* are reported as occurring in giraffe (Round 1968; Fukumoto *et al.* 1996). The remaining species are common nematodes in other ruminant species but constitute the first records for giraffe.

Under natural conditions giraffes are exclusively browsers (Dorst & Dandelot 1972). However, the nematodes found in the giraffe in this study are considered to be mainly those of grazers. The manner in which the animals are maintained in the zoo is probably the reason for this. The zoo can be considered as a type of safari park as the animals are kept in very large paddocks. In addition, they can even change paddocks if they wish to do so. In this respect mouflons, *Ovis musimon*, are frequently found in the paddocks of other animal species. Restricting animals to game reserves and parks facilitates infections between different animal species, and, in addition, many helminths of antelope are not very host specific (Boomker *et al.* 1986).

Although the feeding habits of giraffes, in general, makes it difficult to acquire helminths of other ruminant species, Boomker *et al.* (1986) consider that they could acquire these when forced to graze. In the Aitana zoo lucerne is provided to the herbivores and is placed on the ground. This procedure could explain the presence of nematodes considered uncommon in giraffes, but which are frequently seen in other species in the zoo such as mouflons, fallow deer, *Dama dama* and red deer, *Cervus elaphus*.

Other gastrointestinal parasites previously recorded in giraffes, i.e. *Moniezia expansa* (Pester & Lawrence 1974), hydatid cysts (Kelly, Boray & Dixon 1968; Krecek *et al.* 1990), *Haemonchus mitchelli* (Sachs *et al.* 1973; Krecek *et al.* 1990), *Skrjabinema* spp. (Krecek *et al.* 1990) or *Parabronema skrjabini* (Boomker *et al.* 1986; Krecek *et al.* 1990), were not found in the present study.

SKRJABIN, K.I., SHIKHOBALOVA, N.P., SCHULZ, R.S., POPOVA, T.I., BOEV, S.N. & DELYAURE, S.L. 1961. *Key to parasitic nematodes, Vol. III. Strongylata*. Academy of Sciences of the U.S.S.R. Jerusalem: Israel Program for Scientific Translation.

WISSER, J., TSCHERNER, W. & JANTSCHKE, B. 2001. Infestation with *Camelostongylus mentulatus* in dorcas gazelles (*Gazella dorcas neglecta*) caused lethal abomasitis. *Verhandlungsberichte Erkrankungen Zootierie*, 40:81-86.