Environmental Microbiology (2022) 00(00), 00-00



doi:10.1111/1462-2920.15895

# Coenzyme B<sub>12</sub>-dependent and independent photoregulation of carotenogenesis across Myxococcales

Ricardo Pérez-Castaño <sup>(D, 1</sup> Eva Bastida-Martínez <sup>(D, 1</sup> Jesús Fernández-Zapata <sup>(D, 2</sup> María del Carmen Polanco <sup>(D, 1</sup> María Luisa Galbis-Martínez <sup>(D, 1</sup> Antonio A. Iniesta <sup>(D, 1</sup> Marta Fontes <sup>(D, 1</sup> S. Padmanabhan <sup>(D 2\*</sup> and Montserrat Elías-Arnanz <sup>(D) 1\*</sup>

<sup>1</sup>Departamento de Genética y Microbiología, Área de Genética (Unidad Asociada al IQFR-CSIC), Facultad de Biología, Universidad de Murcia, Murcia, 30100, Spain. <sup>2</sup>Instituto de Química Física "Rocasolano", Consejo Superior de Investigaciones Científicas, Madrid, 28006, Spain.

# Summary

Light-induced carotenogenesis in Myxococcus xanthus is controlled by the B<sub>12</sub>-based CarH repressor and photoreceptor, and by a separate intricate pathway involving singlet oxygen, the B<sub>12</sub>-independent CarH paralogue CarA and various other proteins, some eukaryotic-like. Whether other myxobacteria conserve these pathways and undergo photoregulated carotenogenesis is unknown. Here, comparative analyses across 27 Myxococcales genomes identified carotenogenic genes, albeit arranged differently, with carH often in their genomic vicinity, in all three Myxococcales suborders. However, CarA and its associated factors were found exclusively in suborder Cystobacterineae, with carA-carH invariably in tandem in a syntenic carotenogenic operon, except for Cystobacter/Melittangium, which lack CarA but retain all other factors. We experimentally show B<sub>12</sub>mediated photoregulated carotenogenesis in representative myxobacteria, and a remarkably plastic CarH operator design and DNA binding across Myxococcales. Unlike the two characterized CarH from other phyla, which are tetrameric, Cystobacter CarH (the first myxobacterial homologue amenable to analysis in vitro) is a dimer that combines direct CarH-like  $B_{12}$ -based photoregulation with CarA-like DNA binding and inhibition by an antirepressor. This study provides new molecular insights into  $B_{12}$ -dependent photoreceptors. It further establishes the  $B_{12}$ -dependent pathway for photoregulated carotenogenesis as broadly prevalent across myxobacteria and its evolution, exclusively in one suborder, into a parallel complex  $B_{12}$ -independent circuit.

# Introduction

Light directs many key biological processes but can also cause photooxidative stress by generating highly reactive oxygen species (ROS), like singlet oxygen  $({}^{1}O_{2})$ , which attack proteins, DNA and lipids (Ziegelhoffer and Donohue, 2009; Glaeser et al., 2011). Many organisms, including bacteria, combat photooxidative stress by inducing the synthesis of carotenoids, pigments that quench ROS (Armstrong, 1997; Ziegelhoffer and Donohue, 2009; Rodriguez-Concepción et al., 2018; Sandmann, 2019). How light is sensed and signalled to trigger carotenogenesis has been studied in depth in Myxococcus xanthus (Elías-Arnanz et al., 2011; Padmanabhan et al., 2021), a Gram-negative soil bacterium of the order Myxococcales, currently included within the class Deltaproteobacteria but it was recently proposed that this order be moved to a new and independent phylum with the name Myxococcata (Waite et al., 2020). Members of this order, the myxobacteria, form an important group of ubiguitous, predominantly aerobic, soil bacteria that share complex lifestyles and several traits typical of eukaryotes, such as multicellular development, biosynthesis of specialized lipids and steroids, social behaviour, kin recognition (the ability of an individual cell or organism to identify others as self-like), predation and motility (Muñoz-Dorado et al., 2016; Cao and Wall, 2019; Gallego-García et al., 2019; Hoshino and Gaucher, 2021). Given these eukaryotic-like traits in myxobacteria, the Syntrophy hypothesis envisages an ancient myxobacterium at the origin of eukaryotes (López-García and Moreira, 2020; Hoshino and Gaucher, 2021), and some phylogenetic signals

© 2022 The Authors. *Environmental Microbiology* published by Society for Applied Microbiology and John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Received 14 October, 2021; revised 27 December, 2021; accepted 1 January, 2022. \*For correspondence. E-mail melias@um.es; Tel. (+34) 868 887134; Fax (+34) 868 883963. padhu@iqfr.csic.es; Tel. (+34) 868 884951; Fax (+34) 868 883963

### 2 R. Pérez-Castaño et al.

(A)

supporting the theory can be traced to factors identified in *M. xanthus* light-induced carotenogenesis. Hence, exploring how this response and its unique factors are conserved can provide valuable molecular as well as evolutionary insights, and here we examined this across Myxococcales.

Two light sensing and signalling mechanisms control carotenogenesis in *M. xanthus* (Fig. 1A). One directly senses UV, blue or green light using a single protein, CarH<sub>Mx</sub> (Mx for *M. xanthus*), which is the defining member of a large photoreceptor family whose chromophore is coenzyme  $B_{12}$  or 5'-deoxyadenosylcobalamin (AdoCbl), a biological form of vitamin  $B_{12}$  (Fig. S1, Supporting Information) (Ortiz-Guerrero *et al.*, 2011; Jost *et al.*, 2015b; Padmanabhan *et al.*, 2017; Padmanabhan *et al.*, 2019). The other more complex mechanism

**B<sub>12</sub>-dependent (CarH)** 

involves several factors including CarA<sub>Mx</sub>, an AdoCblindependent CarH<sub>Mx</sub> paralogue and the extensive genetic rewiring that usually shapes the evolution of new regulatory networks (Perez and Groisman, 2009; Romero et al., 2012). It relies on <sup>1</sup>O<sub>2</sub> generated by blue light excitation of protoporphyrin IX. a photosensitizer and heme precursor (Fig. 1A) (Burchard and Dworkin, 1966; Galbis-Martínez et al., 2012). The protein known to act earliest in this pathway is CarF (Fontes et al., 2003), which we recently unmasked as the long-sought desaturase, widespread in metazoa but very rare in bacteria and required for the biosynthesis of plasmalogens, a special type of glycerophospholipids (Gallego-García et al., 2019). These lipids and  ${}^{1}O_{2}$ , through a still unknown mechanism, provoke the release of the extracytoplasmic function  $\sigma$  (ECF- $\sigma$ ) factor CarQ from its membrane-bound

> **Fig. 1.** The two pathways for regulating *M. xanthus* light-induced carotenogenesis.

Light A. Left panel: B<sub>12</sub>-dependent pathway. In the dark, CarH binds to its operator at  $P_{crt}$  (recognized by RNAP- $\sigma^A$ ) to Inner repress expression of the carotenoid membrane synthesis genes (crtE-orf9) of the PPIX carB operon. Light (UV/blue/green) VEPE disrupts CarH-operator binding to allow transcription from Pcrt. Right Lyso-PE panel: B<sub>12</sub>-independent pathway. Fatty CarA represses P<sub>crt</sub> in the dark and its aldehyde action is counteracted by CarS. expressed when protoporphyrin IX (PPIX) exposed to blue light generates photoexcited PPIX (PPIX\*) and <sup>1</sup>O<sub>2</sub>. Plasmalogens (VEPE), whose biosynthesis requires CarF, are essential for inactivation of CarR by a mechanism that may involve VEPE cleavage by <sup>1</sup>O<sub>2</sub> to its lyso-PE form and a fatty aldehyde (Gallego-García et al., 2019) to free CarQ, which associates with RNAP to activate P<sub>1</sub> and. together with the CarD-CarG complex, PORS. B. Molecular mechanism of action of CarH and CarA. Top: Binding of Ado-Cbl to apoCarH (monomer or molten





© 2022 The Authors. Environmental Microbiology published by Society for Applied Microbiology and John Wiley & Sons Ltd., Environmental Microbiology

**B<sub>12</sub>-independent (CarA)** 

anti- $\sigma$  CarR (Galbis-Martínez *et al.*, 2012; Gallego-García *et al.*, 2019). CarQ then associates with RNA polymerase (RNAP) to turn on transcription of the *carQRS* operon (encoding CarQ, CarR and CarS, an antirepressor of CarA<sub>Mx</sub>) and of the carotenogenic gene *crtlb* (Martínez-Argudo *et al.*, 1998; Browning *et al.*, 2003). This requires the CarD–CarG global regulatory complex, with some features reminiscent of multiprotein eukaryotic transcription complexes called enhanceosomes (Peñalver-Mellado *et al.*, 2016; García-Heras *et al.*, 2009; Elías-Arnanz *et al.*, 2010).

Both light signalling pathways converge at  $P_{crt}$  (or  $P_{B}$ ) (Fig. 1A), a promoter recognized by the primary  $\sigma^{A}$ -RNAP holoenzyme that drives expression of the carB operon. which encodes all the carotenoid synthesis enzymes except Crtlb, as well as CarH<sub>Mx</sub> and CarA<sub>Mx</sub> (López-Rubio et al., 2004; Pérez-Marín et al., 2008; Ortiz-Guerrero et al., 2011). CarH<sub>Mx</sub> requires AdoCbl for oligomerization (its precise oligomeric state is unknown) and DNA binding to repress transcription from P<sub>crt</sub>, and to be directly inactivated by light through the disassembly of the oligomer to monomers (Fig. 1B) (Pérez-Marín et al., 2008; Ortiz-Guerrero et al., 2011). In contrast to CarH<sub>Mx</sub>, homologues from species in other bacterial phyla, Thermus thermophilus (CarH<sub>Tt</sub>) and Bacillus megaterium (CarH<sub>Bm</sub>), have been purified and characterized in vitro, and crystal structures of the dark (free or DNAbound) and light-exposed forms have been resolved for CarH<sub>Tt</sub> (Ortiz-Guerrero et al., 2011; Díez et al., 2013; Kutta et al., 2015; Jost et al., 2015a; Jost et al., 2015b; Fernández-Zapata et al., 2018; Miller et al., 2020). In the dark, both proteins form AdoCbl-bound tetramers that bind to operators comprising tandem 11-bp direct repeats, three for CarH<sub>Tt</sub> or four for CarH<sub>Bm</sub>. However, while apo and light-exposed holoCarH<sub>Tt</sub> are both monomers, like CarH<sub>Mx</sub>, apoCarH<sub>Bm</sub> is a molten globule tetramer and light-exposed holoCarH<sub>Bm</sub> is a dimer (Fig. 1B). By contrast, CarA<sub>Mx</sub> is a dimer in the dark or light, with or without B<sub>12</sub>, binds cooperatively to operator DNA as two dimers, and requires expression of CarS in the light for derepression (Fig. 1B) (López-Rubio et al., 2004; Navarro-Avilés et al., 2007; León et al., 2010; Ortiz-Guerrero et al., 2011). CarH<sub>Mx</sub> and CarA<sub>Mx</sub> thus differ in AdoCbl-dependence, oligomerization, and DNA binding modes to control the same promoter and target genes.

CarH homologues are widely distributed in several bacterial phyla, usually as a single copy (Padmanabhan *et al.*, 2019). Hence, it is intriguing that CarH<sub>Mx</sub> coexists with CarA<sub>Mx</sub>, whose function has required coevolution of a cohort of other unique factors. Such duplications and divergence, and their retention or loss are of great interest in understanding transcription factor evolution (Hittinger and Carroll, 2007; Chapal *et al.*, 2019; Kuzmin *et al.*, 2020). We therefore examined conservation of

### Photoregulated carotenogenesis in myxobacteria 3

carotenogenesis and of the two *M. xanthus* photosensory regulatory pathways in all three Myxococcales suborders: Cystobacterineae (includes *M. xanthus*). Sorangiineae and Nannocystineae. This established that: (i) genes for carotenoid biosynthesis enzymes and for CarH are present in all myxobacterial genomes examined except in Anaeromyxobacter dehalogenans and Vulgatibacter incomptus, both with reduced genomes; (ii) that carH in Cystobacterineae always occurs in a large syntenic cluster of carotenogenic genes that in some species also contains, interestingly, a gene for 3-hydroxy-3-methylglutaryl-CoA reductase or HMGR (a separate syntenic operon encodes the primary HMGR and other enzymes of the mevalonate or MVA pathway for the synthesis of isoprenoids, the precursors of carotenoids); (iii) even carotenogenic genes are rearranged thouah in Sorangiineae and Nannocystineae, carH frequently occurs in the genomic vicinity. We provide evidence supporting B<sub>12</sub>-dependent photoregulation of carotenogenesis in representative myxobacteria from all three suborders and its likely mediation by CarH, whose operator design and DNA binding are, we show, remarkably plastic. The parallel CarA pathway and all its associated players appear strictly confined to suborder Cystobacterineae, among which the Cystobacter/ Melittangium genera have CarH and all factors of the CarA pathway yet, surprisingly, not CarA. We purified and characterized Cystobacter CarH (the one myxobacterial homologue thus far amenable to analysis in vitro) and show that it differs from the tetrameric  $CarH_{Tt}$  and Car-H<sub>Bm</sub>, the only others that have been characterized in vitro and which belong to different phyla. Thus, Cystobacter CarH acts as a dimer with CarH-like B12-dependent photosensory mode of action as well as CarA-like DNA binding and its antirepression by CarS. Altogether, this work reveals that B12-based photoregulation of carotenogenesis pervades across Myxococcales and promolecular-evolutionary vides insights into this photoprotective response in these bacteria.

### Results

# CarH is present in all three myxobacterial suborders and CarA only in Cystobacterineae

Myxococcales currently comprise three suborders and 10 families, and we examined 27 complete, publicly accessible myxobacterial genomes for genes encoding CarA/CarH homologues and other elements of photoinduced carotenogenesis identified in *M. xanthus* (Table S1). These myxobacteria are soil-dwelling, except for the aquatic *Enhygromyxa salina*, *Haliangium ochraceum* and *Plesiocystis pacifica*, and obligate aerobes except for *A. dehalogenans*, a facultative anaerobe. Their GC-rich

### 4 R. Pérez-Castaño et al.

(66%-75%) genomes are among the largest in bacteria (~9-16 Mb) but are reduced, presumably by extensive aene loss. in V. incomptus (~4.4 Mb) and A. dehalogenans (~5 Mb) (Huntley et al., 2011; Yamamoto et al., 2014). BLAST searches of these genomes using CarA<sub>Mx</sub> or CarH<sub>Mx</sub> (35% identical) as guery yielded one hit in Sorangiineae and Nannocystineae, and two hits in Cystobacterineae except for a single hit in Cystobacter ferrugineus, Cystobacter fuscus and Melittangium boletus, and none in A. dehalogenans or V. incomptus. In Cystobacterineae with two hits, one was most similar to CarA<sub>Mx</sub> (70%–98% identity, versus 33%–38% to CarH<sub>Mx</sub>) and the other to CarH<sub>Mx</sub> (62%-97% identity, versus 32%-34% to CarA<sub>Mx</sub>), suggesting the first and second sets to be CarA<sub>Mx</sub> and CarH<sub>Mx</sub> orthologs respectively (Fig. 2A and Table S2). Interestingly, the single hits in *C. ferrugineus*, C. fuscus and M. boletus were most similar to CarH<sub>Mx</sub> (57%-60% identity) than to CarA<sub>Mx</sub> (33%-36%), while sequence identities to CarH<sub>Mx</sub> (32%-39%) or to CarA<sub>Mx</sub> (29%-34%) were comparable for hits in Sorangiineae and Nannocystineae (Fig. 2A and Table S2).

Whether the observed hits are bona fide CarA<sub>Mx</sub> or  $CarH_{Mx}$  orthologs was assessed further by sequence analysis and experimentally.  $CarA_{Mx}$  and  $CarH_{Mx}$  have similar architectures, with an N-terminal MerR-type winged-helix DNA binding domain (DBD) and a C-terminal cobalamin-binding domain (CBD) (Fig. 2B). Sequence comparisons revealed patterns in each of these domains closely parallel to those for whole  $CarA_{Mx}$  or  $CarH_{Mx}$ , with the highest sequence identities between CarA<sub>Mx</sub> domains and likely CarA homologues, and CarH<sub>Mx</sub> domains and CarH homologues within Cystobacterineae likelv (Fig. 2A). Our previous studies of CarH<sub>Tt</sub>, CarH<sub>Bm</sub>, Car- $A_{Mx}$  and  $CarH_{Mx}$  established key motifs in both the DBD and CBD (Navarro-Avilés et al., 2007; Jost et al., 2015b; Fernández-Zapata et al., 2018). In the CarH<sub>Tt</sub> structure, the CBD comprises: (i) an N-terminal four-helix bundle subdomain with a  $Wx_9EH$  motif (x = any residue), conserved in CarH<sub>Mx</sub> and CarH<sub>Bm</sub>, which contacts the upper axial Ado group of CarH-bound AdoCbl and is crucial for CarH photoreceptor activity; (ii) a C-terminal five-stranded α/β Rossmann module housing the classic D/ExHx<sub>2</sub>G/



Fig. 2. CarA and CarH homologues in Myxococcales.

A. Heat map of the sequence identities of myxobacterial CarH/A homologues relative to CarA<sub>Mx</sub>, CarH<sub>Mx</sub> and their DBDs or CBDs (see Table S1 for species abbreviations).

B. Left: Structure of the CarH<sub>Tt</sub> protomer (PDB code: 5C8D) with its DBD (DNA recognition helix and wing in dark blue) and its CBD, with bound AdoCbl (sticks in magenta with upper axial Ado group in cyan). Right: Weblogos of myxobacterial CarH/A homologues alongside the corresponding subdomains in CarH<sub>Tt</sub>. Dots indicate DNA contacts (blue), Wx9EH motif (red), lower axial B12binding motif (magenta), dimer contacts (green) and dimer-dimer interface (black), based on the CarH<sub>Tt</sub> and DBD-CarA<sub>Mx</sub> structures, with filled dots indicating residues tested by mutational analysis (Navarro-Avilés et al., 2007; Jost et al., 2015b).

C. Maximum likelihood phylogeny (blue dots >80% bootstrap values) for myxobacterial CarH and CarA homologues.



Px41SxT/Vx22-27GG B12-binding motif, whose His supplies the lower axial ligand of bound AdoCbl (Fig. S1B and Fig. S2) (Jost et al., 2015b; Fernández-Zapata et al., 2018). The CarA<sub>Mx</sub> CBD retains the B12-binding motif and EH of the Wx<sub>9</sub>EH motif but lacks the Trp and four flanking residues (Fig. S2). Accordingly, whether an intact Wx9EH motif co-occurs or not with the classic B12binding motif may help distinguish CarH and CarA homologues. Classification of the Cystobacterineae homologues applying this criterion matched that based on overall sequence homology to CarA<sub>Mx</sub> and CarH<sub>Mx</sub>; moreover, all the single Sorangiineae and Nannocystineae homologues could be assigned as CarH. Consistent with these assignments, CarA homologues form a monophyletic clade within Cystobacterineae, separate from the coexisting CarH or the single Sorangiineae or Nannocystineae homologues (Fig. 2C). Sequence alignment of myxobacterial homologues thus classified as CarA or CarH vielded logos for the signature motifs in each domain shown in Fig. 2B. Besides the presence of a classic B<sub>12</sub>-binding motif in all myxobacterial homologues, and an intact Wx<sub>9</sub>EH motif only in CarH homologues, the logos revealed that the RxWERRY motif in the DNA recognition  $\alpha$ -helix and several other residues implicated in DNA binding are conserved in the predicted DBDs of both CarH and CarA homologues, suggesting a common mode of DNA recognition. Also, many CarH homologues have: Trp of the Wx<sub>9</sub>EH motif followed by a His, which in lightexposed CarH<sub>Tt</sub> is the upper ligand in a bis-His-cobalt linkage; a Glu-Arg or Asp-Arg pair equivalent to the crucial Arg-Asp salt-bridge pair of the CarH<sub>Tt</sub> dimer interface; and one of the two Gly at the dimer-dimer interface important in CarH<sub>Tt</sub> tetramer assembly (the other Gly is often Pro) conserved in CarH as well as CarA homologues (Fig. 2B).

To experimentally test our assignments, we first analysed several CarA and CarH homologues for selfinteractions using the bacterial adenylate cyclase twohybrid (BACTH) assay in E. coli (which cannot synthesize B<sub>12</sub> de novo but can take up and convert exogenous B<sub>12</sub> to AdoCbl for use in vivo). This assay has demonstrated that CarH<sub>Mx</sub> self-interacts via its CBD only in the dark and when B<sub>12</sub> is present, whereas CarA<sub>Mx</sub> self-interacts in the dark or light, irrespective of the presence or not of B<sub>12</sub> (Ortiz-Guerrero et al., 2011). In the BACTH assay, the homologues we assigned as CarA and CarH behaved like CarA<sub>Mx</sub> and CarH<sub>Mx</sub>, respectively, except for the sole C. ferrugineus, C. fuscus and M. boletus homologues, whose behaviour mirrored  $CarA_{Mx}$  rather than  $CarH_{Mx}$ (Fig. 3A). Negative controls with only one fusion protein ruled out autoactivation in the behaviour of the C. fuscus, C. ferrugineus and M. boletus homologues (Fig. S3).

We next checked our assignments by testing if  $CarH_{Mx}$  could be functionally replaced by representative CarH homologues: CarH\_{Sa}, CarH\_{Cfu} and CarH<sub>Cfe</sub> from

S. aurantiaca, C. fuscus and C. ferrugineus, respectively. outside the Myxococcus-Corallococcus branch in Cystobacterineae; CarH<sub>Sc</sub> from S. cellulosum in Sorangiineae; and CarH<sub>Ho</sub> from *H. ochraceum* in Nannocystineae. Each homologue was expressed from a vanillate-inducible promoter in an M. xanthus strain (MR2649) with a P<sub>crt Mx</sub>::/acZ transcriptional fusion and which lacks endogenous CarH, CarA, CarS and PduO, the ATP:corrinoid adenosyltransferase essential for Ado-Cbl generation, to enable controlled exogenous AdoCbl supply. All myxobacterial homologues tested, including CarH<sub>Cfu</sub> and CarH<sub>Cfe</sub>, downregulated reporter P<sub>crt Mx</sub>:: lacZ expression in the dark when AdoCbl was present but not when absent or when the cells were exposed to light, and hence are functionally equivalent to  $CarH_{Mx}$ (Fig. 3B).

We could purify CarH<sub>Cfu</sub> and CarH<sub>Cfe</sub> in the apo form as soluble proteins, unlike other myxobacterial CarH tested, and analyse them in vitro. Consistent with the BACTH assay, in size exclusion chromatography apo- $CarH_{Cfu}$  and apoCarH\_{Cfe} eluted predominantly as dimers  $(M_r \sim 59 \text{ kDa}, \text{ compared to } \sim 35 \text{ kDa} \text{ determined from the})$ sequence or by mass spectrometry) (Fig. 3C). In the presence of AdoCbl, both continued to elute as dimers in the dark ( $M_r$  ~62 and ~68 kDa respectively), but as monomers when exposed to light ( $M_r$  ~31 and ~33 kDa respectively) (Fig. 3C), in contrast to their self-interaction in BACTH analysis. In the latter assay, photolysis of intracellular AdoCbl and consequent poor binding to the two homologues probably causes them to persist as apoproteins and self-interact. The absorbance of the holoproteins at 522 nm (besides at 280 nm) indicated that both the dark and light forms bind to cobalamin (Fig. 3C) at ~1:1 stoichiometry. Moreover, the dark to light photoconversion of their AdoCbl-bound form upon stepwise illumination with green light (Fig. 3D) was similar to Car-H<sub>Tt</sub> or CarH<sub>Bm</sub> (Kutta et al., 2015; Fernández-Zapata et al., 2018; Miller et al., 2020). CarH<sub>Cfu</sub> and CarH<sub>Cfe</sub> are therefore AdoCbI-based CarH photoreceptors but transition from an apodimer to a holodimer to a photolyzed monomer, in a new twist to the oligomeric plasticity of these photoreceptors. Altogether, our analyses indicate that CarH occurs in all three myxobacterial suborders, but CarA only in Cystobacterineae and always with CarH.

# Genes for CarA, CarH and carotenoid synthesis usually occur in genomic proximity and occasionally linked to that for HMGR

To determine whether the link between carotenogenesis and CarH/A is maintained in other myxobacteria, we analysed the genomic context of *carA* and *carH*, and performed a search for carotenoid synthesis genes. We



found the latter in all myxobacteria examined except A. dehalogenans and V. incomptus, which also lack carA and carH (Fig. 4 and Table S2). Within Cystobacterineae, one phylogenetic branch that includes all *Myxococcus*. Corallococcus, Stigmatella, Hyalangium and Archangium species presents a genomic arrangement of carotenoid genes (with crtlb also at a different genomic location), carA and carH matching that in M. xanthus. Synteny is conserved in the branch containing C. ferrugineus, C. fuscus and M. boletus, except for the absence of carA (Fig. 4). In M. xanthus, the primary carotenoid end product in light-grown cells is myxobacton ester (a monocyclic carotenoid), although significant amounts of phytoene (the first C<sub>40</sub> compound in the biosynthetic pathway) are also found (Fig. S4) (Iniesta et al., 2007). This biosynthetic pathway is likely conserved in the other Cystobacterineae with the same set of carotenogenic genes as M. xanthus.

Interestingly, five species within Cystobacterineae (Stigmatella erecta, Hyalangium minutum, Archangium

Fig. 3. Experimental tests of myxobacterial CarH/CarA assignments. A. BACTH analysis of the effect of  $B_{12}$  and light on self-interaction of various myxobacterial CarH/CarA homologues. Cells expressing T25 and T18 fusions of CarH or CarA were spotted on X-Gal-LB plates with or without vitamin  $B_{12}$  and incubated for 48 h in the dark or exposed to light.

Complementation analysis B in M xanthus. Reporter Pcrt Mx::lacZ expression (specific β-galactosidase activity shown as the mean and standard error of three independent measurements) in the dark and in the light with AdoCbl present or absent in M. xanthus strain MR2649 bearing the indicated myxobacterial homoloque, expressed from a vanillateinducible promoter using vanillate at 100 µM (CarH<sub>Mx</sub>, CarH<sub>Sa</sub>, CarH<sub>Cfu</sub>, CarH<sub>Cfe</sub>), 2 µM (CarH<sub>Sc</sub>) or 500 µM (CarH<sub>Ho</sub>).

C. Size-exclusion chromatography elution profiles tracked using absorbance at 280 or 522 nm for CarH<sub>Cfu</sub> and CarH<sub>Cfe</sub> in the apo form in the dark, and with AdoCbl present in the dark (red) or after 5-min exposure to green light (green). Apparent  $M_r$  (kDa) for each eluted peak is on top.

D. UV-visible absorbance spectra for the photoconversion of AdoCbl-bound  $CarH_{Cfu}$  and  $CarH_{Cfe}$  upon stepwise illumination with green light for the times (in seconds) indicated.

gephyra, Archangium violaceum and M. boletus) have a gene encoding HMGR between crtB and crtD in the syntenic carotenoid cluster (Fig. 4 and Table S2). HMGR is an enzyme that acts early in the MVA pathway for producing isopentenyl pyrophosphate (IPP) and its dimethylallyl pyrophosphate (DMAPP) isomer, the C<sub>5</sub> building blocks for the biosynthesis of isoprenoids, including carotenoids (Fig. S4). The MVA pathway is one of the two known for isoprenoid biosynthesis and is used exclusively by archaea, fungi and animals; within bacteria, it is used only by a few species, since the majority employ the alternative methylerythritol 4-phosphate (MEP) path-(Moise et al., 2014; Rodriguez-Concepción wav et al., 2018; Hoshino and Gaucher, 2021). Our observation that a gene for HMGR lies within the carotenogenic cluster in some myxobacteria prompted us to examine isoprenoid biosynthesis pathways in the species studied Interestingly, while A. here. dehalogenans and V. incomptus have the complete gene set for the MEP pathway, all the remaining myxobacteria have MVA



Fig. 4. CarH, CarA and other factors in the  $B_{12}$ -independent pathway identified from myxobacterial genome analysis. The 16S rRNA-based phylogenetic tree for myxobacteria highlighting the three suborders and species, and the genes for *carA* (*A*), *carH* (*H*), and carotenoid synthesis. *B*: *crtB* (phytoene synthase); *C*: *crtC* (neurosporene hydroxylase); *cruA*: CruA-type lycopene cyclase; *cyc*: possible lycopene cyclase; *D*: *crtD* (hydroxyneurosporene dehydrogenase); *E*: *crtE* (geranyl geranyl pyrophosphate synthase); *IIIa/Ib*: *crtI* (phytoene desaturase); *L*: *crtL* (CrtL-type lycopene cyclase); *Y*: lycopene cyclase; fused to CrtU; *Yc*: *crtYc* (CrtYc subunit of heterodimeric lycopene cyclase); *Yd*: *crtYd* (CrtYd subunit of heterodimeric lycopene cyclase); *Ycd*: *crtYcd* (heterodimeric fusion type CrtYcd lycopene cyclase); *Z*: *crtZ* (β-carotene hydroxylase); *6*: *orf6* (possible glycosyltransferase); *9*: *orf9* (possible acyltransferase). Genes *carF*, *carQ*, *carR*, *carS*, *carD* and *carG* are indicated by the red circles if present and by unfilled circles if absent.

pathway genes and are thus among the minority of bacteria employing this typically eukaryotic route for isoprenoid biosynthesis (Hoshino and Gaucher, 2021). The gene for the primary HMGR, a class II type conserved in all myxobacteria (except the two mentioned earlier) tends to occur in a syntenic cluster with genes for enzymes catalysing the subsequent four steps of the MVA pathway (Figs. S4 and S5A, and Table S2). The primary HMGRs in *S. erecta*, *H. minutum*, *A. gephyra*, *A. violaceum* and *M. boletus* share 46%–51% sequence identity (all catalytic residues are conserved; Fig. S5B) with the second HMGRs encoded by the gene within the carotenogenic cluster.

Compared to Cystobacterineae, carotenoid synthesis genes in Sorangiineae and Nannocystineae are often rearranged and/or redistributed as various clusters, with *crtl* existing as a single copy (Fig. 4). Even so, syntenic blocks of carotenogenic genes with *carH* are observed in

the Sorangiineae examined (except for Sandaracinus amylolyticus and Labilithrix luteola), which are split into smaller blocks in Nannocystineae (Fig. 4). Some species appear to conserve only genes acting in the early biosynthetic steps, suggesting that their carotenoid end products may differ. Also, while M. xanthus and other Cystobacterineae (based on protein conservation relative to *M. xanthus*) use a heterodimeric CrtYc–CrtYd cyclase (Iniesta et al., 2008) to catalyse the formation of cyclic carotenoids from acyclic lycopene (Fig. S4), lycopene cyclases of the CruA family were found in Sorangiineae and of the CrtL family in Nannocystineae (Fig. 4). CruAtype cyclases typically occur in green sulfur bacteria and cyanobacteria, and the CrtL-type in cyanobacteria and plants (Maresca et al., 2007). Despite these differences, carH was generally found adjacent to carotenoid synthesis genes in Sorangiineae and Nannocystineae. Neither of these two suborders has a gene encoding HMGR

within a carotenoid synthesis gene cluster. Their only gene for HMGR occurs with MVA pathway genes, and a second copy is found only in *H. ochraceum* and *S. amylolyticus* (Fig. S5A). Interestingly, in the latter and in Nannocystineae, the MVA cluster includes genes for the synthesis of hopanoids (polycyclic triterpenes, like sterols, with six IPP units), which have been functionally and evolutionarily linked to carotenoids (Santana-Molina *et al.*, 2020). All the myxobacterial genomes analysed, excluding the downsized ones in *A. dehalogenans* and *V. incomptus*, thus have carotenoid synthesis genes, with proximal *carH* and/or *carA* in the majority, hinting at light and CarH/CarA-mediated regulation of carotenogenesis in these myxobacteria, as in *M. xanthus*.

# The B<sub>12</sub>-independent CarA pathway evolved solely in Cystobacterineae

We next surveyed myxobacterial genomes for the other essential factors of the M. xanthus B12-independent. CarA-dependent regulatory pathway (CarF, CarQ, CarR, CarS, CarD and CarG), which have singular features. Our search for myxobacterial homologues of CarF<sub>Mx</sub>, the desaturase required for the biosynthesis of plasmalogen lipids typically found in metazoa (Gallego-García et al., 2019), yielded single hits (each encoded by an isolated gene transcribed divergently relative to its flanking genes) in all Cystobacterineae except A. dehalogenans and V. incomptus, and in the other two suborders only in N. exedens (Fig. 4, Fig. S6 and Table S2). All these hits share sequence identities to  $CarF_{Mx}$  of  $\geq 70\%$  among Cystobacterineae (Fig. S6A) and 54% for that in N. exedens, and they conserve the nine histidines shown to be essential for function in  $CarF_{Mx}$  and its metazoan homologues (Gallego-García et al., 2019).

Of the three proteins encoded by the carQRS operon, CarQ belongs to the large ECF- $\sigma$  factor family, whose members are widespread, because use of such alternative  $\sigma$  factors constitutes a fundamental mechanism of signal transduction across eubacteria. Moreover, the number of ECF- $\sigma$  factors in myxobacteria tends to be high (~45 in M. xanthus, ~87 in S. cellulosum, ~118 in P. pacifica) (Abellón-Ruiz et al., 2014). Consequently, unequivocal assignment of genuine CarQ orthologs relied on the simultaneous presence of the far more distinct and CarS. CarR has the unusual CarR six transmembrane-helix topology of DUF1109 family anti-o factors largely restricted to Proteobacteria, some of which have been implicated in heavy metal or oxidative stress responses (Kohler et al., 2012; Masloboeva et al., 2012). However, CarR and its myxobacterial orthologs share low sequence similarity to other DUF1109 factors. Interestingly, CarR is a dual response anti- $\sigma$  that can be

inactivated by the joint action of light-induced  ${}^{1}O_{2}$  and plasmalogens (Galbis-Martínez *et al.*, 2012; Gallego-García *et al.*, 2019) as well as by copper (Moraleda-Muñoz *et al.*, 2005). On the other hand, the CarS antirepressor is a DNA mimic (León *et al.*, 2010) with no sequence homologues beyond Myxococcales. We identified CarR and CarS homologues only in Cystobacterineae and, except for *A. dehalogenans* and *V. incomptus*, in all species analysed in this suborder, with *carR* and *carS* always with *carQ* as a *carQRS* operon, as in *M. xanthus* (Fig. 4, Fig. S6 and Table S2).

Our earlier studies established the unique two-domain architecture of CarD, comprising an N-terminal RNAPinteracting domain, widely distributed among bacteria, which in CarD also interacts with CarG; a C-terminal K/A/ P-rich DNA-binding domain akin to eukarvotic high mobility group A (HMGA) proteins, with four or three repeats of their characteristic AT-hook motifs in CarD<sub>Mx</sub> and CarD<sub>Sa</sub> respectively, but which lacks the AT-hooks and resembles the histone H1 C-terminal region in CarD<sub>Ad</sub> (Padmanabhan et al., 2001; Cayuela et al., 2003; García-Heras et al., 2009; García-Moreno et al., 2010). On the other hand, an inactive version of the metzincin-type zinc-binding metalloprotease motif typically found in Archaea and vertebrates (Marino-Puertas et al., 2017) occurs in CarG, which regulates gene expression by interacting with another transcription factor (CarD) rather than directly with DNA (Peñalver-Mellado et al., 2006). Our search identified CarD and CarG orthologs (encoded in an operon, as in M. xanthus) only among Myxococcales and exclusively in Cystobacterineae, and in every species analysed in this suborder (Fig. 4, Fig. S6 and Table S2). Among CarD orthologs, only CarD<sub>vi</sub> has an H1-like C-terminal domain-like CarD<sub>Ad</sub>, the rest having four AT-hooks or three (in CarD<sub>Sa</sub>, CarD<sub>Se</sub> and CarD<sub>Hm</sub>).

In sum, given the absence of all six factors among Sorangiineae and Nannocystineae (except for CarF in *N. exedens*), the entire B<sub>12</sub>-independent pathway appears to have evolved uniquely in Cystobacterineae, in parallel with the advent of CarA. It is therefore intriguing that all six factors occur in *C. ferrugineus*, *C. fuscus* and *M. boletus*, even though none of the three species have CarA. Despite lacking CarH as well as CarA and four of its associated factors, *A. dehalogenans* and *V. incomptus* retain CarD and CarG, possibly because the two proteins act as a global regulatory complex in processes other than light-induced carotenogenesis (García-Heras *et al.*, 2009; Abellón-Ruiz *et al.*, 2014; Bernal-Bernal *et al.*, 2018).

# Pcrt operator design in Cystobacterineae

Most Cystobacterineae conserve the *M. xanthus* structural and regulatory genes for light-induced carotenogenesis, as well as their genomic arrangement (Fig. 4). This suggests a common strategy to control carotenoid synthesis, with both CarA and CarH-dependent pathways in the *Myxococcus*, *Corallococcus*, *Stigmatella*, *Hyalangium* and *Archangium* species, but only the latter in *C. ferrugineus*, *C. fuscus* and *M. boletus*. A common mode of regulation would likely be reflected by DNA sequence conservation around  $P_{crt}$ , which we analysed next.

DNA binding assays with purified CarA<sub>Mx</sub> revealed its binding as two dimers in a stepwise cooperative manner to an ~55 bp operator at P<sub>crt Mx</sub>, with the proposed CarA<sub>Mx</sub> binding site containing two inverted repeats (Fig. 5A): a perfect one upstream of the -35 promoter region and a second imperfect one overlapping with it (López-Rubio et al., 2004). However, among Cystobacterineae with CarA, only Myxococcus species and C. coralloides conserve a perfect or near-perfect match to the M. xanthus inverted repeats, which steadily degenerate in the rest.  $CarH_{Tt}$  and  $CarH_{Bm}$ , on the other hand, have been shown to bind as AdoCbl-bound tetramers to tandem 11-bp direct repeats (DRs) with the sequence 5'-nnnnTnnACAn-3' (*n* is any base). While  $CarH_{Tt}$  uses three of its DBDs to bind three tandem DRs, one of which includes the -35 promoter element, CarH<sub>Bm</sub> binds to operators comprising four such tandem DRs, or three DRs and one pseudorepeat (hereafter dr, where the TnnACA motif is not fully conserved), which overlap with the -35 or -10promoter elements (Jost et al., 2015b; Fernández-Zapata et al., 2018). Soluble native CarH<sub>Mx</sub> has resisted purification for analysis in vitro, but a functionally equivalent chimeric protein (CTt2) with CarH<sub>Mx</sub> CBD replaced by that of CarH<sub>Tt</sub> recognized the same ~55-bp operator at  $P_{crt Mx}$  as CarA<sub>Mx</sub> (Ortiz-Guerrero et al., 2011; Padmanabhan et al., 2017). Inspection of the ~55-bp DNA segment at P<sub>crt Mx</sub> revealed three 11-bp DRs (DR1, DR3 and DR4) conforming to those identified for  $CarH_{Tt}$  and  $CarH_{Bm}$ , and two drs (dr2 and dr5), the -35 TTGACA promoter element being within DR4 (Fig. 5A and Fig. S7). Sequence comparisons revealed a similar DR1-dr2-DR3-DR4-dr5 arrangement in all other Myxococcus species. C. coralloides, S. aurantiaca and S. erecta, which is dr1-DR2-DR3-DR4-dr5 in H. minutum, C. fuscus, C. ferrugineus and M. boletus, and dr1-dr2-DR3-DR4-dr5 in A. gephyra and A. violaceum (Fig. 5A and Fig. S7). DNA sequence logos from aligning the repeats at each promoter highlight the TnnACA motif, but also reveal variations between closely related species, particularly in the less conserved bases that can contribute additional sequence information to fine-tune DNA binding affinity and specificity (Fig. 5A and Fig. S7) (Jost et al., 2015b; Fernández-Zapata et al., 2018). Yet, despite these variations, all Cystobacterineae CarH homologues tested could replace CarH<sub>Mx</sub> to regulate  $P_{crt Mx}$  in *M. xanthus* (Fig. 3B), demonstrating their functional equivalence as well as DNA-binding plasticity.

# *B*<sub>12</sub>-dependent photoregulation of carotenogenesis, CarH DNA binding and its abrogation by light or CarS in Cystobacter

Among Cystobacterineae, *C. fuscus*, *C. ferrugineus* and *M. boletus* are particularly interesting because, besides CarH, they have all the factors in the CarA pathway except CarA (Fig. 4). Moreover, the *Cystobacter* CarH AdoCbl-bound form is a dimer (Fig. 3C) whereas CarH<sub>Tt</sub> or CarH<sub>Bm</sub> are tetramers, which can have consequences on operator size and/or mode of DNA recognition. We therefore examined how B<sub>12</sub> affects carotenogenesis in *C. fuscus* and *C. ferrugineus* and how purified *Cystobacter* CarH recognizes operator DNA *in vitro*.

First, we checked if cell culture colour depends on vitamin B<sub>12</sub> and/or light. Wild-type *M. xanthus* cell cultures are yellow (due to non-carotenoid pigments called DKxanthenes) (Meiser et al., 2006) in the dark with or without  $B_{12}$  in the growth medium because  $P_{crt Mx}$  is repressed by  $CarA_{Mx}$  under both conditions, and also by CarH<sub>Mx</sub> when  $B_{12}$  is available; but a carA-deleted  $(\Delta carA)$  strain is orange in the dark without B<sub>12</sub>, and yellow when  $B_{12}$  is present since only CarH<sub>Mx</sub> can repress P<sub>crt Mx</sub> (Fig. 5B). Both strains turn red in the light, irrespective of the presence or not of B12, due to the expression of the carB operon as well as of gene crtlb (Pérez-Marín et al., 2008; Ortiz-Guerrero et al., 2011). Like the  $\triangle carA$  M. xanthus strain, C. fuscus and C. ferrugineus were orange without B<sub>12</sub>, yellow or whitish with B<sub>12</sub> in the dark, and more intensely pigmented under light (Fig. 5B). Consistent with these colour changes, expression of the first gene (crtE) in the carotenogenic cluster [estimated by Quantitative Reverse Transcription Polymerase Chain Reaction (gRT-PCR)] was lower in the dark with B<sub>12</sub> present, and higher under light (Fig. 5C). Altogether, these data indicate that B<sub>12</sub> controls lightinduced carotenogenesis in С. fuscus and C. ferrugineus.

We next tested *in vitro* if and how purified CarH<sub>Cfu</sub> and CarH<sub>Cfe</sub> bind to 170-bp DNA probes containing the 55-bp segment mentioned earlier. In gel-shift assays (EMSA), apoCarH<sub>Cfu</sub> or apoCarH<sub>Cfe</sub> yielded only a diffuse smear even at high concentrations, suggesting weak non-specific binding (Fig. S8A). However, in the dark with AdoCbl present, defined retarded bands were observed, which were suggestive of stepwise DNA binding, from a smaller to a larger stable DNA-protein complex as the protein concentration increased (Fig. 5D and Fig. S8A). Binding was cooperative with an affinity (apparent



Fig. 5. B<sub>12</sub>-dependent light-induced carotenogenesis and CarH DNA binding in *Cystobacter*.

A. Alignment of the CarA<sub>Mx</sub>/CarH<sub>Mx</sub> operator DNA sequence (sense strand) at  $P_{ort Mx}$  and the corresponding segments for Cystobacterineae species with CarH. Red dots align with T and ACA of the consensus DR motif, and the span of the CarA<sub>Mx</sub>/CTt2 DNase I footprint at  $P_{ort Mx}$  is indicated; below the logo is a perfect inverted repeat (red arrows) and a less perfect one (purple arrows) found in some *Myxococcus* species. The -35 promoter element within DR4 and the remaining four 11-bp repeats are indicated at the bottom.

B. Left: Colour of cell culture spots on plates with or without vitamin  $B_{12}$  incubated for 48 h in the dark or in the light for *M. xanthus* (wild-type and  $\Delta carA$  strains), *C. fuscus* and *C. ferrugineus* with the corresponding carotenogenic gene cluster and sequence logo (TnnACA segment or equivalent shaded) of the five direct repeats at  $P_{ort}$ .

C. Expression of *crtE* from *C. fuscus* and *C. ferrugineus* measured by qRT-PCR with *rpoD* as the reference gene reported relative to the level in the dark with vitamin  $B_{12}$  present (mean and standard error from three biological replicates).

D. Representative EMSA for the binding of probes P<sub>crt Cfu</sub> and P<sub>crt Cfe</sub> to increasing concentrations of AdoCbl-bound CarH<sub>Cfu</sub> or CarH<sub>Cfe</sub> (0.1, 0.25, 0.5, 1, 5, 10, 100 nM), respectively, in the dark or after 5-min exposure to green light.

 $K_{\rm D} \approx 35 \pm 1$  nM; Fig. S8B) comparable to the estimates reported for CarH<sub>Tt</sub> and CarH<sub>Bm</sub> to their respective operators (Jost *et al.*, 2015b; Fernández-Zapata *et al.*, 2018). Exposure to light eliminated the retarded bands, indicating poor DNA binding by the photolyzed CarH<sub>Cfu</sub> or Car-H<sub>Cfe</sub> monomer (Fig. 5D).

We next mapped the site recognized by  $CarH_{Cfu}$  at  $P_{crt}$ <sub>Cfu</sub> using footprinting analyses. CarH<sub>Cfu</sub> binding yielded a large DNase I footprint, which coincides well with the segment based on alignment with the  $CarA_{Mx}/CarH_{Mx}$  operator (Fig. 5A) and which includes part of dr1 and DR2-dr5. Within the DNase I footprint, four tracts (2–3 nt each) spaced 10–11 nt apart (~1 pitch of double-stranded B-DNA) were protected from hydroxyl radical attack on each strand (Fig. 6A and C). Moreover, CarH<sub>Cfu</sub> binding arrested Exo III at ~4 nt downstream of dr5 (strongly) and

DR4 (weakly) in the sense strand, and at ~2-4 nt upstream of DR2 (strongly) and dr1 (weakly) in the antisense strand (Fig. 6B and C). Taken together, the sizes of the DNase I/Exo III footprints and the four evenly spaced hydroxyl radical footprints are consistent with cooperative binding to one face of a DNA site composed of the three tandem DRs plus one dr (dr1-DR4 or DR2-dr5, both including the -35 promoter region) by two CarH<sub>Cfu</sub> dimers, each using its two DBDs. Accordingly, CarH<sub>Cfu</sub> could bind to 52-bp probes spanning dr1-DR4 or DR2-dr5, yielding the two retarded bands observed with the 170-bp P<sub>crt Cfu</sub> DNA fragment or with a 63-bp probe spanning all five repeats (Figs. 5D and 6D). By contrast, the upper band normally observed at higher CarH<sub>Cfu</sub> concentrations was weak for probes with three DRs (2-4) or two DRs and one dr (1-3, 3-5), and absent for probes with two DRs (2\*-3-4, 2-3-4\*, generated by mutating one of the two outlying DRs of the three DR-probe), with which a stable retarded band nonetheless continued to be observed (Fig. 6D). Moreover, the CarH<sub>Cfu</sub> dimer requires the two repeats to be in tandem, since no binding was detected for probe 2-3\*-4, with the central DR of the three-DR probe mutated (Fig. 6D). Stable formation of the larger complex therefore requires at least four repeats and two CarH<sub>Cfu</sub> dimers, since two DBDs from each dimer would bind to two tandem repeats.

Since CarH<sub>Cfu</sub> can regulate P<sub>crt Mx</sub> in *M. xanthus*, we probed how it binds to P<sub>crt Mx</sub> in vitro. We also tested whether and how CarA<sub>Mx</sub>, a dimer like CarH<sub>Cfu</sub> but B<sub>12</sub>independent, binds to P<sub>crt Cfu</sub>, which lacks the palindromes (Fig. 5A) that were suggested as recognized by CarA<sub>Mx</sub> (López-Rubio et al., 2004). In EMSA, DNase I and Exo III analyses (Fig. S9) both proteins bound in a similar cooperative, stepwise manner to target the same DNA stretch at both P<sub>crt Cfu</sub> or P<sub>crt Mx</sub>, despite variations in the DR sequences and their arrangement and the absence of the inverted repeats in Pcrt Cfu, suggesting that CarA<sub>Mx</sub> also recognizes DRs rather than inverted repeats. Differences in some features, like DNase I hypersensitive sites (usually indicative of DNA distortions from protein binding), possibly reflect intrinsic differences between the two proteins, such as AdoCbl dependence only for CarH<sub>Cfu</sub>. Thus, even though CarH<sub>Cfu</sub> requires AdoCbl for DNA binding and CarA<sub>Mx</sub> does not, both are dimers with similar modes of DNA recognition. In addition, purified CarS<sub>Cfu</sub> abolished DNA binding by AdoCblbound CarH<sub>Cfu</sub> (Fig. 6E), just as CarS<sub>Mx</sub> abolishes DNA binding by CarA<sub>Mx</sub>. CarH<sub>Cfu</sub> therefore combines specific AdoCbl-dependent DNA binding in the dark and direct disruption by light characteristic of its tetrameric homologues CarH<sub>Tt</sub> and CarH<sub>Bm</sub>, with the cooperative stepwise binding as a dimer to operator DNA and its inhibition by the cognate CarS<sub>Cfu</sub> that characterizes the B<sub>12</sub>-independent CarA<sub>Mx</sub> dimer (Fig. 6F).

# *B*<sub>12</sub>-dependent photoregulation of carotenogenesis, CarH operator design and DNA binding in Sorangiineae and Nannocystineae

Sorangiineae and Nannocystineae differ from Cystobacterineae, where the majority of carotenogenic genes and carH occur as a large syntenic operon with a single promoter region that generally contains three DRs, with at least two in tandem, and two drs. Among Sorangiineae, carotenogenic genes are located at several genomic loci and unlinked to carH in S. amvlolvticus and L. luteola, and our search in their putative promoter regions for tandem 11-bp direct repeats as possible CarH binding sites revealed none in S. amylolyticus and only one upstream of carH in L. luteola, with two tandem DRs and one dr (Fig. S10A). In the remaining Sorangiineae, we could identify one DR flanked by two or three drs in the intergenic segments of the divergently expressed crtlcarH and crtE-crtD gene pairs, the latter with three DRs in tandem in S. cellulosum (Pcrt Sc2; Fig. S10A). Similar analyses in Nannocystineae (where carH is isolated in N. exedens but amidst carotenogenic genes in the others) identified at least one putative CarH binding site with two tandem DRs and two adjacent drs in all except E. salina, which has only one DR at both possible sites flanked by 2-3 drs (Fig. S10B). To experimentally test CarH binding to these diverse possible sites, we chose S. cellulosum and H. ochraceum, which we could grow in liquid cultures and/or plates with or without B12, as representative of Sorangiineae and Nannocystineae respectively.

We found that S. cellulosum undergoes B12- and lightdependent cell culture colour changes, being orange in the dark without B12 or when exposed to light but yellow in the dark with B<sub>12</sub> present (Fig. 7A), analogous to the  $\Delta carA$  M. xanthus and the two Cystobacter strains (Fig. 5B). Moreover, expression of the divergent crtlcarH and crtE-crtD genes was repressed in the dark in the presence of B<sub>12</sub> and derepressed upon exposure to light (Fig. 7B), confirming that B<sub>12</sub> and light modulate expression of carotenoid genes in S. cellulosum. We then examined AdoCbl-dependent binding at the crtlcarH (P<sub>crt Sc1</sub>) and crtE-crtD (P<sub>crt Sc2</sub>) intergenic regions in vitro. Because native soluble CarH<sub>Sc</sub> could not be purified, we resorted to the strategy used previously to test CarH<sub>Mx</sub> binding to P<sub>crt Mx</sub> in vitro (Ortiz-Guerrero et al., 2011). For this, the CarH<sub>Sc</sub> DBD (and its specific DNA-binding determinants) and linker were fused to the CarH<sub>Tt</sub> CBD to generate chimera CSc, which was purifiable. In EMSA, binding of CSc to Pcrt Sc1 and Pcrt Sc2 was clearly observed only in the dark and with Ado-Cbl present (Fig. 7C). Consistent with three tandem DRs identified in P<sub>crt Sc2</sub> and only one DR and two drs in P<sub>crt</sub> Sc1, CSc exhibited higher affinity for Pcrt Sc2, since an



Fig. 6. Mapping of the binding site at P<sub>crt Cfu</sub> and mode of action of CarH<sub>Cfu</sub>.

A. Representative DNase I and hydroxyl radical footprints upon binding of 100 nM CarH<sub>Cfu</sub> to the sense and antisense strands of P<sub>crt Cfu</sub>. Lines to the left: DNase I footprints; lines to the right: hydroxyl radical footprints; arrowheads: DNase I hypersensitive sites.
B. Representative Exo III footprints for CarH<sub>Cfu</sub> binding to P<sub>crt Cfu</sub>. Positions of Exo III arrest: horizontal lines on the side (thicker for stronger arrest).

C. Summary of footprint data generated by CarH<sub>Cfu</sub> binding at  $P_{ort Cfu}$ . DNase I footprints: horizontal lines; DNase I hypersensitive sites: arrow-heads; sites protected from hydroxyl radical attack: dots; positions of Exo III arrest: vertical lines (thicker for stronger arrest); -35 element: lettered red; -10 element (part): blue. DR: tandem 11-bp direct repeat with the sequence 5<sup>'</sup>-*nnnnTnnACAn-3*<sup>'</sup> (*n* is any base); dr: 11-bp direct repeat where the T*nnACA* motif is not fully conserved.

Fig. 6. Legend on next page.

eight-fold lower CSc concentration was sufficient for maximal binding to  $P_{crt Sc2}$  compared to  $P_{crt Sc1}$ . Moreover, CSc yielded clearer DNase I, hydroxyl and Exo III footprints at  $P_{crt Sc2}$  than at  $P_{crt Sc1}$  (Fig. 7D–F). Nonetheless, the three tandem repeats in either probe mapped to the ~34–38 bp DNase I footprints and were delimited by positions where Exo III advance was blocked by CSc. Thus, CSc can bind to a site with three tandem DRs ( $P_{crt Sc2}$ ), like CarH<sub>Tt</sub>, but also to one with just one DR and two drs ( $P_{crt Sc1}$ ). Putative –35 and/or –10 promoter elements that could be identified within the footprinted regions of both probes hint at how CarH<sub>Sc</sub> binding at a single site in either probe can simultaneously regulate expression from divergent promoters (Fig. 7F).

A similar analysis was carried out for H. ochraceum, which was also less pigmented in the dark with B<sub>12</sub> present than without, or in the light (Fig. 8A). Like CarH<sub>Sc</sub>, CarH<sub>Ho</sub> was not purifiable as a soluble native protein. We therefore again generated and purified a chimera, CHo, fusing the CarH<sub>Ho</sub> DBD and linker to the CarH<sub>Tt</sub> CBD and tested its binding to a 170-bp probe ( $P_{crt}$  Ho) corresponding to the upstream non-coding region of the carotenogenic gene cluster containing carH (Fig. 8B). CHo could bind to P<sub>crt Ho</sub> only in the dark with AdoCbl present (Fig. 8B) and footprinting analysis mapped the binding site, which overlaps with putative -35 and -10 promoter regions, to four tandem 11-bp repeats comprising two DRs and two drs (Fig. 8C-E). Thus, light and B<sub>12</sub> also appear to regulate via CarH the expression of carotenogenic gene clusters in Sorangiineae and Nannocystineae, despite their distinct arrangements from that in Cystobacterineae. Moreover, the data reveal, for the first time, that even operators with only one or two DRs (with flanking drs) can be recognized by CarH, highlighting unusual plasticity in DNA recognition and operator design.

### Discussion

Studies on *M. xanthus* light-induced carotenogenesis have uncovered new paradigms in bacterial light sensing, signalling and response, and led to the discoveries of widely distributed protein families with novel functions.

Central players in the genetic circuit underlying this M. xanthus response to light are two transcriptional factors with similar domain architectures and DNA target sites, CarH<sub>Mx</sub> and CarA<sub>Mx</sub>, which control the expression of the same genes by distinct mechanisms. While an AdoCbl chromophore directly senses light to allosterically modulate CarH repressor activity, regulation by CarA requires the antirepressor CarS, whose light-induced expression relies on a multifactorial pathway that additionally includes CarF, CarQ, CarR, CarD and CarG. Our present study on the conservation of these elements for photoinduced carotenogenesis across diverse myxobacteria provides molecular and evolutionary insights into these factors and establishes that B<sub>12</sub>-dependent photoregulation of carotenogenesis pervades in this major class of social and predatory soil and marine bacteria.

Our sequence analysis with experimental validation assigned CarH homologues in all the myxobacteria examined, except in two species with reduced genomes, A. dehalogenans and V. incomptus. Both of these also lack genes for carotenoid synthesis and probably rely on ROS detoxifying enzymes and/or light-avoidance mechanisms to combat photooxidative stress. The majority have carH in the genomic neighbourhood of genes for carotenogenesis, which occur as a large syntenic block in Cystobacterineae and as smaller syntenic blocks in Sorangiineae or Nannocystineae. Genomic proximity of carH to carotenogenic genes points to functional, presumably evolutionary, links between them that may be related to CarH's ability to sense light and elicit the photoprotective response over a broader spectral range (UV to green) than most other known photoreceptors. In contrast to CarH, our analysis revealed CarA only in suborder Cystobacterineae, always with CarH, and invariably encoded by the gene immediately upstream of carH in the carotenogenesis gene cluster characteristic of this suborder. That carA and carH exist in tandem and their close phylogenetic relationship suggests that their cooccurrence is likely due to gene duplication. Gene duplication and divergence is a major mechanism for the evolution of new genes or paralogues with overlapping or non-overlapping functions and has been proposed as a key contributor to genomic expansion and sensory com-

D. Top: Sequences (sense strand) of oligonucleotide probes used, with mutations in blue lowercase. Bottom: Binding to the indicated probes (number of repeats shown below) by CarH<sub>Ctu</sub> (50, 100 and 200 nM; 100 and 200 nM for mutant DR2-4 probes).

E. Representative EMSA with CarH<sub>Cfu</sub> (100 nM) and increasing concentrations of its cognate CarS<sub>Cfu</sub> (100, 200, 400, 800 nM, added after prior incubation of CarH<sub>Cfu</sub> with P<sub>crt Cfu</sub>). DNA binding assays in A–E were performed in the dark with AdoCbl present at five-fold excess relative to protein.

F. Model summarizing CarH<sub>Cfu</sub> mode of action. ApoCarH<sub>Cfu</sub> dimers bind to AdoCbl to form active CarH<sub>Cfu</sub> dimers, which bind cooperatively to an operator at P<sub>crt Cfu</sub> with two overlapping sites, each comprising three tandem DRs (filled grey arrows) and one dr (unfilled arrows). Binding of two CarH<sub>Cfu</sub> dimers to dr1–DR4 (as shown) or to DR2–dr5 would block the –35 promoter region and repress transcription. Light (UV/blue/green) cleaves the AdoCbl Co–C bond to liberate the upper axial ligand (unfilled red circles) provoking disassembly of CarH<sub>Cfu</sub> dimers to monomers that retain photolyzed AdoCbl (unfilled red triangles) and do not bind DNA. Alternatively, P<sub>crt Cfu</sub> can be derepressed by CarS<sub>Cfu</sub>, which sequesters CarH<sub>Cfu</sub> to block DNA binding.



Fig. 7. B<sub>12</sub>-dependent light-induced carotenogenesis and CarH DNA binding in S. cellulosum.

A. Colour of S. cellulosum cell culture spots on plates with or without vitamin B<sub>12</sub> incubated for 48 h in the dark or in the light.

B. Expression of the indicated gene measured by qRT-PCR using *rpoD* as the reference gene, and reported relative to the level in the dark with vitamin B<sub>12</sub> present (mean and standard error from three biological replicates).

C. Top: *S. cellulosum* carotenoid synthesis gene cluster with sequence logos (T*nn*ACA segment or equivalent shaded) from the three putative 11-bp repeats at  $P_{crt Sc1}$  (dr1, dr2, DR3) and  $P_{crt Sc2}$  (DR1–DR3). Bottom: Representative EMSA for the binding (in the dark or after 5-min green light exposure in the presence or absence of AdoCbl) of CSc to  $P_{crt Sc1}$  (left gel: 60, 120, 240, 480, 960 nM; right gel: 1000 nM CSc) and to  $P_{crt Sc2}$  (left gel: 10, 20, 40, 80, 120 nM CSc; right gel: 120 nM CSc).

D. Representative DNase I and hydroxyl radical footprints on the sense and antisense strands (relative to the indicated gene) for binding of 1000 nM CSc to P<sub>ort Sc1</sub> and 240 nM CSc to P<sub>crt Sc2</sub> (in the dark and presence of AdoCbl).

E. Representative Exo III footprint on the sense and antisense strands of probes P<sub>crt Sc1</sub> and P<sub>crt Sc2</sub> upon CSc binding as in D.

F. Summary of footprint data at P<sub>crt Sc1</sub> and at P<sub>crt Sc2</sub> upon CSc binding. Proposed promoter elements (red, -35 region; blue, -10 region) are shown (note overlap of the two divergent promoters). Symbols in D–F are as in Fig. 6A–C respectively.

plexity in myxobacteria (Goldman *et al.*, 2006; Schneiker *et al.*, 2007). Tandemly duplicated genes, usually arising from recombination-based mechanisms, are often restricted to closely related species, tend to share similar functions and regulatory elements, and typically

contribute to adaptation and response to environmental stimuli and species-specific traits (Simon *et al.*, 2015; Chapal *et al.*, 2019; Kuzmin *et al.*, 2020). All of these apply to the CarH–CarA pair. Given the ancient prebiotic origin proposed for  $B_{12}$  (Roth *et al.*, 1996; Monteverde



**Fig. 8.**  $B_{12}$ -dependent light-induced carotenogenesis and CarH DNA binding in *H. ochraceum*.

A. Colour of *H. ochraceum* cells streaked on plates with or without vitamin  $B_{12}$  incubated for 48 h in the dark or in the light.

B. Top: *H. ochraceum* carotenoid synthesis gene cluster and sequence logo (TnnACA segment or equivalent shaded) derived from the four repeats at  $P_{crt}$  <sub>Ho</sub>. Bottom: representative EMSA for the binding of CHo to a 221-bp  $P_{crt}$  <sub>Ho</sub> probe in the presence or absence of AdoCbl in the dark (left and middle gels: 2.5, 5, 10, 15 nM CHo) or after 5-min green light exposure (right gel: 10 nM CHo).

C. Representative DNase I and hydroxyl radical footprinting on the sense and antisense strands of probe  $P_{crt Ho}$  with or without CHo in the dark and presence of AdoCbl.

D. Representative Exo III footprinting of CHo binding to probe  $P_{ort}$  Ho as in C.

E. Summary of footprint data on each strand at  $P_{ort\ Ho}$  upon CHo binding. Proposed promoter elements (red, -35 region; blue, -10 region) are shown. Symbols in C–E are as in Fig. 6A–C respectively.

et al., 2017), the phylogeny, and far greater prevalence of CarH over CarA, it is likely that an ancestral AdoCbldependent CarH duplicated and diverged to create a B12independent CarA. The distribution of carH and carA genes in the myxobacterial genomes analysed in this study suggests that such a gene duplication could have occurred in the ancestor of the Myxococcus-Archangium branch or, alternatively, in an even earlier Cystobacterineae predecessor, with subsequent gene loss resulting in just one copy in C. ferrugineus, C. fuscus and M. boletus, and none in A. dehalogenans and V. incomptus.

Transition from CarH to CarA entailed loss of the  $Wx_9EH$  motif, crucial for AdoCbl-specific binding,

oligomerization and DNA binding of CarH, together with changes to ensure that CarA forms stable dimers independently of B<sub>12</sub>. Most remarkably, photosensory regulation by CarA has required coevolution of various singular proteins, some eukaryotic-like, and a signature metazoan lipid. Myxobacteria are the only strictly aerobic bacteria known to have the oxygen-dependent desaturase (CarF) for plasmalogen biosynthesis whose equivalent, despite the enormous evolutionary distance, is specifically conserved in metazoa (Gallego-García et al., 2019). The existence in myxobacteria of factors considered typically eukaryotic point to tangible evolutionary links to eukaryotes and possible horizontal gene transfer events that provide supporting evidence for the Syntrophy

hypothesis, which posits an ancient myxobacterium as crucial in eukaryogenesis (López-García and Moreira, 2020; Hoshino and Gaucher, 2021). Thus, steroid biosynthesis has been proposed to have first evolved in an early myxobacterium in response to the rise of oxygen and later horizontally transferred to an early eukaryote, because myxobacteria form the select set of aerobic bacteria to have both the steroid and its isoprenoid precursor (MVA) biosynthesis pathways homologous to those in eukaryotes (Hoshino and Gaucher, 2021). It is tempting to speculate that this also extends to the oxygen-based biosynthesis of plasmalogens, given the existence of CarF in myxobacteria and in metazoans, and its absence in nearly all other bacteria. However, while eukaryotic-type steroid biosynthesis enzymes appear across all three myxobacterial suborders (Hoshino and Gaucher, 2021), we found CarF almost entirely in Cystobacterineae. Curiously, besides the gene for HMGR that is part of an operon with various other MVA pathway genes, the large syntenic block of carotenogenic genes in some Cystobacterineae includes a second copy of a gene for HMGR. Since HMGR is generally a rate-limiting enzyme in the MVA pathway, its photoinduced co-expression with carotenogenic enzymes possibly ensures ample supply of the earliest precursors (IPP/DMAPP) of carotenoids for light-induced production.

CarR orthologs also appear to have evolved solely in Cystobacterineae to act in concert with plasmalogens and  ${}^{1}O_{2}$  in a photoprotective stress response. They diverge significantly in sequence from other DUF1109 anti-o but may nonetheless retain a role in responses to metals, as observed for CarR<sub>Mx</sub> in the copper response. Moreover, CarR has had to co-evolve with CarQ to recognize it specifically from among the numerous ECF- $\sigma$ factors typically found in myxobacteria. CarS homologues are yet another class of proteins found exclusively in Cystobacterineae. CarS is a small, taxonomically restricted protein acting in a stress response and thus has features typically ascribed to a de novo gene product (Schlotterer, 2015; Van Oss and Carvunis, 2019). Also crucial in the CarA pathway is the highly unique CarD-CarG complex, formed by two proteins encoded by adjacent genes that always co-exist in every (and only in) Cystobacterineae analysed here. In CarD, the fusion of an N-terminal RNAP-interacting module of exclusively bacterial origin with a C-terminal DNA-binding domain that resembles eukaryotic HMGA or histone H1 is intriguing at the evolutionary and functional levels. It has been proposed that the origins of the characteristic K/A/P-rich H1 C-terminal region can be traced to eubacteria, from where it was acquired by horizontal gene transfer and evolved in eukaryotes (Kasinsky et al., 2001). If so, an ancestral CarD could have arisen from fusion of two bacterial modules, one of which was a C-terminal K/A/Prich H1-like region as in *A. dehalogenans/V. incomptus*, from which the remaining Cystobacterineae acquired the HMGA AT-hooks by convergent evolution. Two independent fusion events, one to an H1-like region and another to an HMGA-like region, would be less probable. Since CarD always functions as a single unit with CarG, the two proteins must have co-evolved. Horizontal gene transfer followed by functional divergence may have led to the emergence of CarG, as it lacks any protease activity yet has a zinc-binding metalloprotease motif of the type typically found only in archaeal and metazoan metzincins.

Strikingly, Cystobacter/Melittangium species in Cystobacterineae lack CarA despite having all other known factors of this pathway. The single Cystobacter CarH/A homologue, our data show, is an AdoCbl-based CarH photoreceptor that transitions from an inactive dimeric apo form to active AdoCbl-bound dimers, which become inactive monomers upon photolysis. The contrast with the oligomeric transitions of the only other wellstudied CarH homologues, CarH<sub>Tt</sub> (monomer-tetramermonomer) and CarH<sub>Bm</sub> (tetrameric molten globule-tetramer-dimer) highlights the oligomerization plasticity of these photoreceptors. Oligomerization confers properties crucial for protein function, like cooperative ligand binding, allosteric regulation and stability and changes in oligomerization through evolution usually enable new regulatory modes. But how a given multimer and its functions arise is poorly understood. Extant oxygen-binding haemoglobin tetramers, with their striking cooperativity and regulatory modes, may have evolved by successive gene duplication-divergence events through surprisingly few mutations from an ancestral dimeric precursor, which itself arose from an ancient monomer (Pillai et al., 2020). Single mutations at the interface between the two headto-tail dimers that constitute the CarH<sub>Tt</sub> tetramer suffice to shift the tetramer to a native dimer, which retains AdoCbl-dependent DNA binding, but with altered affinity and cooperativity (Jost et al., 2015b); CarH<sub>Tt</sub> tetramerization traces a kinetic pathway from monomer to dimer to tetramer (Camacho et al., 2021). Hence, few genetic changes may underlie the shift between the AdoCbl-dependent Cystobacter CarH dimer and its tetrameric homologues.

We found that despite the distinct oligomerization, the *Cystobacter* CarH dimer emulates its tetramer homologues in its AdoCbl-dependent binding to one face of a DNA region comprising 11-bp tandem direct repeats. It does so by cooperative binding as two dimers, effectively a tetramer, to two overlapping sites, each composed of three tandem DRs and one dr. Whereas each CarH<sub>Cfu</sub> dimer requires the two repeats to be in tandem, in the

case of the CarH<sub>Tt</sub> tetramer one head-to-tail dimer interacts with two non-contiguous DRs (DR1 and DR3) and the other contacts the central DR (DR2) using only one of its two DBDs (Jost et al., 2015b). Furthermore, CarH<sub>Cfu</sub> and CarA<sub>Mx</sub> recognize each other's operators in vitro and employ a similar DNA binding mode, regardless of the variations between their natural sites and the requirement for AdoCbl by CarH<sub>Cfu</sub> but not by CarA<sub>Mx</sub>; CarH<sub>Cfu</sub> can direct AdoCbl-based regulation of P<sub>crt Mx</sub> in vivo, emphasizing the evolutionary relationship between the two proteins and their operators. Indeed, in every CarHcontaining Cystobacterineae analysed we discerned counterparts to the M. xanthus and Cystobacter operators in size and sequence, generally with three DRs (and two drs) and an identical -35 promoter region embedded in one of them, such that binding of CarH would block access to RNAP. In Sorangiineae and Nannocystineae, CarH operators are less like those in Cystobacterineae. Most species appear to have two operators that differ among themselves, each expected to control two divergent promoters, given the overlap with putative promoter regions. Thus, two CarH operators were experimentally mapped in S. cellulosum, one comprising three tandem DRs and the other just one DR and two drs, while the single operator mapped in H. ochraceum has two DRs and two drs. Even though CarH operators across the three suborders are quite varied, CarH homologues from the three myxobacterial suborders could functionally replace CarH<sub>Mx</sub> for AdoCbl-dependent photoregulation of P<sub>crt Mx</sub> in M. xanthus. This highlights an unusual DNA-binding plasticity of these homologues, possibly providing cells a strategy to fine-tune affinities and operator occupancies for optimal regulation in vivo.

The B<sub>12</sub>-independent CarA pathway, absent in both Sorangiineae and Nannocystineae, affords an alternative mechanism to the B12-dependent CarH one in Cystobacterineae. Parallel mechanisms usually confer a fitness advantage by boosting response efficiency and control, and enable adaptive evolution through cooperation and functional divergence to bestow benefits not gained from just a single mechanism (Chapal et al., 2019; Kuzmin et al., 2020). Direct light-sensing and regulation by AdoCbl-CarH ensures a very rapid and efficient response. However, myxobacteria cannot perform the expensive de novo biosynthesis of the complex Ado-Cbl cofactor, which requires as many as 30 steps, because they do not have all the necessary enzymes (Shelton et al., 2019). They consequently must depend on their environmental niche to acquire B<sub>12</sub>, a frequently transacted resource in microbial consortia (Sokolovskaya et al., 2020), by preving on microorganisms that can supply B<sub>12</sub> or a suitable precursor, or by importing these directly from the milieu. A need to cope with limited B12 probably drove the emergence of the CarA pathway in myxobacteria, which allows them to still regulate the photoprotective response when B<sub>12</sub> is unavailable. An additional adaptative advantage conferred by this pathway is its possible implication in other independent or overlapping responses, as illustrated by its role in the response to copper. Even without CarA, presence of the other factors in the B12-independent pathway would expand the range of responses beyond that to light. Moreover, given that CarQ is an alternative ECF- $\sigma$  factor, its regulon may extend beyond genes for carotenoid synthesis or their specific regulators (our unpublished data point to various genes against oxidative stress). This may explain the occurrence of these other B12-independent pathway factors even in the absence of CarA, as in Cystobacter, and opens the possibility that these other factors may have emerged prior to the appearance of from ancestral B<sub>12</sub>-dependent CarA an CarH photoreceptor.

### **Experimental procedures**

# Bacterial strains, growth conditions and complementation analysis

Bacterial strains, their source and plasmids used in this study are listed in Table S3. E. coli strain DH5a used for plasmid constructs, BL21(DE3) for protein overexpression, and BTH101 (cya<sup>-</sup>) for BACTH analysis, were grown in LB (Luria-Bertani) liquid medium or LBagar (1.5%) plates at 37°C or 25°C/18°C for inducing Hise-tagged protein overexpression. M. xanthus was grown at 33°C in CTT medium (10 g  $L^{-1}$  Bacto-casitone, 10 mM Tris-HCl pH 7.6, 1 mM KH<sub>2</sub>PO<sub>4</sub>, 8 mM MgSO<sub>4</sub>), as reported previously (Ortiz-Guerrero et al., 2011). S. cellulosum was grown at 33°C in M medium (Müller and Gerth, 2006). C. fuscus, C. ferrugineus and *M.* boletus were grown at  $30^{\circ}$ C in 3 g L<sup>-1</sup> Bactocasitone, 1 g  $L^{-1}$  glucose, pH 7.2, and H. ochraceum at 30°C on CY 1:5 plates (0.6 g L<sup>-1</sup> Bacto-casitone, 0.2 g L<sup>-1</sup> yeast extract, 0.27 g L<sup>-1</sup> CaCl<sub>2</sub>·2H<sub>2</sub>O, 21.1 g L<sup>-1</sup> NaCl, 0.6 g L<sup>-1</sup> KCl, 3.6 g L<sup>-1</sup> MgCl<sub>2</sub>·6H<sub>2</sub>O, 0.09 g L<sup>-1</sup> NaHCO<sub>3</sub>, 2.6 g L<sup>-1</sup> MgSO<sub>4</sub>.7H<sub>2</sub>O, 15 g L<sup>-1</sup> Difco agar, pH 7.2). Media were supplemented with 1  $\mu$ g ml<sup>-1</sup> of vitamin B<sub>12</sub> (which is converted intracellularly to AdoCbl) or AdoCbl and, as required, grown under light using three 18-W fluorescent lamps (10 Wm<sup>-2</sup>). Genes for CarA<sub>Av</sub>, CarH<sub>Av</sub>, CarH<sub>Ho</sub>, CarH<sub>Po</sub>, CSc and Cho were synthesized (GenScript). The others examined were PCR-amplified from genomic DNA purchased from DSMZ, Germany (C. coralloides), kindly provided by Prof. Rolf Müller (M. fulvus), or isolated using Wizard genomic DNA purification kit (Promega) from cells (C. fuscus, C. ferrugineus and M. boletus) grown as described above. For complementation analysis, the gene of interest was cloned into

# 18 R. Pérez-Castaño et al.

plasmid pMR3679 (Km<sup>R</sup>), which allows conditional expression from a vanillate-inducible promoter, tunable by inducer concentration (Iniesta et al., 2012). The construct, verified by sequencing, was electroporated into M. xanthus strain MR2649, which harbours a Tn5-lac-132 insertion in the carB operon to serve as a P<sub>crt Mx</sub>:: lacZ reporter probe and lacks endogenous CarH, CarA, CarS and to allow controlled supply of exogenous Ado-Cbl, PduO (Ortiz-Guerrero et al., 2011). Transformants with plasmid integration at a heterologous site by homoloaous recombination were selected using the Km<sup>R</sup> marker. M. xanthus cells were grown to exponential phase (OD<sub>600</sub> ~ 0.8) in the dark in liquid CTT media with vanillate (2-500 µM depending on the CarH homologue), antibiotics and 1  $\mu$ g ml<sup>-1</sup> of AdoCbl as required, diluted into fresh medium of the same composition and grown to OD<sub>600</sub> ~ 0.8, then split into two cultures, one for growth in the dark and the other under light for 8 h. Cells were spotted (5 µl) on plates of the same composition and grown under the same conditions, or collected for Pcrt <sub>Mx</sub>::lacZ β-galactosidase specific activity measurements (in nmol o-nitrophenyl-B-D-galactoside hydrolyzed/minmg<sup>-1</sup> protein reported as the mean and standard error of three biological replicates) in a SpectraMax 340 microtitre plate reader (Molecular Devices), as described previously (Ortiz-Guerrero et al., 2011).

### Genome and phylogenetic analysis

Specific M. xanthus proteins were used as queries in BLASTP analysis of myxobacterial genomes at the JGI integrated microbial genomes (IMG) resource (https://jgi. doe.gov/data-and-tools/img) to identify myxobacterial orthologs in the light response and in the MVA pathway as well as their genome contexts. The default E-value cut-off (1E-5) was used in all the analyses (Table S2 lists percent identities, E-values and other details). Among those retrieved with CarA<sub>Mx</sub> or CarH<sub>Mx</sub> as queries against each genome, the best hits were assigned as true homologues, and all had E-values ≤ 1E-17, percent identities ≥ 29%, contained an N-terminal MerR family DBD and a C-terminal CBD, and were in the 286-330 amino acid residue size range. Myxobacterial carotenoid, isoprenoid and cobalamin biosynthetic pathways were additionally analysed using the KEGG pathways database (https:// www.genome.jp/kegg), BLASTP (https://blast.ncbi.nlm. nih.gov/Blast.cgi), and Conserved Domains search tool (https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi).

Protein sequence identities were visualized with Heatmapper (http://www.heatmapper.ca/). Maximum likelihood phylogenetic trees were generated using MEGA7 (Molecular Evolutionary Genetics Analysis version 7.0; https://www.megasoftware.net), the JTT (Jones–Taylor– Thornton) substitution model and 200 bootstrap iterations after alignment of CarH/A homologues with MUSCLE (in MEGA7) or of 16S rRNA sequences for Myxococcales species with MAFFT (version 7.0: https://www.ebi.ac.uk/ Tools/msa). Trees were visualized and displayed using FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree) and iTOL (version 5: https://itol.embl.de). Putative CarH DNA-binding sites were identified by visual inspection of regions upstream of candidate target genes: and bioinformatic searches between positions -150 and + 25 relative to the annotated translation start codon for sites conforming to three 11-bp direct repeats with an N<sub>4</sub>-T-N<sub>3</sub>-C[C][G] consensus sequence (N is any base), and at least one N<sub>4</sub>-T-N<sub>2</sub>-RCR[G] repeat (R is A or G) using R packages: Biostrings (https://bioconductor.org/packages/ Biostrings), IRanges (Lawrence et al., 2013), dyplr (https://CRAN.R-project.org/package=dplyr) and SeginR (Charif and Lobry, 2007). Genomic data downloaded from Ensembl Bacteria database in FASTA and GFF3 formats were appropriately converted for data handling. WebLogo (https://weblogo.berkeley.edu/logo.cgi) was used to generate logos from sequences aligned with MUSCLE (gaps removed in MEGA7) or ClustalOmega (https://www.ebi.ac.uk/Tools/msa).

### Quantitative RT-PCR

RNA was isolated from 5 ml of myxobacterial cultures grown in liquid media with or without 1  $\mu$ g ml<sup>-1</sup> of vitamin B<sub>12</sub> in the dark or under light, reverse transcribed and analysed by qRT-PCR, as described in detail previously (Fernández-Zapata *et al.*, 2018). Primers used (Table S4) were designed using Primer Express 3.0 software to amplify an ~ 50–150 bp region within each transcript. Target gene expression is reported relative to *rpoD* as reference gene and normalized to the level of expression of the gene in the dark with vitamin B<sub>12</sub> present, as the mean and standard error for three biological replicates.

# BACTH analysis

Two-hybrid analysis of light and B<sub>12</sub>-dependent oligomerization employed the BACTH system (based on protein fusion to the T25 and T18 fragments of *Bordetella pertussis* adenylate cyclase), as described previously (Ortiz-Guerrero *et al.*, 2011). Self-interaction was inferred from the blue colour of cell spots on 40 µg ml<sup>-1</sup> of X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside)-LB plates with or without 1 µg ml<sup>-1</sup> of vitamin B<sub>12</sub> grown in the dark or under light and, as controls, CarH<sub>Mx</sub> and CarA<sub>Mx</sub>. Pairs with just one fusion protein were used as negative controls and no autoactivation was observed, as shown in Fig. S3 for CarH<sub>Cfu</sub>, CarH<sub>Cfe</sub> and CarH<sub>Mb</sub>.

### Protein purification and characterization

Hise-tagged proteins were overexpressed in E. coli BL21 (DE3) using pET15b constructs, purified and stored as reported before (López-Rubio et al., 2002; Ortiz-Guerrero et al., 2011). Native apoproteins purified in a final sizeexclusion chromatography step (in 150 mM NaCl, 50 mM phosphate buffer, pH 7.5, 2 mM  $\beta$ -mercaptoethanol) were concentrated and their identities checked by SDS-PAGE and electrospray ionization-time-of flight mass spectrometry (Agilent 1100 Series HPLC, Supelco Discovery Bio-Wide Pore C5 2.1  $\times$  10 cm, 5  $\mu m$  HPLC column, Ion Trap XCT Plus mass spectrometer with an electrospray interface; mass spectra in the positive ion mode, m/zrange 100 to 2200 for samples separated using a linear 4%-90% water/acetonitrile/formic acid gradient). AdoCbl was always handled under dim light and checked by UVvisible (UV-Vis) spectroscopy. Protein concentrations were determined from absorbance at 280 nm in a Cary 60 spectrophotometer, using molar extinction coefficients  $\epsilon_{280}$  (in M<sup>-1</sup> cm<sup>-1</sup>) of 37,930, 36,440, 30,480, 32,430 and 30,940 for CarH<sub>Cfu</sub>, CarH<sub>Cfe</sub>, CarS<sub>Cfu</sub>, CSc and Cho, respectively, and the Bio-Rad protein assay kit.

# UV–Vis absorption spectroscopy and analytical size exclusion chromatography

UV-visible spectra for AdoCbl-bound CarH<sub>Cfu</sub> and Car-H<sub>Cfe</sub> photoconversion were recorded for samples in the dark and with stepwise green light irradiation (522 nm, 10 Wm<sup>-2</sup>, from a computer-controlled LED array) from 1 to 450 s. Analytical size-exclusion chromatography to estimate the apparent molecular mass  $(M_r)$  was performed in an AKTA HPLC unit using a Superdex200 HR 10/30 column and calibrated yielding log  $M_r = 8.1537$ -0.2316  $V_e$ , where  $V_e$  is the elution volume. Pure protein (100  $\mu$ l, 50–100  $\mu$ M), alone or after 15-min incubation with five-fold molar excess of AdoCbl was injected into the column in the dark or after 5-min green light exposure. Elution (0.4 ml min<sup>-1</sup> flow) was tracked by absorbance at 280, 361 and 522 nm, the peaks collected and analysed for  $M_r$ , and by UV-Vis spectroscopy and SDS-PAGE.

#### DNA binding assays

DNA probes > 100 bp were obtained by PCR using appropriate primers, one of them <sup>32</sup>P-labelled at the 5'end with T4 polynucleotide kinase (Takara). For DNA probes  $\leq$  63 bp, complementary HPLC-purified synthetic oligonucleotides (Merck), one <sup>32</sup>P-labelled at the 5'-end and the other unlabeled, were mixed at a 1:2 ratio, incubated at 100°C for 2 min and hybridized by slow cooling.

### Photoregulated carotenogenesis in myxobacteria 19

EMSA was carried out using 20 ul of samples containing the DNA probe (1.2 nM, ~13,000 cpm) and protein at specific concentrations, with or without a 5-fold excess of AdoCbl relative to protein in 0.1 M KCl, 10 mM MgCl<sub>2</sub>, 0.025 M Tris-HCl (pH 8), 1 mM dithiothreitol, 10% glycerol. 200 ng  $\mu$ l<sup>-1</sup> of bovine serum albumin and 1  $\mu$ g sheared salmon sperm DNA as non-specific competitor. Samples were incubated at 37°C for 30 min and loaded onto a 6% non-denaturing polyacrylamide gel (37.5:1 acrylamide: bisacrylamide) pre-run for 30 min in  $0.5 \times$ TBE buffer (45 mM Tris base, 45 mM boric acid, 1 mM EDTA). All steps were carried out in the dark (under dim light) except when, prior to electrophoresis, samples were exposed for 5 min to green light. Electrophoresis was carried out in the dark for 1.5 h at 200 V and 10°C. Gels were vacuum-dried and scanned using Personal Molecular Imager<sup>TM</sup> (PMI<sup>TM</sup>) FX with Quantity One 4.4 software (BioRad). The fraction bound, estimated from band intensities (quantified using ImageJ, NIH) for the free DNA, was fit to a 3-parameter Hill equation using SigmaPlot (Systat Software) to estimate the apparent equilibrium dissociation constant ( $K_D$ , the protein concentration at half-maximal binding) and Hill coefficient. For DNase I footprinting, 20 µl of samples of a DNA probe (~20,000 cpm with either the sense or the antisense strand 5'-end <sup>32</sup>P-labelled) and protein with AdoCbl at a 5-fold excess in EMSA buffer lacking glycerol were incubated for 30 min at 37°C, then treated with 0.07 units of DNase I (Promega) at 37°C for 2 min and guenched with 25 mM EDTA, all in the dark. DNA was ethanol-precipitated, washed twice with 70% ethanol and dried. For hydroxyl radical footprinting, samples were prepared as for DNase I footprinting but without MgCl<sub>2</sub>, and then treated for 4 min at 25°C with 2 µl each of freshly prepared Fe(II)-EDTA solution (1 mM ammonium iron (II) sulfate, 2 mM EDTA), 0.01 M sodium ascorbate and 0.6% hydrogen peroxide (Merck). The reaction was stopped with 2 µl each of 0.1 M thiourea and 0.5 M EDTA (pH 8), and DNA precipitated and dried as before. For Exo III (3'  $\rightarrow$  5' exodeoxyribonuclease; ThermoFisher Scientific) footprinting, 20 µl samples of the DNA probe 5'-end <sup>32</sup>P-labelled on a specific strand were mixed with protein and a 5-fold excess of AdoCbl, incubated in DNase I buffer for 30 min in the dark, then treated with 2.5–5 U  $\mu$ l<sup>-1</sup> of Exo III for 15 min at 37°C, quenched with 20 mM EDTA, DNA precipitated and dried. Each dried DNA sample for footprinting assays was resuspended in 5  $\mu$ l of formamide loading buffer, incubated at 95°C for 3 min, and loaded onto 6% polyacrylamide-8 M urea sequencing gels. After electrophoresis (2-3 h, 50 mA/60 W/1.90 kV), the gels were vacuum-dried and scanned using PMI<sup>TM</sup> (BioRad). A + G chemical sequencing ladders were used to map the footprints.

### Acknowledgements

This work was supported by the Ministerio de Ciencia, Innovación y Universidades (MCIU)/Agencia Estatal de Investigación (AEI)/European Regional Development Fund (ERDF): grants PGC2018-094635-B-C21 to M.E.-A. and PGC2018-094635-B-C22 to S.P, both funded by MCIU/AEI 10.13039/501100011033 and by 'ERDF A way of making Europe'; by grant 20992/PI/18 to M.E.-A. from Fundación Séneca (Murcia)-Spain; by Ph.D. fellowship contracts to R.P.-C (BES-2016-076883, MINECO/AEI), to J. F.-Z. (BES-2013-063376, MINECO), and to E.B.-M. (FPU17/05685, Ministerio de Educación, Cultura y Deporte). We thank J.A. Madrid and V. López-Egea for technical assistance.

# References

- Abellón-Ruiz, J., Bernal-Bernal, D., Abellán, M., Fontes, M., Padmanabhan, S., Murillo, F.J., and Elías-Arnanz, M. (2014) The CarD/CarG regulatory complex is required for the action of several members of the large set of *Myxococcus xanthus* extracytoplasmic function sigma factors. *Environ Microbiol* **16**: 2475–2490.
- Armstrong, G.A. (1997) Genetics of eubacterial carotenoid biosynthesis: a colorful tale. *Annu Rev Microbiol* 51: 629–659.
- Bernal-Bernal, D., Abellón-Ruiz, J., Iniesta, A.A., Pajares-Martínez, E., Bastida-Martínez, E., Fontes, M., *et al.* (2018) Multifactorial control of the expression of a CRISPR-Cas system by an extracytoplasmic function  $\sigma$ /anti- $\sigma$  pair and a global regulatory complex. *Nucleic Acids Res* **46**: 6726–6745.
- Browning, D.F., Whitworth, D.E., and Hodgson, D.A. (2003) Light-induced carotenogenesis in *Myxococcus xanthus*: functional characterization of the ECF sigma factor CarQ and antisigma factor CarR. *Mol Microbiol* **48**: 237–251.
- Burchard, R.P., and Dworkin, M. (1966) Light-induced lysis and carotenogenesis in *Myxococcus xanthus*. *J Bacteriol* **91**: 535–545.
- Camacho, I.S., Black, R., Heyes, D.J., Johannissen, L.O., Ramakers, L.A.I., Bellina, B., *et al.* (2021) Interplay between chromophore binding and domain assembly by the B<sub>12</sub>-dependent photoreceptor protein, CarH. *Chem Sci* **12**: 8333–8341.
- Cao, P., and Wall, D. (2019) Direct visualization of a molecular handshake that governs kin recognition and tissue formation in myxobacteria. *Nat Commun* **10**: 3073.
- Cayuela, M.L., Elías-Arnanz, M., Peñalver-Mellado, M., Padmanabhan, S., and Murillo, F.J. (2003) The *Stigmatella aurantiaca* homolog of *Myxococcus xanthus* high-mobility-group A-type transcription factor CarD: insights into the functional modules of CarD and their distribution in bacteria. *J Bacteriol* **185**: 3527–3537.
- Chapal, M., Mintzer, S., Brodsky, S., Carmi, M., and Barkai, N. (2019) Resolving noise-control conflict by gene duplication. *PLoS Biol* **17**: e3000289.
- Charif, D., and Lobry, J.R. (2007) SeqinR 1.0–2: a contributed package to the R project for statistical computing devoted to biological sequences retrieval and analysis. In *Structural Approaches to Sequence Evolution: Molecules, Networks, Populations*, Bastolla, U., Porto, M., Roman, H.

E., and Vendruscolo, M. (eds). Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 207–232.

- Díez, A.I., Ortiz-Guerrero, J.M., Ortega, A., Elías-Arnanz, M., Padmanabhan, S., and García de la Torre, J. (2013) Analytical ultracentrifugation studies of oligomerization and DNA-binding of TtCarH, a *Thermus thermophilus* coenzyme B<sub>12</sub>-based photosensory regulator. *Eur Biophys J* **42**: 463–476.
- Elías-Arnanz, M., Padmanabhan, S., and Murillo, F.J. (2010) The regulatory action of the myxobacterial CarD/CarG complex: a bacterial enhanceosome? *FEMS Microbiol Rev* **34**: 764–778.
- Elías-Arnanz, M., Padmanabhan, S., and Murillo, F.J. (2011) Light-dependent gene regulation in nonphototrophic bacteria. *Curr Opin Microbiol* **14**: 128–135.
- Fernández-Zapata, J., Pérez-Castaño, R., Aranda, J., Colizzi, F., Polanco, M.C., Orozco, M., *et al.* (2018) Plasticity in oligomerization, operator architecture, and DNA binding in the mode of action of a bacterial B<sub>12</sub>-based photoreceptor. *J Biol Chem* **293**: 17888–17905.
- Fontes, M., Galbis-Martínez, L., and Murillo, F.J. (2003) A novel regulatory gene for light-induced carotenoid synthesis in the bacterium *Myxococcus xanthus*. *Mol Microbiol* **47**: 561–571.
- Galbis-Martínez, M., Padmanabhan, S., Murillo, F.J., and Elías-Arnanz, M. (2012) CarF mediates signaling by singlet oxygen, generated via photoexcited protoporphyrin IX, in *Myxococcus xanthus* light-induced carotenogenesis. *J Bacteriol* **194**: 1427–1436.
- Gallego-García, A., Monera-Girona, A.J., Pajares-Martínez, E., Bastida-Martínez, E., Pérez-Castaño, R., Iniesta, A.A., *et al.* (2019) A bacterial light response reveals an orphan desaturase for human plasmalogen synthesis. *Science* **366**: 128–132.
- García-Heras, F., Padmanabhan, S., Murillo, F.J., and Elías-Arnanz, M. (2009) Functional equivalence of HMGA- and histone H1-like domains in a bacterial transcriptional factor. *Proc Natl Acad Sci U S A* **106**: 13546–13551.
- García-Moreno, D., Abellón-Ruiz, J., García-Heras, F., Murillo, F.J., Padmanabhan, S., and Elías-Arnanz, M. (2010) CdnL, a member of the large CarD-like family of bacterial proteins, is vital for *Myxococcus xanthus* and differs functionally from the global transcriptional regulator CarD. *Nucleic Acids Res* **38**: 4586–4598.
- Glaeser, J., Nuss, A.M., Berghoff, B.A., and Klug, G. (2011) Singlet oxygen stress in microorganisms. *Adv Microb Physiol* **58**: 141–173.
- Goldman, B.S., Nierman, W.C., Kaiser, D., Slater, S.C., Durkin, A.S., Eisen, J.A., *et al.* (2006) Evolution of sensory complexity recorded in a myxobacterial genome. *Proc Natl Acad Sci U S A* **103**: 15200–15205.
- Hittinger, C.T., and Carroll, S.B. (2007) Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* **449**: 677–681.
- Hoshino, Y., and Gaucher, E.A. (2021) Evolution of bacterial steroid biosynthesis and its impact on eukaryogenesis. *Proc Natl Acad Sci U S A* **118**: e2101276118.
- Huntley, S., Hamann, N., Wegener-Feldbrugge, S., Treuner-Lange, A., Kube, M., Reinhardt, R., *et al.* (2011) Comparative genomic analysis of fruiting body formation in Myxococcales. *Mol Biol Evol* 28: 1083–1097.

- Iniesta, A.A., Cervantes, M., and Murillo, F.J. (2007) Cooperation of two carotene desaturases in the production of lycopene in *Myxococcus xanthus*. *FEBS J* 274: 4306– 4314.
- Iniesta, A.A., Cervantes, M., and Murillo, F.J. (2008) Conversion of the lycopene monocyclase of *Myxococcus xanthus* into a bicyclase. *Appl Microbiol Biotechnol* **79**: 793–802.
- Iniesta, A.A., García-Heras, F., Abellón-Ruiz, J., Gallego-García, A., and Elías-Arnanz, M. (2012) Two systems for conditional gene expression in *Myxococcus xanthus* inducible by isopropyl-β-D-thiogalactopyranoside or vanillate. *J Bacteriol* **194**: 5875–5885.
- Jost, M., Fernández-Zapata, J., Polanco, M.C., Ortiz-Guerrero, J.M., Chen, P.Y., Kang, G., *et al.* (2015b) Structural basis for gene regulation by a B<sub>12</sub>-dependent photoreceptor. *Nature* **526**: 536–541.
- Jost, M., Simpson, J.H., and Drennan, C.L. (2015a) The transcription factor CarH safeguards use of adenosylcobalamin as a light sensor by altering the photolysis products. *Biochemistry* **54**: 3231–3234.
- Kasinsky, H.E., Lewis, J.D., Dacks, J.B., and Ausió, J. (2001) Origin of H1 linker histones. *FASEB J* **15**: 34–42.
- Kohler, C., Lourenço, R.F., Avelar, G.M., and Gomes, S.L. (2012) Extracytoplasmic function (ECF) sigma factor σ<sup>F</sup> is involved in *Caulobacter crescentus* response to heavy metal stress. *BMC Microbiol* **12**: 210.
- Kutta, R.J., Hardman, S.J., Johannissen, L.O., Bellina, B., Messiha, H.L., Ortiz-Guerrero, J.M., *et al.* (2015) The photochemical mechanism of a B<sub>12</sub>-dependent photoreceptor protein. *Nat Commun* 6: 7907.
- Kuzmin, E., VanderSluis, B., Nguyen Ba, A.N., Wang, W., Koch, E.N., Usaj, M., *et al.* (2020) Exploring wholegenome duplicate gene retention with complex genetic interaction analysis. *Science* **368**: eaaz5667.
- Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., *et al.* (2013) Software for computing and annotating genomic ranges. *PLoS Comput Biol* **9**: e1003118.
- León, E., Navarro-Avilés, G., Santiveri, C.M., Flores-Flores, C., Rico, M., González, C., *et al.* (2010) A bacterial antirepressor with SH3 domain topology mimics operator DNA in sequestering the repressor DNA recognition helix. *Nucleic Acids Res* **38**: 5226–5241.
- López-García, P., and Moreira, D. (2020) The Syntrophy hypothesis for the origin of eukaryotes revisited. *Nat Microbiol* **5**: 655–667.
- López-Rubio, J.J., Elías-Arnanz, M., Padmanabhan, S., and Murillo, F.J. (2002) A repressor-antirepressor pair links two loci controlling light-induced carotenogenesis in *Myxococcus xanthus. J Biol Chem* **277**: 7262–7270.
- López-Rubio, J.J., Padmanabhan, S., Lázaro, J.M., Salas, M., Murillo, F.J., and Elías-Arnanz, M. (2004) Operator design and mechanism for CarA repressor-mediated down-regulation of the photoinducible *carB* operon in *Myxococcus xanthus*. *J Biol Chem* **279**: 28945–28953.
- Maresca, J.A., Graham, J.E., Wu, M., Eisen, J.A., and Bryant, D.A. (2007) Identification of a fourth family of lycopene cyclases in photosynthetic bacteria. *Proc Natl Acad Sci U S A* **104**: 11784–11789.

- Marino-Puertas, L., Goulas, T., and Gomis-Rüth, F.X. (2017) Matrix metalloproteinases outside vertebrates. *Biochim Biophys Acta Mol Cell Res* **1864**: 2026–2035.
- Martínez-Argudo, I., Ruiz-Vázquez, R.M., and Murillo, F.J. (1998) The structure of an ECF-sigma-dependent, lightinducible promoter from the bacterium *Myxococcus xanthus. Mol Microbiol* **30**: 883–893.
- Masloboeva, N., Reutimann, L., Stiefel, P., Follador, R., Leimer, N., Hennecke, H., *et al.* (2012) Reactive oxygen species-inducible ECF  $\sigma$  factors of *Bradyrhizobium japonicum*. *PLoS One* **7**: e43421.
- Meiser, P., Bode, H.B., and Muller, R. (2006) The unique DKxanthene secondary metabolite family from the myxobacterium *Myxococcus xanthus* is required for developmental sporulation. *Proc Natl Acad Sci U S A* **103**: 19128–19133.
- Miller, N.A., Kaneshiro, A.K., Konar, A., Alonso-Mori, R., Britz, A., Deb, A., *et al.* (2020) The photoactive excited state of the B<sub>12</sub>-based photoreceptor CarH. *J Phys Chem B* **124**: 10732–10738.
- Moise, A.R., Al-Babili, S., and Wurtzel, E.T. (2014) Mechanistic aspects of carotenoid biosynthesis. *Chem Rev* **114**: 164–193.
- Monteverde, D.R., Gómez-Consarnau, L., Suffridge, C., and Sañudo-Wilhelmy, S.A. (2017) Life's utilization of B vitamins on early Earth. *Geobiology* **15**: 3–18.
- Moraleda-Muñoz, A., Pérez, J., Fontes, M., Murillo, F.J., and Muñoz-Dorado, J. (2005) Copper induction of carotenoid synthesis in the bacterium *Myxococcus xanthus*. *Mol Microbiol* 56: 1159–1168.
- Müller, R., and Gerth, K. (2006) Development of simple media which allow investigations into the global regulation of chivosazol biosynthesis with *Sorangium cellulosum* So ce56. *J Biotechnol* **121**: 192–200.
- Muñoz-Dorado, J., Marcos-Torres, F.J., García-Bravo, E., Moraleda-Muñoz, A., and Pérez, J. (2016) Myxobacteria: moving, killing, feeding, and surviving together. *Front Microbiol* **7**: 781.
- Navarro-Avilés, G., Jiménez, M.A., Pérez-Marín, M.C., González, C., Rico, M., Murillo, F.J., *et al.* (2007) Structural basis for operator and antirepressor recognition by *Myxococcus xanthus* CarA repressor. *Mol Microbiol* 63: 980–994.
- Ortiz-Guerrero, J.M., Polanco, M.C., Murillo, F.J., Padmanabhan, S., and Elías-Arnanz, M. (2011) Lightdependent gene regulation by a coenzyme B<sub>12</sub>-based photoreceptor. *Proc Natl Acad Sci U S A* **108**: 7565– 7570.
- Padmanabhan, S., Elias-Arnanz, M., Carpio, E., Aparicio, P., and Murillo, F.J. (2001) Domain architecture of a high mobility group A-type bacterial transcriptional factor. *J Biol Chem* 276: 41566–41575.
- Padmanabhan, S., Jost, M., Drennan, C.L., and Elías-Arnanz, M. (2017) A new facet of vitamin B<sub>12</sub>: gene regulation by cobalamin-based photoreceptors. *Annu Rev Biochem* 86: 485–514.
- Padmanabhan, S., Monera-Girona, A.J., Pérez-Castaño, R., Bastida-Martínez, E., Pajares-Martínez, E., Bernal-Bernal, D., *et al.* (2021) Light-triggered carotenogenesis in *Myxococcus xanthus*: new paradigms

### 22 R. Pérez-Castaño et al.

in photosensory signaling, transduction and gene regulation. *Microorganisms* **9**: 1067.

- Padmanabhan, S., Pérez-Castaño, R., and Elías-Arnanz, M. (2019) B<sub>12</sub>-based photoreceptors: from structure and function to applications in optogenetics and synthetic biology. *Curr Opin Struct Biol* **57**: 47–55.
- Peñalver-Mellado, M., García-Heras, F., Padmanabhan, S., García-Moreno, D., Murillo, F.J., and Elías-Arnanz, M. (2006) Recruitment of a novel zinc-bound transcriptional factor by a bacterial HMGA-type protein is required for regulating multiple processes in *Myxococcus xanthus*. *Mol Microbiol* **61**: 910–926.
- Perez, J.C., and Groisman, E.A. (2009) Evolution of transcriptional regulatory circuits in bacteria. *Cell* **138**: 233–244.
- Pérez-Marín, M.C., Padmanabhan, S., Polanco, M.C., Murillo, F.J., and Elías-Arnanz, M. (2008) Vitamin B<sub>12</sub> partners the CarH repressor to downregulate a photoinducible promoter in *Myxococcus xanthus*. *Mol Microbiol* 67: 804–819.
- Pillai, A.S., Chandler, S.A., Liu, Y., Signore, A.V., Cortez-Romero, C.R., Benesch, J.L.P., *et al.* (2020) Origin of complexity in haemoglobin evolution. *Nature* 581: 480–485.
- Rodriguez-Concepción, M., Avalos, J., Bonet, M.L., Boronat, A., Gómez-Gómez, L., Hornero-Mendez, D., *et al.* (2018) A global perspective on carotenoids: metabolism, biotechnology, and benefits for nutrition and health. *Prog Lipid Res* **70**: 62–93.
- Romero, I.G., Ruvinsky, I., and Gilad, Y. (2012) Comparative studies of gene expression and the evolution of gene regulation. *Nat Rev Genet* **13**: 505–516.
- Roth, J.R., Lawrence, J.G., and Bobik, T.A. (1996) Cobalamin (coenzyme B<sub>12</sub>): synthesis and biological significance. *Annu Rev Microbiol* **50**: 137–181.
- Sandmann, G. (2019) Antioxidant protection from UV- and light-stress related to carotenoid structures. *Antioxidants* (*Basel*) **8**: 219.
- Santana-Molina, C., Rivas-Marin, E., Rojas, A.M., and Devos, D.P. (2020) Origin and evolution of polycyclic triterpene synthesis. *Mol Biol Evol* **37**: 1925–1941.
- Schlotterer, C. (2015) Genes from scratch-the evolutionary fate of *de novo* genes. *Trends Genet* **31**: 215–219.

- Schneiker, S., Perlova, O., Kaiser, O., Gerth, K., Alici, A., Altmeyer, M.O., et al. (2007) Complete genome sequence of the myxobacterium Sorangium cellulosum. Nat Biotechnol 25: 1281–1289.
- Shelton, A.N., Seth, E.C., Mok, K.C., Han, A.W., Jackson, S. N., Haft, D.R., and Taga, M.E. (2019) Uneven distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative genomics. *ISME J* **13**: 789–804.
- Simon, S., Ruhl, M., de Montaigu, A., Wotzel, S., and Coupland, G. (2015) Evolution of CONSTANS regulation and function after gene duplication produced a photoperiodic flowering switch in the Brassicaceae. *Mol Biol Evol* 32: 2284–2301.
- Sokolovskaya, O.M., Shelton, A.N., and Taga, M.E. (2020) Sharing vitamins: Cobamides unveil microbial interactions. *Science* **369**: eaba0165.
- Van Oss, S.B., and Carvunis, A.R. (2019) *De novo* gene birth. *PLoS Genet* **15**: e1008160.
- Waite, D.W., Chuvochina, M., Pelikan, C., Parks, D.H., Yilmaz, P., Wagner, M., *et al.* (2020) Proposal to reclassify the proteobacterial classes *Deltaproteobacteria* and *Oligoflexia*, and the phylum *Thermodesulfobacteria* into four phyla reflecting major functional capabilities. *Int J Syst Evol Microbiol* **70**: 5972–6016.
- Yamamoto, E., Muramatsu, H., and Nagai, K. (2014) Vulgatibacter incomptus gen. Nov., sp. nov. and Labilithrix luteola gen. Nov., sp. nov., two myxobacteria isolated from soil in Yakushima Island, and the description of Vulgatibacteraceae fam. Nov., Labilitrichaceae fam. Nov. and Anaeromyxobacteraceae fam. Nov. Int J Syst Evol Microbiol 64: 3360–3368.
- Ziegelhoffer, E.C., and Donohue, T.J. (2009) Bacterial responses to photo-oxidative stress. *Nat Rev Microbiol* **7**: 856–863.

### **Supporting Information**

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

#### Appendix S1: Supporting Information

**Table S2**: Genome and gene product details formyxobacteria examined in this study