

Invited Review

Determinants of axonal regeneration

J. Frisé

Department of Neuroscience, Karolinska Institute, Stockholm, Sweden

Summary. Axons often regrow to their targets and lost functions may be restored after an injury in the peripheral nervous system. In contrast, axonal regeneration is generally very limited after injuries in the central nervous system, and functional impairment is usually permanent. The regenerative capacity depends on intrinsic neuronal factors as well as the interaction of neurons with other cells. Glial cells may, in different situations, either support or inhibit axonal growth. This review discusses the molecular mechanisms that are involved in promoting and inhibiting axonal regeneration in the nervous system after injuries.

Key words: Extracellular matrix, Macrophage, Nerve regeneration, Neurotrophin

Introduction

Transection of a nerve fiber triggers a complex set of events in the nerve cell body, in the axon proximal and distal to the lesion, in supporting glial cells and other non-neuronal cells. The neuronal perikaryon swells, the nucleus is displaced against the cell membrane, and the Nissl substance disintegrates, a sign of increased protein synthesis. This neuronal reaction to axotomy occurs to a different extent in different types of neurons and also depends on factors such as the distance to the cell body. Many proteins normally synthesized by a neuron will no longer be produced, or will be produced at low levels after axotomy, while other proteins that are normally synthesized only at low levels or not at all, will be expressed at high levels. This shift in gene expression in response to injury probably occurs, at least in part, in order to optimize the neuronal capacity for regeneration (reviewed in Aldskogius and Svensson, 1993). After transection of a nerve fiber the axonal membrane will rapidly seal and an end swelling, the growth cone, forms. Thin filopodia emerging from the axonal endbulb can sense molecular cues in the environment and transmit signals to the growth cone, presumably resulting in the decision of which path to choose (Davenport et al.,

1993). When an outgrowing axon has reestablished contact with the target, neuronal protein synthesis will return to normal. The alteration in neuronal molecular phenotype after axotomy seems to be triggered by the disrupted retrograde transport of target derived factor(s), since the phenotype can be reversed toward the normal if the required trophic factor is supplied (see e.g. Fitzgerald et al., 1985). However, axons often fail to regrow and reinnervate the target, and moreover, neurons may even die in response to transection of the axon. The susceptibility of a neuron to death after transection of the axon is greater during development than in adulthood, probably as a result of decreasing dependence on trophic support for survival in the adult animal.

While regeneration in the peripheral nervous system (PNS) is often rapid and successful, regeneration in the adult mammalian central nervous system (CNS) is generally not seen. After an injury in the PNS, debris is rapidly cleared from the denervated distal stump, through a process known as Wallerian degeneration. The Schwann cells in the distal nerve stump will line up to form channels, known as the bands of Bügner, through which the severed axons can regrow to their targets (Fig. 1). In contrast, in the injured CNS, axon-growth is in most cases limited to the scar that will be formed by astrocytes at the site of the injury. The Wallerian degeneration distal to a lesion in the CNS is much slower than in the PNS, and a dense scar will form there with time (Fig. 1). There may be important intrinsic differences between PNS and CNS neurons in response to injury. However, non-neuronal factors seem to be at least as important. Here follows an overview of some of the suggested key players in promoting and inhibiting axonal regeneration. This review focuses on differences between the adult mammalian PNS and CNS in their response to traumatic injuries. Factors that are likely to be key determinants of the capacity of axonal regeneration are discussed, and a few examples are mentioned in each case.

Intrinsic neuronal regenerative capacity

Aguayo and collaborators demonstrated that axons of CNS neurons can grow long distances through PNS tissue by grafting segments of peripheral nerve to the

injured brain, spinal cord, and optic nerve (Reviewed by Aguayo et al., 1991). This clearly demonstrated that the lack of regeneration within the CNS is not due entirely to a weak intrinsic neuronal regenerative capacity. Instead, this indicates that the CNS environment fails to stimulate axon regrowth or that the CNS contains axon-regrowth inhibiting molecules. The latter theory was also supported by the lack of PNS axon-growth in a CNS graft (Aguayo et al., 1978). However, intrinsic neuronal factors may indeed limit axonal regrowth in the CNS in some situations (reviewed by Fawcett, 1992). For example, the capacity of CNS neurons to innervate a peripheral nerve graft inserted in the injured spinal cord, brain or optic nerve decreases with increasing distance between the neuronal cell body and the injury (David and Aguayo, 1981; Benfey and Aguayo, 1982; Richardson et al., 1984; Vidal-Sanz et al., 1987). This is probably a result of a weaker triggering of regenerative programs in the neuron when the lesion is far from the cell body, as indicated by the fact that the growth-associated protein GAP-43 is induced in retinal ganglion cells only if the axotomy is within the first mm of the optic nerve (Doster et al., 1991). The distance between the cell body and the site of axotomy does not seem to be an important determinant of axonal regeneration in the PNS, since axons can regrow successfully even after the axon is cut close to the target. Other intrinsic

neuronal differences that may render CNS neurons less likely than PNS neurons to regenerate successfully may be related to the facts that central neurons are more likely to die after axotomy than PNS nerve cells (Sunderland, 1978), and that the formation of ectopic terminals on nearby neurons may impede axonal regrowth in the CNS (Bernstein and Bernstein, 1971).

That axons from neurons in the brain and spinal cord can grow in peripheral nerve grafts indicates that the extracellular environment provided by glial and other cell types may be crucial for the regeneration process. In different situations non-neuronal cells may either stimulate axon-growth by e.g. secreting neurotrophic molecules or extracellular matrix molecules or actively inhibit axonal regeneration by repelling growth cones.

Neurotrophic molecules

Much of the research on the role of neurotrophic factors in regeneration has focused on the neurotrophin family. This family consists of the prototypical neurotrophic factor, nerve growth factor (NGF) and the structurally and functionally related molecules brain-derived neurotrophic factor (BDNF), neurotrophin 3 (NT-3) and NT-4. The neurotrophins exert their functions by binding to specific cell surface receptors. The first identified neurotrophin-binding molecule is a widely expressed protein which can bind all neurotrophins with low affinity (designated p75 or low-affinity neurotrophin receptor). However, a group of tyrosine kinase receptors denoted trkA, trkB and trkC seem to be more important for mediating the effects of neurotrophins. NGF binds to trkA, BDNF and NT-4 selectively activate trkB, and the physiological receptor for NT-3 seems to be trkC (reviewed by Barbacid, 1995). The existence of alternative forms of the trkA, trkB and trkC tyrosine kinase receptors, as well as truncated forms of trkB and trkC which lack the tyrosine kinase domain add an extra level of complexity. The neuro-trophins, by activating their cognate receptors, stimulate differentiation and neurite outgrowth, and support the survival of different, but overlapping, groups of neurons (reviewed by Snider, 1994).

There is a wealth of circumstantial evidence implicating neurotrophins in PNS regeneration. When a peripheral nerve is transected, the normally low levels of NGF, BDNF and NT-4 in the nerve increase dramatically in the distal nerve stump (Heumann et al., 1987a; Meyer et al., 1992; Funakoshi et al., 1993). Moreover, some neurotrophins are expressed at increased levels in target tissues after denervation, i.e. BDNF mRNA expression is strongly elevated in muscle (Funakoshi et al., 1993; Koliatsos et al., 1993) and NGF mRNA levels increase in skin (Mearow et al., 1993).

Schwann cells normally express significant levels of truncated trkB and trkC receptors lacking the tyrosine kinase domain, but very low levels of p75. This expression pattern of neurotrophin receptors is reversed in denervated Schwann cells so that high levels of p75




Fig. 1. Schematic drawing depicting the main cellular reactions after injuries in the PNS and CNS. In the PNS, the Schwann cells in the distal nerve stump line up to form channels through which the axons can regrow. In the CNS, clearance of axonal debris and myelin is slow distal to the injury, and astrocytes form a dense scar. The axonal growth in the injured CNS is limited to the site of the lesion.

Determinants of axonal regeneration

are expressed and *trkB* and *trkC* expression is strongly decreased (Taniuchi et al., 1986, 1988; Heumann et al., 1987b; Frisén et al., 1993; Funakoshi et al., 1993). See Fig. 2 for a summary of changes in neurotrophin and neurotrophin receptor expression after injury.

It has been suggested that p75 can bind NGF on the surface of reactive Schwann cells and present NGF to regrowing axons (reviewed by Johnson et al., 1988). Indeed, NGF bound to distal segments of previously severed sciatic nerves can enhance sensory neurite growth in vitro (Sandrock and Matthew, 1987a). Since p75 can bind all neurotrophins with similar affinity, it is reasonable to extend this hypothesis to the other neurotrophins. The decrease in *trkB* and *trkC* mRNA in the distal nerve segment after transection limits competition for ligand, thus enabling binding of neurotrophins with low affinity to Schwann cell surfaces.

Most studies regarding the role of neurotrophins in peripheral nerve regeneration have hitherto focused on NGF sensitive nerve cells, such as sensory and sympathetic neurons. The weak sensory axon regeneration in C57BL/Ola mice (a mouse strain in which macrophage recruitment is virtually absent after injury and NGF-synthesis increases only marginally after injury) can be stimulated by exogenous NGF, supporting a role for NGF in sensory axon regrowth (Brown et al., 1991). Moreover, the rate of sensory axon regeneration through silastic tubes has been reported to increase when NGF is supplied (Derby et al., 1993). However, other reports suggest a lesser role for NGF in nerve regeneration. p75 and *trkA* expression as well as retrograde NGF-transport decrease in dorsal root ganglia neurons after axotomy (Raivich et al., 1991; Verge et al., 1992), and regeneration of sensory axons is not impeded when endogenous NGF is inactivated with anti-NGF antibodies (Rich et al., 1984; Diamond et al., 1987, 1992). Additionally, it does not appear that NGF is

involved in survival of sensory neurons after injury (Rich et al., 1984) or regrowth of lesioned sympathetic axons (Gloster and Diamond, 1992). Thus, although NGF is synthesized at highly elevated levels after injury, it does not seem to be involved in sensory or sympathetic axon regeneration. However, data have been presented which indicate that NGF is involved in lesion-induced collateral sprouting of undamaged sensory neurons (which increase their p75 mRNA expression) and sympathetic neurons (Mearow et al., 1991; Diamond et al., 1992b; Gloster and Diamond, 1992). The decrease in p75 mRNA in sensory neurons after axotomy may indicate that neurotrophins are not involved in sensory axon regeneration. However, *trkB* and *trkC* mRNA are increased in dorsal root ganglia neurons after axotomy (Ernfors et al., 1993) suggesting that BDNF, NT-3, and/or NT-4 could be important for sensory neurons after injury.

BDNF promotes the survival of axotomized motoneurons in neonatal rats (Sendtner et al., 1992a; Yan et al., 1992; Koliatsos et al., 1993). In contrast to *trkA* mRNA in sensory neurons, *trkB* expression by motoneurons increases after axotomy (Piehl et al., 1994; Kobayashi et al., 1996). The increase in *trkB* mRNA probably results in an accumulation of *trkB* receptors in the axonal growth cone, which could make motoneurons more sensitive to local and target-derived neurotrophin. A recent study in genetically engineered mice lacking *trkB* receptors, has indeed demonstrated that *trkB* is required for the survival of some facial motoneurons after axotomy (Alcántara et al., 1997).

The expression of neurotrophins and neurotrophin receptors is altered also after injuries in the central nervous system (see Fig. 2). NGF levels are transiently elevated at CNS injuries (Bakhit et al., 1991; Ishikawa et al., 1991; Lindholm, 1992). NGF synthesis in astrocytes can be induced in vitro by several cytokines and other molecules (Gadient, 1990; Lindholm et al., 1990; Spranger et al., 1990; Pechán et al., 1992, 1993; Zafra et al., 1992; Ladenheim, 1993), and the increase in NGF at a CNS injury may be due to increased levels of macrophage-derived cytokines. BDNF mRNA has been found to increase in neurons in the brain after ischemia and hypoglycemia (Lindvall et al., 1992) but not in glial cells after traumatic injuries (Lindholm et al., 1992). p75 and truncated *trkB* receptors are expressed at high levels by reactive astrocytes as well as by leptomeningeal cells at CNS injuries (Brunello et al., 1990; Frisén et al., 1992, 1993a,c; Risling et al., 1992; Beck et al., 1993).

At this point, only speculations can be made regarding the function of non-neuronal expression of p75 and truncated *trkB* receptors in the injured CNS. It is unlikely that truncated *trkB* receptors can transduce a signal to the cell since they lack the catalytic tyrosine kinase domain. It is also doubtful whether p75 on glial cells can mediate any biological effect of neurotrophins, although evidence has been presented for NGF signaling and internalization via p75 in glial cells (Kahle and Hertel, 1992; Carter et al., 1996). Both p75 and




Fig 2. Schematic drawing illustrating alterations in immunoreactivity and/or mRNA for neurotrophins and neurotrophin receptors after central and peripheral nerve injury. See the text for references.

truncated *trkB* receptors could perhaps serve a function by binding and accumulating neurotrophins at the site of the injury, thus making them available for axonal endbulbs. Neurotrophins bound to the surfaces of non-neuronal cells at a CNS lesion could possibly stimulate axonal growth. However, the immunoglobulin-like domains in the ectodomain of *trk* family receptors can inhibit neurite growth (Tannahill et al., 1995), and the high expression of truncated *trkB* receptors by reactive astrocytes may thus possibly inhibit axon-growth.

In addition to the neurotrophins, many other neurotrophic factors have been suggested to be of importance after neuronal injury. Moreover, several growth factors, best known for their effects on cells outside the nervous system, can stimulate axonal growth or promote neuronal survival and have been implicated in axonal regeneration, although the roles of these factors have not yet been studied as thoroughly as the effects of neurotrophins (see e.g. Raivich and Kreutzberg, 1993). The expression of ciliary neurotrophic factor (CNTF) decreases dramatically in the distal stump of a severed peripheral nerve (Sendtner et al., 1992b), but is expressed in reactive astrocytes after CNS injury (Ip et al., 1993). CNTF lacks a signal sequence and does not seem to be secreted from cells. However, CNTF may possibly be released from damaged cells, and the retrograde transport of this molecule is indeed increased after injury (Curtis et al., 1993). Glial cell-line derived neurotrophic factor (GDNF), a potent neurotrophic factor which is required for the survival of certain neurons during development (Moore et al., 1996; Sánchez et al., 1996), is another example of a neurotrophic factor that is produced at increased levels in injured peripheral nerves (Trupp et al., 1995; Hammarberg et al., 1996).

However, in spite of the wealth of circumstantial evidence implicating several neurotrophic factors in axonal regrowth after injury, there are in fact data suggesting that none of them may be required for regeneration. Axons can grow into acellular peripheral nerve segments, in which all cells have been killed and thus no neurotrophic factors can be synthesized (Ide et al., 1983; Sketelj et al., 1989).

Extracellular matrix

Acellular nerve segments predominantly consist of extracellular matrix tubes, and axon-growth in this environment suggests that extracellular matrix proteins may be of importance for regeneration. Much of the attention in this context has focused on laminin and related molecules, since this glycoprotein synthesized by Schwann cells is one of the strongest neurite outgrowth promoting matrix molecules known to date. Other matrix molecules which may affect neurite growth include for example collagen, fibronectin and tenascin (see Martini, 1994). Several different approaches have been used to study the role of laminin in axon-growth in mammals *in vivo*. Morphological studies have demonstrated that cut

PNS axons regrow in close relation to laminin (Bignami et al., 1984; Liesi, 1985; Salonen et al., 1987; Kuecherer-Ehret et al., 1990). Other studies have shown the effect of blocking the interaction between axons and laminin. These experiments have demonstrated impeded axonal growth in acellular nerve segments pretreated with anti-laminin (or anti-fibronectin antisera) (Wang et al., 1992a,b), as well as weak axon-ingrowth into extracellular matrix-coated tubes in the presence of antibodies to the laminin/collagen receptor $\alpha 1\beta 1$ -integrin (Toyota et al., 1990). Furthermore, the re-innervation of the denervated iris is slower in the presence of antibodies to a laminin-proteoglycan complex (Sandrock and Matthew, 1987b). Taken together, these data suggest that laminin is important in axonal regeneration in the PNS.

In the CNS, astrocytes express laminin during development and reexpress laminin after injuries in the adult. However, laminin is only synthesized locally at injuries and not in degenerating tracts undergoing Wallerian degeneration (Liesi et al., 1984). Axonal sprouts which grow into the scar tissue that forms at the lesion after a spinal cord injury are closely related to laminin expressing astrocytes (Risling et al., 1993; Frisé et al., 1995a). Furthermore, when neuronal cells are cultured on spinal cord cryostat sections, the neurons attach to laminin-rich regions of the sections and orient their neurites after laminin in the tissue (Frisé et al., 1995a). These data suggest that laminin may be involved in lesion-induced sprouting in the injured spinal cord. However, no relation was found between axonal sprouting and laminin in the injured optic nerve (Giftochristos and David, 1988). Possible explanations for this discrepancy may be differences in axonal and/or glial properties in the spinal cord and optic nerve. Differences in laminin-sensitivity in the different neuronal cell types could possibly be of importance; indeed, whereas sensory neurons (which project to the spinal cord) remain laminin-sensitive in adult animals (Unsicker et al., 1985), retinal ganglion cell laminin-receptors decrease after innervation of the target tissue during development, and the neurite-outgrowth promoting effect of laminin is lost (Cohen et al., 1986, 1989). Moreover, different populations of astrocytes in various parts of the CNS may respond differently to injuries, and the ability of astrocytes to support neurite growth *in vitro* depends on which part of the CNS the glial cells were taken from (reviewed by Wilkin et al., 1990; Hatten et al., 1991).

Other matrix molecules which are upregulated after CNS injuries include, among others tenascin, collagen, and fibronectin (Egan and Vijayan, 1991; McKeon et al., 1991; Bartsch et al., 1992; Laywell et al., 1992; Risling et al., 1993) all of which may be of importance for axon growth. The expression and role of cell adhesion molecules belonging to e.g. the immunoglobulin and cadherin families, which could be important for neuron-glia interactions (Neugebauer et al., 1988; Tomaselli et al., 1988), remains to be studied in the injured nervous

system. Several cell adhesion molecules have been implicated in developmental and regenerative axon-glia interactions (Reichardt et al., 1990).

Axon-growth inhibiting molecules

Through the years theories regarding cell extension and migration have mostly focused on stimulating cues in the environment, but more recently growth control by repulsion has gained increasing interest. Schwab and collaborators have provided a large body of evidence for the presence of neurite growth-inhibiting molecules within the brain and spinal cord (reviewed in Schwab et al., 1993). One of the first strong indications for the presence of axon-growth inhibiting activity was the finding that cultured neurons can attach to and send out neurites on slices of peripheral nerves and CNS gray matter, but not on CNS white matter even in the presence of neurotrophic factors (Schwab and Thoenen, 1985; Carbonetto, 1987). Cocultures of neurons and glial cells demonstrated that neurites readily grew over astrocytes but when the growth cones came in contact with oligodendrocytes they not only ceased to grow, but collapsed and were retracted (Schwab and Caroni, 1988). That oligodendrocytes are important inhibitors *in vivo* is supported by studies demonstrating that regeneration in the embryonic spinal cord is enhanced when the onset of myelination is delayed (Keirstead et al., 1992) or oligodendrocyte precursor cells are eliminated by x-irradiation (Savio and Schwab, 1990). Recently, some functional recovery was made possible in spinal cord transected paraplegic rats when axons were directed from the white matter to the gray matter with PNS grafts bridging the lesion (Cheng et al., 1996), thus probably circumventing the inhibitory oligodendrocytes in the white matter.

Two oligodendrocyte-derived molecules that are at least in part responsible for this dramatic effect on axon-growth are myelin-associated proteins designated NI-35 and NI-250 (Caroni and Schwab, 1988). They seem to cause growth cone collapse and retraction by binding to neuronal receptors which, via G-proteins, induce rapid influx of extracellular calcium (Bandtlow et al., 1993; Igarashi et al., 1993). Inhibition of axonal growth *in vivo* by these oligodendrocyte proteins is indicated by the fact that transected cortico-spinal and septo-hippocampal axons show extensive axon-growth in the presence of neutralizing antibodies to NI-35 and NI-250 (Schnell and Schwab, 1990, 1993; Cadelli and Schwab, 1991). Myelin-associated glycoprotein (MAG) is another oligodendrocyte-derived molecule that inhibits axonal growth *in vitro* (McKerracher et al., 1994; Mukhopadhyay et al., 1994), but it may not be a major inhibitor *in vivo* (Bartsch et al., 1995). Several other molecules produced by oligodendrocytes that can inhibit axon-growth have been described, but their significance remains to be established (Schwab et al., 1993).

Myelin-related axonal-growth inhibiting molecules

thus seem to be very important inhibitors of axonal regeneration in the CNS. However, myelin is removed from degenerating tracts with time, but the axons still do not regrow, suggesting that other factors may be important. The finding that the initially strong regenerative response of retinal ganglion cells ceases within a few weeks after injury (Thanos and Vanselow, 1989), led to the suggestion that the slow myelin removal in Wallerian degeneration in the CNS (where myelin is cleared away only after the regenerative attempts by retinal neurons have stopped) is a major reason for the lack of regeneration in the CNS (David et al., 1990). However, neurons fail to attach *in vitro* to sections of tracts undergoing Wallerian degeneration in the spinal cord even long after the injury, at a time when little myelin can be detected (Frisén et al., 1994). Furthermore, lack of axon ingrowth in the adult spinal cord dorsal horn after dorsal root injury, where few oligodendrocytes are found (Carlstedt, 1985), indicates that factors other than myelin-related molecules may counteract CNS regeneration. Indeed, other axon-growth inhibiting factors (e.g. glial hyaluronate-binding protein, GHAP) are expressed by astrocytes in areas of Wallerian degeneration, but not, or at substantially lower levels, in astrocytic scars formed at traumatic injuries (Mansour et al., 1990; Bovolenta et al., 1993).

In the last few years, a plethora of novel axon-growth inhibiting molecules have been identified as a result of searches for molecules involved in axonal guidance during the development of the nervous system. Emerging evidence strongly implicate several of these proteins, for example semaphorins, netrins and ligands for Eph-related tyrosine kinase receptors, in axonal pathfinding (Goodman, 1996), but very little is yet known regarding their possible role in axonal re-generation.

Cooperation with non-nervous system cells

Transection of a peripheral nerve results in the rapid disintegration of the barrier between the blood and the nerve tissue, i.e. the blood-nerve barrier, distal to the injury (Olsson, 1966; Mellick and Cavanagh, 1968). This enables the fast recruitment of mononuclear leukocytes which phagocytize degrading axons and myelin (see Beuche and Fride, 1984) (Fig. 3). The importance of macrophages in the repair after injuries in the nervous system is indicated by, for example, the finding that regeneration of sensory axons is very slow in the mouse strain C57BL/Ola, in which macrophage recruitment to an injured nerve is almost absent (Lunn et al., 1989). Macrophages which invade an injured nerve seem, via interleukin-1, to induce NGF-synthesis by non-neuronal cells (Lindholm et al., 1987; Heumann et al., 1987b), and the weak sensory axon regrowth in C57BL/Ola mice can be supported by exogenous NGF (Brown et al., 1991).

After an injury in the CNS, macrophages rapidly invade the lesion area. However, macrophage infiltration

distal to a CNS injury is slow (Fig. 3), resulting in the delayed clearance of debris in tracts undergoing Wallerian degeneration (Frisén et al., 1994). Macrophages may be detrimental or beneficial following CNS injury. Suppression of macrophage invasion after transient spinal cord ischemia in rabbits improves functional recovery and promotes motoneuron survival (Giulian and Robertson, 1990). A possible role for macrophages could be to phagocytize debris from axotomized neurons undergoing retrograde cell death. Furthermore, it is well known that a large number of synapses on motoneurons disappear after axotomy. In addition to phagocytosis of necrotic neuronal somata, macrophages could therefore be involved in phagocytosis of terminals on moto-neurons (Blinzinger and Kreutzberg, 1968). Morphological indications of macrophage phagocytosis of neuronal elements have been demonstrated on seemingly intact neurites during development (Innocenti et al., 1983) as well as on axons in sciatic nerve-end neuromata (Frisén et al., 1993b). An alternative way in which macrophages may negatively affect neurons unaffected by the initial trauma is by secreting neurotoxic molecules (Théry et al., 1991). In contrast, macrophages can counteract the inhibitory action of optic nerve white matter on neurite growth *in vitro*, and could therefore be important in axonal sprouting in the injured CNS (David et al., 1990). Macrophages are thus implicated in injury reactions in both the PNS and CNS, in part by acting as regulators of some of the other determinants of axonal regeneration mentioned above.

The glial scar - a physical barrier for axonal




Fig. 3. Disruption of the barrier to blood and recruitment of macrophages is different in the PNS and in the CNS. In the PNS, the blood-nerve barrier is disrupted throughout the distal segment of a severed nerve. This enables rapid invasion of macrophages which engulf myelin and axonal degradation products. In the CNS, the disruption of the blood-brain barrier is limited to the lesion area. Macrophages will rapidly enter the CNS at the lesion and clear this area from debris, but macrophage recruitment and removal of debris distal to the lesion is very slow.

The glial scar - a physical barrier for axonal regrowth?

After lesions in the CNS, astrocytes both at the injury and in denervated tracts undergo hypertrophy and increase the synthesis of the intermediate filament proteins glial fibrillary acidic protein, nestin and vimentin (Eddleston and Mucke, 1993; Frisén et al., 1995b). Within a few weeks the astrocytes form a dense network of processes, referred to as the glial scar. The fact that this tissue appears compact histologically led to the suggestion that it may constitute a physical barrier for axon regrowth (reviewed in Reier et al., 1989). An axon-growth inhibitory effect of astrocytes has been supported for example by studies on regeneration of the central branch from primary sensory neurons: after a dorsal root crush in the adult rat, the sensory axons regrow to the dorsal root transitional zone, where most axons then stop growing and form terminals on astrocytes (Carlstedt, 1985). When the axons make contact with astrocytes, the neurofilament synthesis in the neuronal cell body ceases. Astrocytes have therefore been suggested to activate an inherent neuronal physiological stop mechanism (Liuzzi and Lasek, 1987; Liuzzi and Tedeschi, 1992).

Despite the wealth of data suggesting axon-growth inhibition by astrocytes, these glial cells are believed to be important in axonal growth promotion and guidance during embryogenesis. Furthermore, astrocytes in culture form a suitable substrate for cocultured neurons, and neurites readily grow over the astrocytes (reviewed in Hatten et al., 1991). Moreover, already by the turn of the century scientists such as Santiago Ramón y Cajal noticed that after injuries in the CNS, axons sent sprouts into the scar formed at the lesion, although they failed to grow beyond this scar. The axonal sprouts in the scar tissue are closely associated with astrocytes. In many situations scar tissue sprouts seem to be aborted within one month after an injury (Ramón y Cajal, 1928). However, several studies have demonstrated that axonal sprouts in the spinal cord can persist for more than a year in the glial scar formed after an incision (Risling et al., 1983, 1992; Frisén et al., 1993a). After a ventral funiculus incision or after ventral root avulsion and replantation, many motoneuron axons manage to traverse the glial scar and reinnervate the ventral root, and may even reinnervate the target tissue in a functional manner (Cullheim et al., 1989; Carlstedt et al., 1993, 1995).

Ramón y Cajal hypothesized from the fact that axonal sprouts grow into the scar tissue formed at lesions that stimulatory substances for axon growth may be present in the scar (Ramón y Cajal, 1928). Many decades later this hypothesis proved to be true (Nieto-Sampedro et al., 1982). Among the neurotrophic factors that have been described to be expressed at increased levels in glial scars are NGF (Ishikawa et al., 1991; Lindholm et al., 1992), basic fibroblast growth factor (Frautschy et al., 1991; Ishikawa et al., 1991; Logan et al., 1992; Koshinaga et al., 1993), insulin-like growth

factor 1 (IGF-1; García-Estrada et al., 1992), transforming growth factor- β (Lindholm, 1992), and CNTF (Ip et al., 1993). Furthermore, expression of neurite-growth promoting extracellular matrix molecules such as laminin, collagen, fibronectin, and tenascin may be of importance (discussed above). The majority of these axon-growth promoting molecules are synthesized by astrocytes. In addition to these axon growth-promoting molecules, axon growth-inhibiting activity is also found at CNS lesions (Rudge and Silver, 1990; McKeon et al., 1991).

One explanation to the contradictory reports on the role of astrocytes and glial scars on axonal elongation is that astrocytes react differently to different types of injuries. Glial scars are formed both at the site of an injury, such as an incision, and distal to this point where the distal segments of the transected axons degenerate, i.e. Wallerian degeneration. These two types of scars, the type that is formed at the site of a penetrating injury and the type that forms in a tract undergoing Wallerian degeneration, differ in many respects, not least in their capacity to promote axonal growth. The increased synthesis of axon-growth promoting molecules is limited to the type of glial scar that forms at the site of the injury, and does not take place in the scar that forms in the zone undergoing Wallerian degeneration. The differential capacity to promote axon-growth in these two types of glial scars has been demonstrated in *in vitro* experiments where neuronal cells have been cultured on tissue sections from the injured CNS. Neurons can attach and extend neurites on astrocytes in sections from a glial scar formed at a penetrating injury, but fail to attach to sections of a denervated tract undergoing Wallerian degeneration (David et al., 1990; Frisén et al., 1994).

The regulation of the astrocytic phenotype with regard to neurite-growth promotion is poorly understood. Possible factors leading to a change in astrocyte properties after injury may for example be the direct trauma to the cell, a change in the extracellular environment caused by the blood-brain barrier disruption with concomitant influx of blood cells and plasma, or signals from severed neurons or glial cells. *In vitro* studies have identified molecules able to induce synthesis of NGF, BDNF, bFGF, and laminin by astrocytes (Gadient et al., 1990; Lindholm et al., 1990; Spranger et al., 1990; Pechán et al., 1992, 1993; Toru-Delbauffe et al., 1992; Zafra et al., 1992; Baghdassarian et al., 1993; Ladenheim et al., 1993). Interestingly, some data suggest that macrophages may be involved in the regulation of astrocytic properties after injury. Local high levels of TGF- β at the lesion site, probably at least in part derived from macrophages invading the site after CNS injury, may inhibit glial proliferation and induce laminin- and NGF-synthesis by astrocytes (Lindholm et al., 1990, 1992; Toru-Delbauffe et al., 1992; Baghdassarian et al., 1993).

The ingrowth of axons into CNS scar tissue demonstrates that the adult mammalian CNS has a potential to support axon-growth. This might perhaps

involve reactivation of developmental programs used to support and guide growing axons during embryogenesis. An understanding of the molecular basis for the axon growth promotion by reactive glial cells, and the elucidation of the signals which induce this permissive glial phenotype may in the future lead to strategies to manipulate glial features and promote axonal regeneration.

Is the blood-brain barrier a barrier for axonal regeneration?

In the intact brain and spinal cord, as well as in peripheral nerves, the nervous tissue is protected from the outer environment by a barrier to the blood. The barrier between the blood and the CNS, the blood-brain barrier (BBB), resides in the endothelium of blood vessels in the CNS: these endothelial cells do not take up molecules by endocytosis to the same extent as do those outside the CNS and they are interconnected by tight junctions which hinder passive diffusion of molecules through the vessel wall. These features are believed to be induced in the endothelial cells by soluble factors from neighboring astrocytes (Janzer and Raff, 1987; Lobrinus et al., 1992).

The role of the barrier between blood and nervous tissue in the context of nerve regeneration is intriguing. In most, if not all, regenerating systems no barrier is present during axon regrowth. In lower animals such as amphibians, in which CNS regeneration is possible, the BBB is disrupted in the denervated part of an injured tract (Kiernan and Contestabile, 1980). In mammals the blood-nerve barrier is similarly disrupted distal to a peripheral nerve lesion (Olson, 1966; Mellick and Cavanagh, 1968). Moreover, olfactory receptor neurons residing in the nasal mucosa are exchanged throughout life in mammals, and axons from new receptor cells grow to the glomeruli in the olfactory bulb (Graziadei and Monti Graziadei, 1979). The BBB is incomplete in the superficial olfactory bulb where the axon ingrowth to the CNS occurs (Balin et al., 1986). After injuries in the CNS, the BBB is disrupted locally at the site of the injury, and not, as in the PNS, throughout the denervated area (Olsson, 1966; Mellick and Cavanagh, 1968) (Fig. 3). This defect in the BBB is usually repaired within a month (Kiernan, 1985), which correlates with the retraction of lesion-induced sprouts in the CNS (Ramón y Cajal, 1928). However, in the situation of spinal cord injuries, where axonal sprouts may persist for much longer times, the BBB defect is much more persistent (Risling et al., 1989, 1990; Frisén et al., 1993a).

Is the correlation between axon-growth and absence of a barrier to the blood merely a coincidence or can the contact between the blood and nervous tissue be causally related to axonal growth? The absence of a complete barrier could enable molecules as well as cells to pass from the blood to the nervous tissue. In a peripheral nerve the rapid disruption of the blood-nerve barrier makes it possible for macrophages to enter the nerve

distal to an injury (Fig. 3). These blood-derived cells phagocytize myelin and axonal debris, and may secrete potent cytokines and growth factors. Macrophages rapidly enter the lesion area after brain or spinal cord trauma, but fail initially to enter the denervated tract (see above and Fig. 3). Macrophages may also be of importance in the injured CNS not least because of their ability to neutralize inhibitory properties in the mammalian CNS (David et al., 1990). Macrophage-derived cytokines may also be intimately involved in the glial reactions seen at the injury. In addition, molecules that normally circulate in plasma, such as growth factors and cytokines, which may enter the injured spinal cord at a site of failing blood-brain barrier, could be important for axon growth.

Future perspectives

Very much has been learned about axonal regeneration during the past few decades, and the industrious activity in this field is likely to increase our understanding substantially during the coming years. The first step on the road to enhancing axonal regeneration in the CNS must be to promote neuronal survival and axonal growth. Elucidating the mechanisms behind injury-induced nerve cell death and determinants of intrinsic neuronal regenerative capacity will be of fundamental importance. Other pivotal studies for the near future will be further characterization of axon growth-inhibiting molecules in the CNS. This will facilitate the search for neuronal receptors transmitting signals which cause axons to stop growing. Identification of neuronal receptors for CNS axon growth-inhibiting molecules, and the development of antagonist drugs to such receptors may prove instrumental in the treatment of brain and spinal cord injuries. Increased knowledge regarding the regulation of astrocyte features, enabling the targeted induction of an axon growth permissive phenotype after injury may also prove valuable. However, even if axon growth can be induced in the adult CNS, will these axons be able to navigate to their appropriate target? Misdirected growth of axons and innervation of inappropriate targets may very well constitute a worse situation than no axon growth. Axonal pathfinding during embryogenesis is directed by distinct molecular cues in the path of the growing axons and on the target cells. The molecular nature of some of these axon guidance cues have recently been unveiled. Are these guidance cues only expressed during embryogenesis, or are they present also in the adult tissues (or are they reexpressed after injury) so that they could direct regrowing axons to the appropriate target? Some experiments give reason to hope that this is the case (Wizenmann et al., 1993; Bähr and Wizenmann, 1996). The identification of molecules that support or inhibit axonal growth have resulted in a major leap forward in our understanding of axonal regeneration, and it is likely that as more molecular key players in axonal growth are being unveiled that it soon

may be possible to restore severed neural circuits in the injured mammalian CNS.

Acknowledgments. I am indebted to Drs. D. Clarke, M. Jaramillo and I. Silos-Santiago for thoughtful discussions and suggestions on the manuscript. The experiments from the authors lab described here were supported by the Swedish Medical Research Council, Stiftelsen Elsa och Sigurd Goljes minne, Hans och Loo Ostermans fond, Åke Wibergs stiftelse, Jeanssonss stiftelse, Tore Nilssons fond and the Swedish Medical Society.

References

- Aguayo A.J., Dickson R., Trecarten J., Attiwell M., Bray G.M. and Richardson P. (1978). Ensheathment and myelination of regenerating PNS fibres by the transplanted optic nerve glia. *Neurosci. Lett.* 9, 97-104.
- Aguayo A.J., Rasminsky M., Bray G.M., Carbonetto S., McKerracher L., Villegas-Pérez M.P., Vidal-Sanz M. and Carter D.A. (1991). Degenerative and regenerative responses of injured neurons in the central nervous system of adult mammals. *Phil. Trans. R. Soc. Lond.* 331, 337-343.
- Alcántara S., Frisén., del Rio J.A., Soriano E., Barbacid M and Silos-Santiago I. (1997). *trkB* signalling is required for postnatal survival of CNS neurons and protects hippocampal and motor neurons from axotomy-induced cell death. *J. Neurosci.* 17, 3623-3633.
- Aldskogius H. and Svensson M. (1993). Neuronal and glial cell responses to axon injury. *Adv. Struct. Biol.* 2, 191-223.
- Baghdassarian D., Toru-DelBauffe D., Gavaret J.M. and Pierre M. (1993). Effects of transforming growth factor- β 1 on the extracellular matrix and cytoskeleton of cultured astrocytes. *Glia* 7, 193-202.
- Bähr M. and Wizenmann A. (1996). Retinal ganglion cell axons recognize specific guidance cues present in the deafferented adult rat superior colliculus. *J. Neurosci.* 16, 5106-5116.
- Bakhit C., Armanini M., Wong W.L.T., Bennet G.L. and Wrathall J.R. (1991). Increase in nerve growth factor-like immunoreactivity and decrease in choline acetyltransferase following contusive spinal cord injury. *Brain Res.* 554, 264-271.
- Balin B.J., Broadwell R.D., Salzman M. and El-Kalliny M. (1986). Avenues for entry of peripherally administered protein to the central nervous system in mouse, rat, and squirrel monkey. *J. Comp. Neurol.* 251, 260-280.
- Bandtlow C.E., Schmidt M.F., Hassinger T.D., Schwab M.E. and Kater S.B. (1993). Role of intracellular calcium in NI-35-evoked collapse of neuronal growth cones. *Science* 259, 80-83.
- Barbacid M. (1995). Structural and functional properties of the TRK family of neurotrophin receptors. *Ann. NY Acad. Sci.* 766, 442-458.
- Bartsch U., Bartsch S., Dörries U. and Schachner M. (1992). Immunohistochemical localization of tenascin in the developing and lesioned adult mouse optic nerve. *Eur. J. Neurosci.* 4, 338-352.
- Bartsch U., Bandtlow C.E., Schnell L., Bartsch S., Spillman A.A., Rubin B.P., Hillenbrand R., Montag D., Schwab M.E. and Schachner M. (1995). Lack of evidence that myelin-associated glycoprotein is a major inhibitor of axonal regeneration in the CNS. *Neuron* 15, 1375-1381.
- Beck K.D., Lamballe F., Klein R., Barbacid M., Schauwecker P.E., McNeill T.H., Finch C.E., Hefti F. and Day J.R. (1993). Induction of noncatalytic *trkB* neurotrophin receptors during axonal sprouting in

Determinants of axonal regeneration

- the adult hippocampus. *J. Neurosci.* 13, 4001-4014.
- Benfey M. and Aguayo A.J. (1982). Extensive elongation of axons from rat brain into peripheral nerve grafts. *Nature* 296, 150-152.
- Bernstein J.J. and Bernstein M.E. (1971). Axonal regeneration and formation of synapses proximal to the site of lesion following hemisection of the rat spinal cord. *Exp. Neurol.* 30, 336-351.
- Beuche W. and Friede R.L. (1984). The role of non-resident cells in Wallerian degeneration. *J. Neurocytol.* 13, 767-796.
- Bignami A., Chi N.H. and Dahl D. (1984). Laminin in rat sciatic nerve undergoing Wallerian degeneration. *J. Neuropathol. Exp. Neurol.* 43, 94-103.
- Blinzinger K. and Kreutzberg G. W. (1968). Displacement of synaptic terminals from regenerating motoneurons by microglial cells. *Z. Zellforsch. Mikrosk. Anat.* 85, 145-157.
- Bovolenta P., Wandosell F. and Nieto-Sampedro M. (1993). Characterization of a neurite outgrowth inhibitor expressed after CNS injury. *Eur. J. Neurosci.* 5, 454-465.
- Brown M.C., Perry V.H., Lunn E.R., Gordon S. and Heumann R. (1991). Macrophage dependence of peripheral sensory nerve regeneration: possible involvement of nerve growth factor. *Neuron* 6, 359-370.
- Brunello N., Reynolds M., Wrathall J.R. and Mocchetti I. (1990). Increased nerve growth factor receptor mRNA in contused rat spinal cord. *Neurosci. Lett.* 118, 238-240.
- Cadelli D. and Schwab M.E. (1991). Regeneration of lesioned septo-hippocampal acetylcholinesterase-positive axons is improved by antibodies against the myelin-associated neurite growth inhibitors NI-35/250. *Eur. J. Neurosci.* 3, 825-832.
- Carbonetto S., Evans D. and Cochar P. (1987). Nerve fiber growth in culture on tissue substrata from central and peripheral nervous systems. *J. Neurosci.* 7, 610-620.
- Carlstedt T. (1985). Regenerating axons form terminals at astrocytes. *Brain Res.* 347, 188-191.
- Carlstedt T., Grane P., Hallin R.G. and Norén G. (1995). Return of function after spinal cord implantation of avulsed spinal nerve roots. *Lancet* 346, 1323-1325.
- Carlstedt T., Hallin R., Hedström K. and Nilsson-Remahl I. (1993). Functional recovery in primates with brachial plexus injury after spinal cord implantation of avulsed ventral roots. *J. Neurol. Neurosurg. Psych.* 56, 649-654.
- Caroni P. and Schwab M.E. (1988). Two membrane protein fractions from rat central myelin with inhibitory properties for neurite growth and fibroblast spreading. *J. Cell Biol.* 106, 1281-1288.
- Carter B.D., Kaltschmidt C., Kaltschmidt B., Offenhäuser N., Böhm-Mattei R., Baeuerle P.A. and Barde Y.-A. (1996). Selective activation of NF- κ B by nerve growth factor through the neurotrophin receptor p75. *Science* 272, 542-545.
- Cheng H., Cao Y. and Olson L. (1996). Spinal cord repair in adult paraplegic rats: partial restoration of hind limb function. *Science* 273, 510-513.
- Cohen J., Burne J.F., Winter J. and Bartlett P. (1986). Retinal ganglion cells lose response to laminin with maturation. *Nature* 322, 465-467.
- Cohen J., Nurcombe V., Jeffrey P. and Edgar D. (1989). Developmental loss of functional laminin receptors on retinal ganglion cells is regulated by their target tissue, the optic tectum. *Development* 107, 381-387.
- Cullheim S., Carlstedt T., Lindå H., Risling M. and Ulfhake B. (1989). Motoneurons reinnervate skeletal muscle after ventral root implantation into the spinal cord of the cat. *Neuroscience* 29, 725-733.
- Curtis R., Adryan K.M., Zhu Y., Harkness P.J., Lindsay R.M. and DiStefano P.S. (1993). Retrograde axonal transport of ciliary neurotrophic factor is increased by peripheral nerve injury. *Nature* 365, 253-255.
- Davenport R.W., Dou P., Rehder V. and Kater S.B. (1993). A sensory role for neuronal growth cone filopodia. *Nature* 361, 721-724.
- David S. and Aguayo A.J. (1981). Axonal elongation into peripheral nervous system "bridges" after central nervous system injury in adult rats. *Science* 214, 931-933.
- David S., Bouchard C., Tsatas O. and Giftchristos N. (1990). Macrophages can modify the nonpermissive nature of the adult mammalian central nervous system. *Neuron* 5, 463-469.
- Derby A., Engelman V.W., Friedrich G.E., Neises G., Rapp S.R. and Roufa D.G. (1993). Nerve growth factor facilitates regeneration across nerve gaps: morphological and behavioural studies in rat sciatic nerve. *Exp. Neurol.* 119, 176-191.
- Diamond J., Coughlin M., Macintyre L., Holmes M. and Vishieau B. (1987). Evidence that endogenous β nerve growth factor is responsible for the collateral sprouting, but not the regeneration, of nociceptive axons in adult rats. *Proc. Natl. Acad. Sci. USA* 84, 6596-6600.
- Diamond J., Foerster A., Holmes M. and Coughlin M. (1992a). Sensory nerves in adult rats regenerate and restore sensory function to the skin independently of endogenous NGF. *J. Neurosci.* 12, 1467-1476.
- Diamond J., Holmes M. and Coughlin M. (1992b). Endogenous NGF and nerve impulses regulate the collateral sprouting of sensory axons in the skin of the adult rat. *J. Neurosci.* 12, 1454-1466.
- Doster S.K., Lozano A.M., Aguayo A.J. and Willard M.B. (1991). Expression of the growth associated protein GAP-43 in adult rat retinal ganglion cells following axon injury. *Neuron* 6, 635-647.
- Eddleston M. and Mucke L. (1993). Molecular profile of reactive astrocytes - implications for their role in neurologic disease. *Neuroscience* 54, 15-36.
- Egan R.A. and Vijayan V.K. (1991). Fibronectin immunoreactivity in neural trauma. *Brain Res.* 568, 330-334.
- Ernfors P., Rosario C.M., Merlio J.-P., Grant G., Aldskogius H. and Persson H. (1993). Expression of mRNAs for neurotrophin receptors in the dorsal root ganglion and spinal cord during development and following peripheral or central axotomy. *Mol. Brain Res.* 17, 217-226.
- Fawcett J.W. (1992). Intrinsic neuronal determinants of regeneration. *TINS* 15, 5-8.
- Fitzgerald M., Wall P.D., Goedert M. and Emson P.C. (1985). Nerve growth factor counteracts the neurophysiological and neurochemical effects of chronic sciatic nerve section. *Brain Res.* 332, 131-141.
- Frautschy S.A., Walicke P.A. and Baird A. (1991). Localization of basic fibroblast growth factor and its mRNA after CNS injury. *Brain Res.* 553, 291-299.
- Frisén J., Verge V.M.K., Cullheim S., Persson H., Fried K., Middlemas D.S., Hunter T., Hökfelt T. and Risling M. (1992). Increased levels of trkB mRNA and trkB protein-like immunoreactivity in the injured rat and cat spinal cord. *Proc. Natl. Acad. Sci. USA* 89, 11282-11286.
- Frisén J., Fried K., Sjögren A.M. and Risling M. (1993a). Growth of ascending spinal axons in CNS scar tissue. *Int. J. Dev. Neurosci.* 4, 461-475.
- Frisén J., Risling M. and Fried K. (1993b). Distribution and axonal relations of macrophages in a neuroma. *Neuroscience* 55, 1003-1013.
- Frisén J., Verge V.M.K., Fried K., Risling M., Persson H., Trotter J.,

Determinants of axonal regeneration

- Hökfelt T. and Lindholm D. (1993c). Characterization of glial trkB receptors: differential response to injury in the central and peripheral nervous system. *Proc. Natl. Acad. Sci. USA* 90, 4971-4975.
- Frisén J., Haegerstrand A., Fried K., Piehl F., Cullheim S. and Risling M. (1994). Adhesive/repulsive properties in the injured spinal cord: relation to myelin phagocytosis by macrophages. *Exp. Neurol.* 129, 183-193.
- Frisén J., Haegerstrand A., Risling M., Fried K., Elde R., Hökfelt T. and Cullheim S. (1995a). Spinal axons in CNS scar tissue are closely related to laminin-immunoreactive astrocytes. *Neuroscience* 65, 293-304.
- Frisén J., Johansson C.B., Török C., Risling M. and Lendahl U. (1995b). Rapid, widespread, and longlasting induction of nestin contributes to the generation of glial scar tissue after CNS injury. *J. Cell Biol.* 131, 453-464.
- Funakoshi H., Frisén J., Barbany G., Timmusk T., Zachrisson O., Verge V.M.K. and Persson H. (1993). Differential expression of mRNAs for neurotrophins and their receptors following axotomy of the sciatic nerve. *J. Cell Biol.* 123, 455-466.
- Gadient R.A., Cron K.C. and Otten U. (1990). Interleukin-1B and tumor necrosis factor- α synergistically stimulate nerve growth factor (NGF) release from cultured rat astrocytes. *Neurosci. Lett.* 117, 335-340.
- García-Estrada J., García-Segura L.M. and Torres-Aleman I. (1992). Expression of insulin-like growth factor I by astrocytes in response to injury. *Brain Res.* 592, 343-347.
- Giftchristos N. and David S. (1988). Laminin and heparan sulphate proteoglycan in the lesioned adult mammalian central nervous system and their possible relationship to axonal sprouting. *J. Neurocytol.* 17, 385-397.
- Giulian D. and Robertson C. (1990). Inhibition of mononuclear phagocytes reduces injury in the spinal cord. *Ann. Neurol.* 27, 33-42.
- Gloster A. and Diamond J. (1992). Sympathetic nerves in adult rats regenerate normally and restore pilomotor function during an anti-NGF treatment that prevents their collateral sprouting. *J. Comp. Neurol.* 326, 363-374.
- Goodman C. S. (1996). Mechanisms and molecules that control growth cone guidance. *Annu. Rev. Neurosci.* 19, 341-377.
- Graziadei P.P.C. and Monti Graziadei G.A. (1979). Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of the olfactory sensory neurons. *J. Neurocytol.* 8, 1-18.
- Hammarberg H., Piehl F., Cullheim S., Fjell J., Hökfelt T. and Fried K. (1996). GDNF mRNA in Schwann cells and DRG satellite cells after chronic sciatic nerve injury. *Neuroreport* 7, 857-860.
- Hatten M.E., Liem R.K.H., Shelanski M.L. and Mason C. A. (1991). Astroglia in CNS injury. *Glia* 4, 233-243.
- Heumann R., Korsching S., Bandtlow C. and Thoenen H. (1987a). Changes of nerve growth factor synthesis in non-neuronal cells in response to sciatic nerve transection. *J. Cell Biol.* 104, 1623-1631.
- Heumann R., Lindholm D., Bandtlow C., Meyer M., Radeke M.J., Misko T.P., Shooter, E. and Thoenen H. (1987b). Differential regulation of mRNA encoding nerve growth factor and its receptor in rat sciatic nerve during development, degeneration, and regeneration: Role of macrophages. *Proc. Natl. Acad. Sci. USA* 84, 8735-8739.
- Ide C., Tohyama K., Yokota R., Nitoro T. and Onodera S. (1983). Schwann cell basal lamina and nerve regeneration. *Brain Res.* 288, 61-75.
- Igarashi M., Strittmatter S.M., Vartanian T. and Fishman M.C. (1993). Mediation by G proteins of signals that cause collapse of growth cones. *Science* 259, 77-79.
- Innocenti G.M., Clarke S. and Koppel H. (1983). Transitory macrophages in the white matter of the developing visual cortex. II. Development and relations with axonal pathways. *Dev. Brain Res.* 11, 55-66.
- Ip N.Y., Wiegand S.J., Morse J., and Rudge J.S. (1993). Injury-induced regulation of ciliary neurotrophic factor mRNA in the adult brain. *Eur. J. Neurosci.* 5, 25-33.
- Ishikawa R., Nishikori K. and Furukawa S. (1991). Appearance of nerve growth factor and acidic fibroblast growth factor with different time courses in the cavity-lesioned cortex of the rat brain. *Neurosci. Lett.* 127, 70-72.
- Janzer R.C. and Raff M.C. (1987). Astrocytes induce blood-brain barrier properties in endothelial cells. *Nature* 325, 253-257.
- Johnson E.M.J., Taniuchi M. and DiStefano P.S. (1988). Expression and possible function of nerve growth factor receptors on schwann cells. *TINS* 11, 299-304.
- Kahle P. and Hertel C. (1992). Nerve growth factor (NGF) receptor on rat glial cell lines; evidence for NGF internalization via p75NGFR. *J. Biol. Chem.* 267, 13917-13923.
- Keirstead H.S., Hasan S.J., Muir G.D. and Steeves J.D. (1992). Suppression of the onset of myelination extends the permissive period for the functional repair of embryonic spinal cord. *Proc. Natl. Acad. Sci. USA* 89, 11664-11668.
- Kiernan J.A. (1985). Axonal and vascular changes following injury to the rat's optic nerve. *J. Anat.* 141, 139-154.
- Kiernan J.A. and Contestabile A. (1980). Vascular permeability associated with axonal regeneration in the optic system of the goldfish. *Acta Neuropathol.* 51, 39-45.
- Kobayashi N.R., Bedard A.M., Hincke M.T. and Tetzlaff W. (1996). Increased expression of BDNF and trkB mRNA in rat facial motoneurons after axotomy. *Eur. J. Neurosci.* 8, 1018-1029.
- Koliatsos V.E., Clatterbuck R.E., Winslow J.W., Cayouette M.H. and Price D. (1993). Evidence that brain-derived neurotrophic factor is a trophic factor for motor neurons *in vivo*. *Neuron* 10, 359-367.
- Koshinaga M., Sanon H.R. and Whittemore S.R. (1993). Altered acidic and basic fibroblast growth factor expression following spinal cord injury. *Exp. Neurol.* 120, 32-48.
- Kuecherer-Ehret A., Graeber M.B., Edgar H., Thoenen H. and Kreutzberg G.W. (1990). Immunoelectron microscopic localization of laminin in normal and regenerating mouse sciatic nerve. *J. Neurocytol.* 19, 101-109.
- Ladenheim R.G., Lacroix I., Foignant-Chaverot N., Strosberg A.D. and Couraud P.O. (1993). Endothelins stimulate c-fos and nerve growth factor expression in astrocytes and astrocytoma. *J. Neurochem.* 60, 260-266.
- Laywell E.D., Dörries U., Bartsch U., Faissner A., Schachner M. and Steindler D.A. (1992). Enhanced expression of the developmentally regulated extracellular matrix molecule tenascin following adult brain injury. *Proc. Natl. Acad. Sci. USA* 89, 2634-2638.
- Liesi P. (1985). Laminin-immunoreactive glia distinguish regenerative adult CNS systems from non-regenerative ones. *EMBO J.* 4, 2505-2511.
- Liesi P., Kaakkola S., Dahl D. and Vaheri A. (1984). Laminin is induced in astrocytes of adult brain by injury. *EMBO J.* 3, 683-686.
- Lindholm D., Heumann R., Meyer M. and Thoenen H. (1987). Interleukine-1 regulates synthesis of nerve growth factor in non-neuronal cells of rat sciatic nerve. *Nature* 330, 658-659.
- Lindholm D., Hengerer B., Zafra F. and Thoenen H. (1990).

Determinants of axonal regeneration

- Transforming growth factor- β 1 stimulates expression of nerve growth factor in the rat CNS. *Neuroreport* 1, 9-12.
- Lindholm D., Castrén E., Kiefer R., Zafra F., and Thoenen H. (1992). Transforming growth factor- β 1 in the rat brain: increase after injury and inhibition of astrocyte proliferation. *J. Cell Biol.* 117, 395-400.
- Lindvall O., Ernfors P., Bengzon J., Kokaia Z., Smith M.-L., Siesjö B.K. and Persson H. (1992). Differential regulation of mRNAs for nerve growth factor, brain-derived neurotrophic factor, and neurotrophin 3 in the adult rat brain following cerebral ischemia and hypoglycemic coma. *Proc. Natl. Acad. Sci. USA* 89, 648-652.
- Liuzzi F.J. and Lasek R.J. (1987). Astrocytes block axonal regeneration in mammals by activating the physiological stop pathway. *Science* 237, 642-645.
- Liuzzi F.J. and Tedeschi B. (1992). Axo-glial interactions at the dorsal root transitional zone regulate neurofilament synthesis in axotomized sensory neurons. *J. Neurosci.* 12, 4783-4792.
- Lobrinus J.A., Juillerat-Jeanneret L., Darekar P., Schlosshauer B. and Janzer R.C. (1992). Induction of the blood-brain barrier specific HT7 and neurothelin epitopes in endothelial cells of the chick chorio-allantoic vessels by a soluble factor derived from astrocytes. *Dev. Brain Res.* 70, 207-211.
- Logan A., Frautschy S.A., Gonzalez A.-M. and Baird A. (1992). A time course for the focal elevation of synthesis of basic fibroblast growth factor and one of its high-affinity receptors (flg) following a localized cortical brain injury. *J. Neurosci.* 12, 3828-3837.
- Lunn E.R., Perry V.H., Brown M.C., Rosen H. and Gordon S. (1989). Absence of Wallerian degeneration does not hinder regeneration in peripheral nerve. *Eur. J. Neurosci.* 1, 27-33.
- Mansour H., Asher R., Dahl D., Labkovsky B., Perides G. and Bignami A. (1990). Permissive and non-permissive reactive astrocytes: immunofluorescence study with antibodies to the glial hyaluronate-binding protein. *J. Neurosci. Res.* 25, 300-311.
- Martini R. (1994). Expression and functional roles of neural cell surface molecules and extracellular matrix. *J. Neurocytol* 23, 1-28.
- McKeon R.J., Schreiber R.C., Rudge J.S. and Silver J. (1991). Reduction of neurite outgrowth in a model of glial scarring following CNS injury is correlated with the expression of inhibitory molecules on reactive astrocytes. *J. Neurosci.* 11, 3398-3411.
- McKerracher L., David S., Jackson D.L., Kottis V., Dunn R.J. and Braun P.E. (1994). Identification of myelin-associated glycoprotein as a major myelin-derived inhibitor of neurite growth. *Neuron* 13, 805-811.
- Mearow K., Ross G., Kril Y., Gloster A., Holmes M. and Diamond J. (1991). Changes in NGF receptor and GAP43 mRNA associated with collateral sprouting and regeneration of dorsal cutaneous nerves in the rat. *Soc. Neurosci. Abstr.* 17, 1312.
- Mearow K.M., Kril Y. and Diamond J. (1993). Increased NGF mRNA expression in denervated rat skin. *Neuroreport* 4, 351-354.
- Mellick R.S. and Cavanagh J.B. (1968). Changes in blood vessel permeability during degeneration and regeneration in peripheral nerves. *Brain* 91, 141-160.
- Meyer M., Matsuoka I., Wetmore C., Olsson L. and Thoenen H. (1992). Enhanced synthesis of brain-derived neurotrophic factor in the lesioned peripheral nerve: different mechanisms are responsible for the regulation of BDNF and NGF mRNA. *J. Cell Biol.* 119, 45-54.
- Moore M.W., Klein R.D., Farinas I., Sauer H., Armanini M., Philips H., Reichardt L.F., Ryan A.M., Carver-Moore K. and Rosenthal A. (1996). Renal and neuronal abnormalities in mice lacking GDNF. *Nature* 382, 76-79.
- Mukhopadhyay G., Doherty P., Walsh F.S., Crocker P.R. and Filbin M.T. (1994). A novel role for myelin-associated glycoprotein as an inhibitor of axonal regeneration. *Neuron* 13, 757-767.
- Neugebauer K.M., Tomaselli K.J., Lilien J. and Reichardt L.F. (1988). N-cadherin, N-CAM and integrins promote retinal neurite outgrowth on astrocytes in vitro. *J. Cell Biol.* 107, 1177-1187.
- Nieto-Sampedro M., Lewis E.R., Cotman C.W., Manthorpe M., Skaper S.D., Barbin G., Longo F.M. and Varon S. (1982). Brain injury causes a time-dependent increase in neuronotrophic activity at the lesion site. *Science* 217, 860-861.
- Olsson Y. (1966). Studies on vascular permeability in peripheral nerves. *Acta Neuropathol.* 7, 1-15.
- Pechán P.A., Chowdhury K. and Seifert W. (1992). Free radicals induce gene expression of NGF and bFGF in rat astrocyte culture. *Neuroreport* 3, 469-472.
- Pechán P.A., Chowdhury K., Gerder W. and Seifert W. (1993). Glutamate induces the growth factors NGF, bFGF, the receptor GFGR-1 and c-fos mRNA expression in rat astrocyte culture. *Neurosci. Lett.* 153, 111-114.
- Piehl F., Frisén J., Risling M., Hökfelt T. and Cullheim S. (1994). Increased trkB mRNA expression by axotomized motoneurons. *Neuroreport* 5, 697-700.
- Raivich G., Hellweg R. and Kreutzberg G.W. (1991). NGF receptor-mediated reduction in axonal NGF uptake and retrograde transport following sciatic nerve injury and during regeneration. *Neuron* 7, 151-164.
- Raivich G. and Kreutzberg G.W. (1993). Peripheral nerve regeneration: role of growth factors and their receptors. *Int. J. Dev. Neurosci.* 11, 311-324.
- Ramón y Cajal S. (1928). Degeneration and regeneration of the nervous system. Hafner. New York.
- Reichardt L.F., Bossy B., Carbonetto S., De Curtis I., Emmet C., Hall D.E., Ignatius M.J., Lefcort F., Napolitano E., Large T., Neugebauer K.M. and Tomaselli K.J. (1990). Neuronal receptors that regulate axon-growth. *Cold Spring Harbor Symp. Quant. Biol.* 55, 341-350.
- Reier P.J., Eng L.F. and Jakeman L. (1989). Reactive astrocytes and axonal outgrowth in the injured CNS: Is gliosis really an impediment to regeneration. In: *Neural regeneration and transplantation*. Seil F.J. (ed). New York Alan R. Liss, Inc. pp 183-209.
- Rich K.M., Yip H.K., Osborne P.A., Schmidt R.E. and Johnson Jr, E.M. (1984). Role of nerve growth factor in the adult dorsal root ganglia neuron and its response to injury. *J. Comp. Neurol.* 230, 110-118.
- Richardson P.M., Issa V.M.K. and Aguayo A.J. (1984). Regeneration of long spinal axons in the rat. *J. Neurocytol.* 13, 165-182.
- Risling M., Cullheim S. and Hildebrand C. (1983). Reinnervation of the ventral root L7 from ventral horn neurons following intramedullary axotomy in adult cats. *Brain Res.* 280, 15-23.
- Risling M., Fried K., Lindå H., Carlstedt T. and Cullheim S. (1993). Regrowth of motor axons following spinal cord lesions: distribution of laminin and collagen in the CNS scar tissue. *Brain Res. Bull.* 30, 405-414.
- Risling M., Fried K., Lindå H., Cullheim S. and Meier M. (1992). Changes in nerve growth factor receptor-like immunoreactivity in the spinal cord after ventral funiculus lesion in adult cats. *J. Neurocytol.* 21, 79-93.
- Risling M., Lindå H., Cullheim S. and Franson P. (1989). A persistent defect in the blood brain barrier after ventral funiculus lesion in adult cats - implications for CNS regeneration. *Brain Res.* 494, 13-21.
- Risling M., Sjögren A.M., Lindå H., Cullheim S., Carlstedt T. and Thelestam M. (1990). Demonstration of a persistent defect in the blood-brain barrier after ventral root implantation into the spinal cord

Determinants of axonal regeneration

- using immunogold technique and staphylococcal alpha-toxin. In: Pathophysiology of the blood-brain barrier. Johansson B.B., Owman C. and Widner H. (eds). Elsevier. Amsterdam. pp 555-559.
- Rudge J. S. and Silver J. (1990). Inhibition of neurite growth on astroglial scars *in vitro*. *J. Neurosci.* 10, 3594-3603.
- Salonen V., Peltonen J., Rönttö M. and Virtanen I. (1987). Laminin in traumatized peripheral nerve: basement membrane changes during degeneration and regeneration. *J. Neurocytol.* 16, 713-720.
- Sánchez M., Silos-Santiago I., Frisén J., He B., Lira S. and Barbacid M. (1996). Renal agenesis and the absence of enteric neurons in mice lacking GDNF. *Nature* 382, 70-73.
- Sandrock A.W. and Matthew W.D. (1987a). Substrate-bound nerve growth factor promotes neurite growth in peripheral nerve. *Brain Res.* 425, 360-363.
- Sandrock A.W.J. and Matthew W.D. (1987b). An *in vitro* neurite-promoting antigen functions in axonal regeneration *in vivo*. *Science* 237, 1605-1608.
- Savio T. and Schwab M.E. (1990). Lesioned corticospinal tract axons regenerate in myelin-free rat spinal cord. *Proc. Natl. Acad. Sci. USA* 87, 4130-4133.
- Schnell L. and Schwab M.E. (1990). Axonal regeneration in the rat spinal cord produced by an antibody against myelin-associated neurite growth inhibitors. *Nature* 343, 269-272.
- Schnell L. and Schwab M.E. (1993). Sprouting and regeneration of lesioned corticospinal tract fibres in the adult rat spinal cord. *Eur. J. Neurosci.* 5, 1156-1171.
- Schwab M. and Caroni P. (1988). Oligodendrocytes and CNS myelin are nonpermissive substrates for neurite growth and fibroblast spreading *in vitro*. *J. Neurosci.* 8, 2381-2393.
- Schwab M.E., Kapfhammer J.P. and Bandtlow C.E. (1993). Inhibitors of neurite growth. *Annu. Rev. Neurosci.* 16, 565-595.
- Schwab M.E. and Thoenen H. (1985). Dissociated neurons regenerate into sciatic but not optic nerve explants in culture irrespective of neurotrophic factors. *J. Neurosci.* 5, 2415-2423.
- Sendtner M., Holtmann B., Kolbeck R., Thoenen H. and Barde Y.-A. (1992a). Brain-derived neurotrophic factor prevents the death of motoneurons in newborn rats after nerve section. *Nature* 360, 757-759.
- Sendtner M., Stöckli K.A. and Thoenen H. (1992b). Synthesis and localization of ciliary neurotrophic factor in the sciatic nerve of the adult rat after lesion and during regeneration. *J. Cell Biol.* 118, 139-148.
- Sketelj J., Brescjanac M. and Popovic M. (1989). Rapid growth of regenerating axons across the segments of sciatic nerve devoid of Schwann cells. *J. Neurosci. Res.* 24, 153-162.
- Snider W.D. (1994). Functions of the neurotrophins during nervous system development: what the knockouts are teaching us. *Cell* 77, 627-638.
- Spranger M., Lindholm D., Bandtlow C., Heumann R., Gnahn H., Näher-Noé M. and Thoenen H. (1990). Regulation of nerve growth factor (NGF) synthesis in the rat central nervous system: comparison between the effects of interleukin-1 and various growth factors in astrocyte cultures and *in vivo*. *Eur. J. Neurosci.* 2, 69-76.
- Sunderland S. (1978). Nerves and nerve injuries. 2 Ed. Churchill Livingstone. London.
- Taniuchi M., Clark H.B. and Johnson E.M.J. (1986). Induction of nerve growth factor receptor in Schwann cells after axotomy. *Proc. Natl. Acad. Sci. USA* 83, 4094-4098.
- Taniuchi M., Clark H.B., Schweitzer J.B. and Johnson E.M.J. (1988). Expression of nerve growth factor receptors by Schwann cells of axotomized peripheral nerves: ultrastructural location, suppression by axonal contact, and binding properties. *J. Neurosci.* 8, 664-681.
- Tannahill L., Klein R. and Schachner M. (1995). The neurotrophin receptors TrkA and TrkB are inhibitory for neurite growth. *Eur. J. Neurosci.* 7, 1424-1428.
- Thanos S. and Vanselow J. (1989). Adult retinal ganglion cells retain the ability to regenerate their axons up to several weeks after axotomy. *J. Neurosci. Res.* 22, 144-149.
- Théry C., Chamak B. and Mallat M. (1991). Cytotoxic effect of brain macrophages on developing neurons. *Eur. J. Neurosci.* 3, 1155-1164.
- Tomaselli K.J., Neugebauer K.M., Bixby J.L., Lilien J. and Reichardt L.F. (1988). N-cadherin and integrins: two receptor systems that mediate neuronal process outgrowth on astrocyte surfaces. *Neuron* 1, 33-43.
- Toru-Delbauffe D., Baghdassarian D., Both D., Bernard R., Rouget P. and Pierre M. (1992). Effects of TGF- β 1 on the proliferation and differentiation of an immortalized astrocyte cell line: Relationship with extracellular matrix. *Exp. Cell Res.* 202, 316-325.
- Toyota B., Carbonetto S. and David S. (1990). A dual laminin/collagen receptor acts in peripheral nerve regeneration. *Proc. Natl. Acad. Sci. USA* 87, 1319-1322.
- Trupp M., Rydén M., Jörnvall H., Funakoshi H., Timmusk T., Arenas E. and Ibáñez C.F. (1995). Peripheral expression and biological activities of GDNF, a new neurotrophic factor for avian and mammalian peripheral neurons. *J. Cell Biol.* 130, 137-148.
- Unsicker K., Skaper S.D., Davis G.E., Manthorpe M. and Varon S. (1985). Comparison of the effects of laminin and the polyornithine-binding neurite promoting factor from RN22 Schwannoma cells on neurite regeneration from cultured newborn and adult rat dorsal root ganglion neurons. *Dev. Brain Res.* 17, 304-308.
- Verge V.M.K., Merlio J.P., Grondin J., Erfors P., Persson H., Riopelle R.J., Hökfelt T. and Richardson P.M. (1992). Colocalization of NGF binding sites, trk mRNA, and low-affinity NGF receptor mRNA in primary sensory neurons: responses to injury and infusion of NGF. *J. Neurosci.* 12, 4011-4022.
- Vidal-Sanz M., Bray G.M., Villegas-Perez M.P., Thanos S. and Aguayo A.J. (1987). Axonal regeneration and synapse formation in the superior colliculus by retinal ganglion cells in the adult rat. *J. Neurosci.* 7, 2894-2909.
- Wang G.-Y., Hirai K.-I. and Shimada H. (1992a). The role of laminin, a component of Schwann cell basal lamina, in rat sciatic nerve regeneration within antiserum-treated nerve grafts. *Brain Res.* 570, 116-125.
- Wang G.-Y., Hirai K.-I., Shimada H., Taji S. and Zhong S.-Z. (1992b). Behaviour of axons, Schwann cells and perineural cells in nerve regeneration within transplanted nerve grafts: effects of anti-laminin and anti-fibronectin antisera. *Brain Res.* 583, 216-226.
- Wilkin G.P., Marriott D.R. and Cholewinski A.J. (1990). Astrocyte heterogeneity. *TINS* 13, 43-46.
- Wizenmann A., Thies E., Klostermann S., Bonhoeffer F. and Bähr M. (1993). Appearance of target-specific guidance information for regenerating axons after CNS lesions. *Neuron* 11, 975-983.
- Yan Q., Elliott J. and Snider W.D. (1992). Brain-derived neurotrophic factor rescues spinal motor neurons from axotomy-induced cell death. *Nature* 360, 753-755.
- Zafra F., Lindholm D., Castrén E., Hartikka J. and Thoenen H. (1992). Regulation of brain-derived neurotrophic factor and nerve growth factor mRNA in primary cultures of hippocampal neurons and astrocytes. *J. Neurosci.* 12, 4793.