

Presence and distribution of 5HT-, VIP-, NPY-, and SP-immunoreactive structures in adult mouse lung

C. Verástegui, J. Fernández-Vivero, A. Prada, F. Rodríguez, A. Romero,
M. González-Moreno and J.M. de Castro

Department of Morphological Sciences, Faculty of Medicine, University of Cádiz, Spain

Summary. A large number of biologically active substances have been identified and characterised in the respiratory tract of several mammals. These substances (amines and peptides) exert important regulatory influences on respiratory functions, and they act as neurotransmitters/neuromodulators, both being released from nerve terminals as neuroendocrine cells. However, these substances can also have other effects which suggest a paracrine action. Thus, to understand the role of amines and peptides in the lung, it is important to explore their localisation in different species. By using immunocytochemical staining methods we have studied the morphology and distribution of serotonin-, Substance P-, neuropeptide Y- and VIP-like immunoreactivity in the adult mouse lung. Moreover a pretreatment with colchicine, pargyline and 5-hydroxytryptophan as staining enlargement method was made. A widespread distribution of isolated endocrine cells and neuroepithelial bodies containing 5HT-like immunoreactivity was recorded within the lung. NPY-like immunoreactive nerve fibres were localised in the airway smooth muscle and surrounding the blood vessels. VIP-like immunoreactivity was revealed in single cells as well as in some nerve fibres and ganglia around the blood vessels and in the bronchial smooth muscle. SP-like IR was observed in nerve fibres located in the smooth muscle of the airways, surrounding bronchi and bronchioli but not next to the intrapulmonary blood vessels. Their localisation both in cells and nerve fibres of the respiratory system suggests that they play a role in the regulatory function of the mouse respiratory tract, exerting their influence by endocrine, paracrine, neurosecretory pathways or a combination of all of these.

Key words: Serotonin, Substance P, Neuropeptide Y, Vasoactive intestinal peptide, Lung

Introduction

The lung is supplied by the motor nerves comprising the sympathetic and parasympathetic autonomic nervous system and by sensory nerves that originate from sensory ganglia of the vagus nerve or from dorsal root ganglia (Richardson, 1979; Doidge and Satchell, 1982; Laitinen et al., 1985; Lundberg et al., 1988). It is now recognised that in addition to classic adrenergic and cholinergic innervation in airways, there is a "third nervous system" that is neither adrenergic nor cholinergic (NANC) (Campbell, 1971; Lundberg et al., 1988).

The function of the bronchial endocrine cells is to detect changes in the oxygen level of the airways. These cells respond to hypoxic conditions by releasing secretory substances. Some of these cells are connected with nerve fibres, whereas others are free of them. All these substances could act either via endocrine, paracrine or humoral pathways (Lauweryns and Peuskens, 1972; Lauweryns and Cokelaere, 1973a,b; Cutz et al., 1985a).

Chemical and immunocytochemical studies confirmed the simultaneous occurrence of neuropeptides and amines in neuroendocrine (NE) cells and led to the concept that the coexistence of different bioactive substances in these cells is the rule rather than the exception (Wharton et al., 1978; Cutz et al., 1981; Keith and Ekman, 1988).

In the respiratory tract, several biologically active substances (amines and peptides) have been identified in human and the other mammals. These substances are in relationships with adrenergic (Richardson, 1979; Doidge and Satchell, 1982; Laitinen et al., 1985), cholinergic (El-Bermani et al., 1982; Amenta et al., 1983), or sensory neural structures (Lauweryns and Van Lommel, 1983; Lundberg et al., 1988). Those neural structures could be both nerve fibres, isolated NE cells and neuroepithelial bodies (NEBs). These NEBs are cellular clusters described by Lauweryns and Peuskens in 1972 with a marked axonic confluence in their cellular basal pole. The storage of serotonin and peptide substances in epithelial NE cells may explain some mechanisms of

*Immunoreactive structures in mouse lung***Table 1.** Immunocytochemical localisation of biologically active substances described in the lungs of various species of adult mammals. The distribution and intensity of immunostaining for these substances varies between species with some showing relatively high levels of expression, and others intermediate or weak immunostaining.

	BB	NPY	CAL	SP	LEU-E	GAL	CGRP	VIP	CCK	5HT	REFERENCES
Human	C	F	C	F	F, C	F	F	F	C	C	Wharton et al., 1978; Polak and Bloom, 1982; Lundberg et al., 1984; Sheppard et al., 1984; Lauweryns and Van Rast, 1987; Cutz et al., 1989; Dey and Zhu, 1993; Wang and Cutz, 1993
Monkey					C, F				C	C	Polak and Bloom, 1982; Lauweryns and Van Rast, 1987; Wang and Cutz, 1993
Pig		F		F	F, C	F	F	F	C	C	Wharton et al., 1979; Polak and Bloom, 1982; Cheung et al., 1985; Lauweryns and Van Rast, 1987; Martling et al., 1990; Wang and Cutz, 1993
Dog						F		F	C	C	Dey et al., 1981a; Polak and Bloom, 1982; Cheung et al., 1985; Wang and Cutz, 1993
Cat		F		F		F, G		F			Dey et al., 1981a; Polak and Bloom, 1982; Said, 1982; Lundberg et al., 1984; Sheppard et al., 1984; Uddman et al., 1984; Dey and Zhu, 1993
Guinea pig	F	F		F	F	F	F	F		C	Wharton et al., 1979; Polak and Bloom, 1982; Lundberg et al., 1984; Sheppard et al., 1984; Uddman et al., 1984; Cheung et al., 1985; Shimosegawa et al., 1989
Rabbit					F			F	C	C	Lauweryns et al., 1973; Said, 1982; Shimosegawa et al., 1989; Wang and Cutz, 1993
Hamster							C		C	C	Polak and Bloom, 1982; Wang and Cutz, 1993
Rat		F	C	F	F	F	F, C	F	C		Polak and Bloom, 1982; Said, 1982; Lundberg et al., 1984; Sheppard et al., 1984; Uddman et al., 1994; Cheung et al., 1985; Cadieux et al., 1986; Shimosegawa et al., 1989; Wang and Cutz, 1993
Mouse		F		F							Sheppard et al., 1984; Uddman et al., 1984

BB: bombesin; NPY: neuropeptide Y; CAL: calcitonin; SP: substance P; LEU-E: leu-enkephalin; GAL: galanin; CGRP: calcitonin gene-related peptide; VIP: vasoactive intestinal peptide; CCK: cholecystokinin; 5HT: serotonin; F: fibres; C: neuroendocrine cells; G: ganglion cells.

local or paracrine regulation mediated by them (e.g. mucous secretion, smooth muscle tone, vasomotor function). In a wide sense, these amines and peptides may act as the integration of the pulmonary activity, and may play a role as neuroendocrine regulators and neurotransmitters (Wasano and Yamamoto, 1978; Wharton et al., 1978; Cutz et al., 1981; Uddman et al., 1984; Dayer et al., 1985).

For the NEBs, a paracrine function has been suggested, but they also serve as functional neuro-receptor organs based on their strategic localisation (Lauweryns and Peuskens, 1972; Lauweryns and Cokelaere, 1973a). However, the exact function of the NEBs remains unknown. They may respond to numerous stimuli as baroreceptors or chemoreceptors. This possible role has been studied in various animal species (Lauweryns and Cokelaere, 1973a,b; Lauweryns and Goddeeris, 1975; Lauweryns et al., 1977; Wasano and Yamamoto, 1981; Keith and Will, 1982).

In the lung, the localisation and concentration of several neurotransmitters-neuromodulators are different. According to the mammal species and also in the same species these concentrations and localisations are age-dependent. Table 1 provides data on the immunocytochemical localisation of biologically active substances which have been described in the lung of various adult mammals. This interspecies variability suggests that functions and roles of the neurotransmitters-neuromodulators probably differ between the different species.

This points out the importance to determine the presence, localisation and distribution of these substances at pulmonary level in each animal. For this reason, we have chosen as experimental animal the adult mouse. Although the mouse has widely been used for studying the control of the intrinsic neuroendocrine system in many organs, little is known about the distribution at the pulmonary level of the substances acting in this control system.

The existence of serotonin in the mammalian respiratory tract has been well documented (Lauweryns et al., 1973, 1982; Cutz et al., 1975; Eaton and Fedde, 1977; Dayer et al., 1985; Bock de et al., 1986).

In addition to serotonin, VIP and substance (SP) are the two peptides so far demonstrated by immunocytochemistry in the mammalian pulmonary nervous system (Uddman et al., 1978; Wharton et al., 1979; Dey et al., 1981a; Ghatei et al., 1982). VIP-immunoreactivity (-IR), though present in several species, is found more abundantly in the cat (Dey et al., 1981a; Said, 1982), and it is the most abundant peptide isolated from human lung (Polak and Bloom, 1986). There is accumulating evidence that VIP may be the neurotransmitter in NANC nerves, but its role cannot be confirmed until specific blockers are developed.

SP was initially detected in nerves of the respiratory tract of the guinea-pig (Nilsson et al., 1977; Terenghi et al., 1983; Saria et al., 1987; Shimosegawa et al., 1989), and this peptide has been to also exist in the respiratory

Immunoreactive structures in mouse lung

tract of humans (Lundberg et al., 1984).

Neuropeptide Y (NPY) has been found in the lungs of several mammals, including human (Sheppard et al., 1984). This neurotransmitter has been demonstrated in sympathetic adrenergic nerves, mostly in relation to blood vessels (Lundberg et al., 1982).

It is now clear that the neuropeptides have a wide influence on the pathophysiological aspects of the lung. SP is a potent endogenous bronchoconstrictor, whereas VIP is a potent endogenous bronchodilator (Said, 1982; Lilly et al., 1993), and NPY may act as a vasoconstrictor. There is abundant evidence that these neuropeptides are released in the lung in a variable conditions and that they have the capacity to modulate the bronchoactivity of the same stimuli that release them. So, VIP seems to be a good candidate as neurotransmitter in airway NANC nerves and the sensory neuropeptide SP may make an important contribution to the physiopathological features of asthma. The recent development of the potential and selective tachykinin receptor antagonists opens new possibilities for therapy of airway diseases. However, the lack of specific and potent antagonists against most of the neuropeptides makes it difficult to define their physiological role in the control of the respiratory tract.

In this report we use several drugs to enlarge the immunostaining. In some samples we used colchicine to inhibit the fast axonal flow at neural cells which causes an unphysiological accumulation of many substances including transmitters (Hökfelt et al., 1978). Moreover, the animal pretreatment with the precursor 5-hydroxytryptophan and with pargyline, the inhibitor of monoamine-oxidase, increased the amount and the intensity at the 5HT-immunoreactive cells.

This work contributes to establish the morphological basis of the mouse pulmonary innervation for the selected neuroregulators, and the point of the scope of light microscopic studies is to establish the coexistence of these substances in the same nerve fibres or NE cells by comparing serial sections or by double-labelling techniques.

Materials and methods

Animals and tissue treatment

Twenty adult male mice (*Mus musculus*, Swiss OF-1) were studied in our experiments.

Ten animals received a total dose of 10 mg/kg (i.p.) of colchicine, 24 h before being killed. The other ten did not receive this treatment. In both groups, two animals were used for immunohistochemical detection of 5HT and these animals received a total dose of 100 mg/kg (i.p.) of 5-hydroxytryptophan and pargyline (100 mg/kg), respectively 60 and 90 min before sacrifice. Animals were anaesthetized with ether inhalation. The lungs were fixed by perfusion through the right ventricle with a fixative solution (4% phosphate-buffered paraformaldehyde, pH 7.4). The samples were then postfixed by immersion in the same fixative for 2h at room

temperature. The fixed tissues were then washed for two hours in 0.05 M TRIS-HCl buffer, pH 7.6, with 0.25% Triton X-100 (Sigma, St Louis, MO).

Immunohistochemistry

After rinsing, serial sections (50-100 µm) were obtained by cryostat sectioning and processed by using the streptavidin-biotin immunoperoxidase method (Bonnard et al., 1984). After treatment with peroxidase blocking reagent and the nonspecific blocking reagent (pig normal serum), sections were exposed to a rabbit anti-5HT, anti-VIP, anti-NPY and anti-SP polyclonal commercial antisera (Cambridge Research Biochemicals, U.K.) diluted 1:1000, 1:2000, 1:500 and 1:1000 respectively in a moisture chamber overnight at 4 °C. The second step was developed by using the avidin-biotin-peroxidase complex (Sigma) diluted 1:50 for 20 min at room temperature. The sections were revealed by rising in a 0.05% diaminobenzidine solution in presence of 0.01% H₂O₂ in the washing buffer under microscopic control. The negative control procedures were carried out by omitting one or more steps of the method and by using sections incubated with inactivated antisera with added excess antigen.

Results

We noted in all animals which received an intraperitoneal dose of colchicine 24 h before killing, the amount and the intensity of the immunoreactive structures was increased.

Serotonin

A widespread distribution of isolated endocrine cells and neuroepithelial bodies with 5HT-like IR was recorded in the lung. The immunoreactive isolated NE cells were located in the epithelium of large bronchi and bronchioles, next to the basal lamina, but they did not establish any contact with the airway lumen (Fig. 1). Their shape was pyramidal and they had a voluminous nucleus. Furthermore, we identified serotonin immunoreactive NEBs located preferentially in the bronchial and bronchiolar bifurcations (Fig. 2). The NEBs were composed of a variable number of serotonergic cells (3 to 50) that were in close contact with the basal lamina, reaching the airway lumen, and which nuclei were round, pale and with a basal disposition.

Our results showed that in the animals which received the pretreatment with the precursor 5-hydroxytryptophan and with the inhibitor of monoamine oxidase pargyline, the amount and the intensity of the 5HT-immunoreactive cell level was increased.

Neuropeptide Y

A rich supply of fine, varicose nerve fibres for this

Immunoreactive structures in mouse lung

mediator were identified in the lung of the adult mouse. These fibres were preferentially located in the smooth muscle of the airways, in the subepithelial layer and

around bronchi and bronchioles (Fig. 3). Furthermore, we observed, the presence of NPY-immunoreactive nerves at the vascular level. Those immunoreactive




Fig. 1. An isolated 5HT-immunoreactive cell displaying a pyramidal shape at bronchial epithelium level in contact with the basal lamina (arrowhead). x 1,000




Fig. 2. Adult mouse bronchial bifurcation epithelium revealing 5HT-IR in a NEB which is extending from the basement membrane to the bronchial airway lumen (arrowheads). E: Epithelium; B: Bronchus. x 1,000

Immunoreactive structures in mouse lung

fibres of large calibre had a sinuous tract and penetrated through the vascular wall, reaching the endothelium (Fig. 4). Some of them ended in the

vascular lumen forming a bulk that was constituted by the reunion of different converging fibres at this level.




Fig.3. NPY immunoreactive nerve fibres beneath the smooth muscle of the mouse lung. x 1,000




Fig.4. A network of NPY immunoreactive nerve fibres surrounds a blood vessel. x 1,000

Substance P

Nerve fibres displaying SP-IR were observed distributed within bronchi and bronchioli walls. The

immunoreactive fibres showed different morphologies (e.g. varicose, filiform). Different calibre and sometimes multiple branching was also observed. In some nerves, SP-immunoreactive fibres coexisted with non-immuno-




Fig. 5. SP immunoreactive nerve fibres with a peribronchial trajectory (arrow). A nerve fibre can be seen at subepithelial level (arrowhead). x 1,000




Fig. 6. SP immunoreactive nerve fibres with different calibers which give branchlets penetrating the bronchial epithelium (arrowhead). x 1,000




Fig. 7. VIP immunoreactive cells at bronchial epithelium in close contact with the basal lamina, where they are arranged between the cells of the epithelium. AL: airway lumen. x 1,000

reactive fibres (Fig. 5). The nervous fibres possessed a peribronchial distribution giving branches which penetrated the bronchial epithelium into the epithelial cells (Fig. 6). We also identified immunoreactive fibres with a long course next to the major bronchi, and these emitted branches at the bifurcation levels. These branches followed the course of the bronchi and bronchioles. No immunoreactivity was observed next to the intrapulmonary blood vessels.

Vasoactive intestinal peptide

We observed VIP-IR in the adult mouse lung both in nerve fibres and in ganglion cells. The IR could also be observed in NE cells of the bronchial epithelium. These cells were mainly located at the bronchial bifurcation and they were in close contact with the airway lumen and with the basal lamina of the epithelium. The cellular morphology was pyramidal with a basal pole bigger than the apical one. They had a large nucleus and a granular cytoplasmic staining. They were arranged in the cells of the epithelium (Fig. 7). Furthermore, it was possible to identify some immunoreactive nervous fibres, basically in the vascular walls and with a perivascular disposition (Fig. 8). The VIP-IR was also observed in the intrinsic ganglia located under the bronchial epithelium and next to blood vessels. They were formed by a variable number of cells, immunoreactive and non-immunoreactive cells coexisting. We also observed VIP-immunoreactive nerve fibres, some of which contact with the ganglion cells in the vicinity of these structures




Fig. 8. VIP immunoreactive nerve fibres at vascular level. x 1,000

(Fig. 9).

Discussion

This work shows that immunoreactive SP, NPY and VIP are widely distributed in nerve fibres surrounding bronchi, bronchioli and blood vessels and immunoreactive 5HT and VIP are presented in NE cells and NEBs at the level of bronchial bifurcation in the adult mouse lung. In agreement with our results, some drugs, such as colchicine, have been used extensively to enhance the detectability of monoamines and peptides in the immunocytochemical techniques.

The understanding of the action of these neurotransmitters-neuromodulators is difficult. Until a few years ago, most of the known peptide hormones were thought to be produced exclusively by endocrine glands. The peptides of the lung are diffusely distributed, and it is likely that the principal tissue upon which these hormones act is the lung itself. Serotonin has been reported to have a contractile effect on smooth muscles of the digestive tract in rats, guinea pigs and rabbits (Kedzierska, 1970). In the breathing lung, it has been proposed that the role of serotonin is to control the mucus production or ciliary function (Dey et al., 1981b). Our results show that the distribution of 5HT NE cells and NEBs at the bronchial epithelium are in agreement with previous data in other mammals (Lauweryns et al., 1973, 1974, 1982; Eaton and Fedde, 1977; Sonstegard et al., 1982). Even though the distribution and amount of 5-

HT cells in the respiratory epithelium in fetus and newborns was bigger than adults (Eaton and Fedde, 1977; Sonstegard et al., 1982; Redick and Hung, 1984; Cutz et al., 1985b; Bock de et al., 1986), the intensity of the immunocytochemical staining and the amount of the immunoreactive cells can be increased in fetal animals following injection of 5-L-hidroxytryptophane, the amine precursor, to the mother during pregnancy (Hage, 1974; Lauweryns et al., 1982), or directly to the specimen for study (Ericson et al., 1972; Cutz et al., 1974, 1975; Lauweryns et al., 1977; Marchesky and Kleinerman, 1983; Carabba et al., 1985).

On the other hand, a rich supply of nerve fibres containing VIP were previously seen close to blood vessels, glands and nonvascular smooth muscle (Luts et al., 1993). However, an interesting finding was the presence of VIP-IR in NE cells of the bronchial epithelium. The placement of endocrine cells throughout the airways suggests that they are involved in the maintenance of important lung functions. The presence of VIP-IR in some local ganglia located under the bronchial epithelium and next to blood vessels suggests that VIP nerves come from these local parasympathetic ganglia.

Substance P-IR was found within nerve fibres, in the airway muscle, around blood vessels and beneath the bronchial epithelium (Luts et al., 1993); this IR being most abundant in the guinea pig (Terenghi et al., 1983; Saria et al., 1987; Shimosegawa et al., 1989). The distribution of the SP-IR fibres in the mouse lung was




Fig. 9. VIP intrapulmonar ganglia immunoreactive (arrows) located under the bronchial epithelium, and next to blood vessels (asterisk). Original magnification x 1,000

Immunoreactive structures in mouse lung

different from described in other species. Thus, the SP innervation is dense in the guinea pig and rat tracheo-bronchial smooth muscle (Lundberg et al., 1988) but only a few SP immunoreactive fibres were found at peripheral airways level. This is in disagreement with our results since we found a rich SP immunoreactive supply at bronchial and bronchiolar epithelium.

NPY has been found in several mammalian lungs, and it is primarily localised in blood vessels (Sheppard et al., 1984). The distribution of NPY is similar to the sympathetic nerves, and it is probably a co-transmitter of norepinephrine and a potent constrictor of vascular smooth muscle (Lundberg et al., 1982). NPY may also play an important role in regulating pulmonary and bronchial vessels, but it probably has less important influence on the airway tone. In conclusion, the dense innervation by NPY-containing fibres (smooth muscle and vascular innervation) suggests that the sympathetic control of airways may not only involve noradrenaline as transmitter.

A comparison with the known localisation of these neuroregulators in the lung of various species revealed that the distribution of NPY-, SP-, VIP-, and 5HT-IR does not correspond to that studied to date in mouse lung. Only NPY and SP have been reported in mouse lung (Sheppard et al., 1984; Uddman et al., 1984). However, to the best of our knowledge, this is the first time that VIP and 5HT have been immunocytochemically demonstrated in the mouse lung in NE cells (VIP and 5HT), NEBs (5HT), nerve fibres (VIP) and ganglia (VIP).

References

- Amenta F., Cavalotti C., Ferrante F. and Tonelli F. (1983). Cholinergic innervation of the human pulmonary circulation. *Acta Anat.* 117, 58-64.
- Bock de V., Yoshizaki K. and Solomon S. (1986). Serotonin content of rabbit lung and small intestine during perinatal development. *Life Sci.* 38, 431-435.
- Bonnard C., Papermaster D.S. and Kraehenbuhl J.P. (1984). The extravidin-biotin bridge technique: Application in light and electron microscope immunocytochemistry. In: Immunolabeling for electron microscopy. J.M. Polak and I.M. Vamdeell (eds). Elsevier. Oxford. pp 95-112.
- Cadieux A., Springall D.R., Mulderry P.K., Rodrigo J., Ghatei M.A., Terenghi G., Bloom S.R. and Polak J.M. (1986). Occurrence, distribution and ontogeny of CGRP immunoreactivity in the rat lower respiratory tract: Effect of capsaicin treatment and surgical denervations. *Neuroscience* 19, 605-627.
- Campbell G. (1971). Autonomic innervation of the lung musculature of a toad (*Bufo marinus*). *Comp. Gen. Pharmacol.* 2, 281-286.
- Carabba V.H., Sorokin S.P. and Hoyt R.F. Jr. (1985). Development of Neuroepithelial bodies in intact and cultured lung of fetal rats. *Am. J. Anat.* 173, 1-27.
- Cheung A., Polak J.M., Bauer F.E., Cadieux A., Christofides N.D., Springall D.R. and Bloom S.R. (1985). Distribution of galanin immunoreactivity in the respiratory tract of pig, guinea pig, rat and dog. *Torax* 40, 889-896.
- Cutz E., Chan W., Wong V. and Conen P.E. (1974). Endocrine cells in rat fetal lungs. Ultrastructural and histochemical study. *Lab. Invest.* 30, 358-464.
- Cutz E., Chan W., Wong V. and Conen P.E. (1975). Ultrastructure and fluorescence biochemistry of endocrine (APUD-type) cells in tracheal mucosa of human and various animals species. *Cell Tissue Res.* 158, 425-437.
- Cutz E., Chan W. and Track N.S. (1981). Bombesin, calcitonin and leu-enkephalin immunoreactivity in endocrine cells of human lung. *Experientia* 37, 765-767.
- Cutz E., Gillan J.E. and Bryan A.C. (1985a). Neuroendocrine cells in the developing human lung: Morphologic and functional considerations. *Pediatr. Pulm.* 1(Suppl.), S21-S29.
- Cutz E., Yeger H., Wong V., Bienkowski E. and Chan W. (1985b). In vitro characteristics of pulmonary neuroendocrine cells isolated from rabbit fetal lung. I. Effects of culture media and nerve growth factor. *Lab. Invest.* 52, 672-683.
- Dayer A.M., De Mey J. and Will J.A. (1985). Localization of somatostatin-, bombesin and serotonin-like immunoreactivity in the lung of the fetal Rhesus monkey. *Cell Tissue Res.* 239, 621-625.
- Dey R.D. and Zhu W. (1993). Origin of galanin in nerves of cat airways and colocalization with vasoactive intestinal peptide. *Cell Tissue Res.* 273, 193-200.
- Dey R.D., Shannon W.A. Jr. and Said S.I. (1981a). Localization of VIP-immunoreactive nerves in airways and pulmonary vessels of dogs, cats and human subjects. *Cell Tissue Res.* 220, 231-238.
- Dey R.D., Echt R. and Dinerstein R. (1981b). Morphology, histochemistry and distribution of serotonin-containing cells in tracheal epithelium of adult rabbit. *Anat. Rec.* 199, 23-31.
- Doidge J.M. and Satchell D.G. (1982). Adrenergic and nonadrenergic inhibitory nerves in mammalian airways. *J. Auton. Nerv. System*, 5, 83-99.
- Eaton J.A. Jr. and Fedde M.R. (1977). Identification of two populations of biogenic amine containing cells in the mouse lung. *Cell Tissue Res.* 176, 243-249.
- El-Bermani A.W., Blomquist E.I. and Montvilo J.A. (1982). Distribution of pulmonary cholinergic nerves in the rabbit. *Thorax* 37, 703-710.
- Ericson L.E.R., Hakanson B. and Larson C.H. (1972). Fluorescence an electron microscopic of the amine storing enterocromaffin-like cells in the tracheal epithelium of mouse. *Z. Zellforsch.* 124, 532-545.
- Ghatei M.A., Sheppard M.N., O'Shaughnessy D.J., Adrian T.E., McGregor G.P., Polak J.M. and Bloom S.R. (1982). Regulatory peptides in the mammalian respiratory tract. *Endocrinology* 111, 1248-1254.
- Hage E. (1974). Histochemistry and fine structure of endocrine cells in fetal lungs of the rabbit, mouse and guinea pig. *Cell Tissue Res.* 149, 513-524.
- Hökfelt T., Ljungdahl A., Steinbusch H., Verhafstad A., Nilsson G., Brodi E., Pernow B. and Goldstein M. (1978). Immunohistochemical evidence of substance P-like immunoreactivity in some 5-hydroxytryptamine-containing neurons in the rat central nervous system. *Neuroscience* 3, 517-538.
- Kedzierska L. (1970). The influence of serotonin and histamine on contractility of the smooth muscle of the digestive tract in the experimental animals. *Arch. Immunol. Ther. Exp.* 18, 239-257.
- Keith I.M. and Ekman R. (1988). Calcitonin gene-related peptide in hamster lung and its coexistence with serotonin. A chemical and immunocytochemical study. *Regul. Pept.* 22, 315-323.
- Keith I.M. and Will J.A. (1982). Dynamics of the neuroendocrine cell regulatory peptide system in the lung. Specific overview and new results. *Exp. Lung Res.* 3, 387-402.

Immunoreactive structures in mouse lung

- Laitinen A., Partanen M., Hervonen A., Petohuikko M. and Laitinen L.A. (1985). Electron microscopic study on the innervation of the human lower respiratory tract. Evidence of adrenergic nerves. *Eur. J. Respir. Dis.* 67, 209-215.
- Lauweryns J.M. and Cokelaere M. (1973a). Intrapulmonary neuroepithelial bodies, hypoxia-sensitive neuro (chemo-) receptors. *Experientia* 29, 1384-1386.
- Lauweryns J.M. and Cokelaere M. (1973b). Hypoxia-sensitive neuroepithelial bodies, intrapulmonary secretory neuroreceptors, modulated by the CNS. *Z. Zellforsch.* 145, 521-540.
- Lauweryns J.M. and Goddeeris P. (1975). Neuroepithelial bodies in the human child and adult lung. *Am. Rev. Respir. Dis.* 3, 468-476.
- Lauweryns J.M. and Peuskens J.C. (1972). Neuro-epithelial bodies (neuroreceptor or secretory organs?) in human infant bronchial and bronchiolar epithelium. *Anat. Rec.* 172, 471-481.
- Lauweryns J.M. and Van Lommel A. (1983). The intrapulmonary intraepithelial bodies after vagotomy, demonstration of their sensory neuroreceptor like innervation. *Experientia* 39, 1123-1124.
- Lauweryns J.M. and Van Ranst L. (1987). Leu-7 immunoreactivity in human, monkey and pig bronchopulmonary neuroepithelial bodies and neuroendocrine cells. *J. Histochem. Cytochem.* 35, 687-691.
- Lauweryns J.M., Cokelaere M. and Theunynck P. (1973). Serotonin producing NEB in rabbit respiratory mucosa. *Science* 180, 410-413.
- Lauweryns J.M., Cokelaere M., Theunynck P. and Deleersnyder M. (1974). Neuroepithelial bodies in mammalian respiratory mucosa, light optical, histochemical and ultrastructural studies. *Chest* 65, 22S-29S.
- Lauweryns J.M., Cokelaere M., Deleersnyder M. and Liebens M. (1977). Intrapulmonary neuroepithelial bodies in newborn rabbits. Influence of hypoxia, hyperoxia, hypercapnia, nicotina, reserpine, L-dopa and 5-HTP. *Cell Tissue Res.* 182, 425-440.
- Lauweryns J.M., De Bock V., Verhofstad A.A. and Steinbusch H.W. (1982). Immunohistochemical localization of serotonin in intrapulmonary neuro-epithelial bodies. *Cell Tissue Res.* 226, 215-223.
- Lilly C.M., Drazen J.M. and Shore S.A. (1993). Peptidase modulation of airway effects of neuropeptides. *Proc. Soc. Exp. Biol. Med.* 203, 388-404.
- Lundberg J.M., Terenius L., Hökfelt T., Martling C.R., Tatemoto K., Mutt V., Polak J.M., Blomm S. and Goldstein M. (1982). Neuropeptide Y (NPY)-like immunoreactivity in peripheral noradrenergic neurons and effects of NPY on sympathetic function. *Acta physiol. Scand.* 116, 477-480.
- Lundberg J.M., Hökfelt T., Martling C.R., Saria A. and Cuello C. (1984). Substance P-immunoreactive sensory nerves in the lower respiratory tract of various mammals including man. *Cell Tissue Res.* 235, 251-261.
- Lundberg J.M., Martling C.R. and Hökfelt T. (1988). Airway innervation. In: *Handbook of chemical neuroanatomy. Vol. 6. The peripheral nervous system.* Björklund A. and Hökfelt T. (eds). Elsevier, Amsterdam-New York-Oxford. pp 391-444.
- Luts A., Uddman R., Alm P., Basterra J. and Sundler F. (1993). Peptide-containing nerve fibres in human airways, distribution and coexistence pattern. *Int. Arch. Allergy. Immunol.* 101, 52-60.
- Marcheusky A.M. and Kleinerman J. (1983). Quantitative and immunocytochemical studies of APUD cells in airways and gut. *Lung* 161, 67-76.
- Martling C.R., Matran R., Alving K., Hökfelt T. and Lundberg J.M. (1990). Innervation of lower airways and neuropeptide effects on bronchial and vascular tone in the pig. *Cell Tissue Res.* 260, 223-233.
- Nilsson G., Dahlberg K., Brodin E., Sundler F. and Strandberg K. (1977). Distribution and constrictor effects of substance P in guinea pig tracheobronchial tissue. In: *Substance P.* Euler V.S. von and Pernow B. (eds). Raven Press. New York. pp 57-61.
- Polak J.M. and Bloom S.R. (1982). Distribution of the regulatory peptides in the respiratory tract of man and mammals. In: *Systemic role of regulatory peptides.* Bloom S.R., Polak J.M. and Lindenlaub E. (eds). F.K. Schattauer Verlag. Stuttgart. pp 241-269.
- Polak J.M. and Bloom S.R. (1986). Regulatory peptides of the gastrointestinal and respiratory tracts. *Arch. Int. Pharmacodyn. Ther.* 280, 16-49.
- Redick M.L. and Hung K.S. (1984). Quantitation of pulmonary neuroepithelial bodies in pre and postnatal rabbits. *Cell Tissue Res.* 238, 583-587.
- Richardson J.B. (1979). Nerve supply to the lungs. *Am. Rev. Respir. Dis.* 119, 785-802.
- Said S.I. (1982). Vasoactive intestinal peptide (VIP and the lung). In: *Systemic role of regulatory peptides.* Bloom S.R., Polak J.M. and Lindenlaub E. (eds). F.K. Schattauer Verlag. Stuttgart. pp 293-305.
- Saria A., Martling C.R., Theodorsson-Norheim E., Gamse R., Hua X.Y. and Lundberg J.M. (1987). Coexisting peptides in capsaicin-sensitive sensory neurons. Release and actions in the respiratory tract of the guinea pig. *Act. Physiol. Hung.* 69, 421-424.
- Sheppard M.N., Polak J.M., Allen J.M. and Bloom S.R. (1984). Neuropeptide Tyrosine (NPY), a newly discovered peptide is present in the mammalian respiratory tract. *Thorax* 39, 326-330.
- Shimosegawa T., Foda H.D. and Said S.I. (1989). Immunohistochemical demonstration of enkephalin-containing nerve fibers in guinea pig and rat lungs. *Am. Rev. Respir. Dis.* 140, 441-445.
- Sonstegard K.S., Mailman R.B., Cheek J.M., Tomlin T.E. and DiAugustine R.P. (1982). Morphological and cytochemical characterization of neuroepithelial bodies in fetal rabbit lung. I-Studies of isolated NEB. *Exp. Lung Res.* 3, 348-377.
- Terenghi G., Mc Gregor G.P., Bhattacharji S., Wharton J., Bloom S.R. and Polak J.M. (1983). Vagal origin of substance P-containing nerves in the guinea pig lung. *Neurosci. Lett.* 36, 229-235.
- Uddman R., Alumets J., Densert O., Hakanson R. and Sundler F. (1978). Occurrence and distribution of VIP nerves in the nasal mucosa and tracheobronchial wall. *Acta Otolaryngol.* 86, 443-448.
- Uddman R., Sundler F. and Emson P. (1984). Occurrence and distribution of neuropeptide Y-immunoreactive nerves in the respiratory tract and middle ear. *Cell Tissue Res.* 237, 321-327.
- Wang Y.Y. and Cutz E. (1993). Localization of colecystokinin-like peptide in neuroendocrine cells of mammalian lungs, a light and electron microscopic immunohistochemical study. *Anat. Rec.* 236, 198-205.
- Wasano K. and Yamamoto T. (1978). Monoamine-containing granulated cell in the frog lung. *Cell Tissue Res.* 192, 201-209.
- Wasano K. and Yamamoto T. (1981). A scanning and transmission electron-microscopic study on NEB in the neonatal mouse lung. *Cell Tissue Res.* 216, 481-490.
- Wharton J., Polak J.M., Bloom S.R., Hakanson R. and Sundler F. (1978). Bombesin-like immunoreactivity in the lung. *Nature* 273, 769-770.
- Wharton J., Polak J.M., Bloom S.R., Will J.A. and Pearse A.G.E. (1979). Substance P-like immunoreactive nerves in mammalian lung. *Invest. Cell. Pathol.* 2, 3-10.