

An Approach to the Factors Related to Sperm Capacitation Process

Rebeca López-Úbeda and Carmen Matás*

Department of Physiology, University of Murcia, Campus Mare Nostrum, 30071, Murcia, Spain

Abstract

This review summarizes some information about the different ways in relation to sperm capacitation. On one hand, the classical pathway that define the functional changes that occur in sperm during *in vitro* capacitation with special emphasis on the factors that lead to the tyrosine Phosphorylation (PY), and on the other hand, molecules and process that are involved in new mechanisms involved in this event like reactive species, especially Nitric Oxide (NO) and protein nitrosylation.

Keywords: Sperm capacitation; *In vitro*; Protein nitrosylation; Phosphorylation

Introduction

After mating or artificial insemination, millions of sperm are deposited in the female genital tract, of which only a small proportion is able to reach the caudal portion of the isthmus (Figure 1A). This sperm population encounters a sticky secretion of glycoprotein that modifies the sperm surface [1]. Motility decreases in this viscous medium and facilitates the sperm adhesion to the epithelium (Figure 1B). Sperm remains in the caudal portion of the oviductal isthmus, during the pre and peri-ovulatory time, forming the Sperm Reservoir (SR) [2,3]. This binding is a reversible process and the oviductal microenvironment signals stimulate sequential release of a limited number of sperm from the SR to the ampullary-isthmic junction. This ensures fertilization of oocytes in a time interval (Figure 1C), even if ovulation occurs over a long period of time [4]. However, the mechanisms which sperm are released from SR are unknown.

During the passage of sperm through the female genital tract, the spermatozoa undergo functional and molecular changes which confer ability to fertilize the oocyte (Figure 1). This process is known as sperm capacitation [5].

Capacitation is a complex process, which appears to be controlled by crosstalk between different pathways [6,7]. The most notable event is an increase in protein tyrosine phosphorylation [8,9] although an oxidative process also has been shown, including the nitric oxide-dependent pathway [10,11]. In this sense, some papers have shown that NOS (Nitric Oxide Synthase) is present in the oviduct [12-14], oocyte, and cumulus and corona cells [15,16] of different species [12,17,18]. NOS isoforms are hormonally regulated in the oviduct and expresses differently throughout the oestrous cycle. In the oviduct, Nitric Oxide (NO) has been shown to regulate contractility [19], ciliary beating of the ciliated epithelial cells, the sperm motility or even inducing chemotaxis [20]. For this reason NO also module sperm capacitation although the pathway is not known totally.

Functional Changes and Molecular Pathways during Sperm Capacitation

During the capacitation process, spermatozoa undergo a series of functional changes, which enables them to bind to the extracellular matrix of the oocyte and consequently require the acrosome reaction. Although the latter is under discussion as recently shown by Jin et al. [21] that the acrosome reaction in mouse sperm occurs before binding to the zona pellucida. Besides, the pattern of movement of sperm flagellum changes allowing penetration of the zona pellucida [22].

Capacitation process implied several changes sequentially. Some of these changes are rapid and occur at the moment of ejaculation. Others require a longer period of time in the female genital tract (*in vivo*) or in a medium capable of supporting this process (*in vitro*). All these processes (both rapid and slow), appear to be regulated by protein kinase A (PKA) and HCO₃⁻, Soluble Adenylate Cyclase (SACY or sAC), and Cyclic Adenosine 3'5 'Monophosphate (cAMP) participate in this process (revised by [23]).

Traditionally, Reactive Oxygen Species (ROS) are considered to be injurious by products of cellular metabolism but also fundamentally participants in cell signalling and regulation mechanisms [24]. This apparent paradox also is true for spermatozoa, which are particularly susceptible to ROS-induced damage because their plasma membranes contain relatively large amounts of polyunsaturated fatty acids and their cytoplasm contains relatively low concentrations of scavenging enzymes [25], but require low concentration of ROS to acquire the fertilizing ability [6,26]. The essential role of ROS as modulators of capacitation is recognized in human [27], bovine [28], and mouse [27], and boar spermatozoa [29,30].

Some Authors Consider that Capacitation Occurs in Two Steps, Fast and Slow [23]

Facts during fast sperm capacitation

An early event during capacitation is the activation of sperm motility. Although sperm stored in the cauda epididymis being practically immobile consume oxygen in large proportions. The flagellum movement starts immediately after sperm are released from the epididymis and contact has been made with seminal plasma. This is due to exposure of sperm to the HCO₃⁻ [31].

Facts during slow sperm capacitation


In contrast to the rapid activation of motility, other processes

*Corresponding author: Carmen Matás, Facultad de Veterinaria, Campus de Espinardo, University of Murcia, Murcia 30100, Spain, Tel: 34868887256; Fax: 34868884147; E-mail: cmatas@um.es

Received February 06, 2015; Accepted February 17, 2015; Published February 25, 2015

Citation: López-Úbeda R, Matás C (2015) An Approach to the Factors Related to Sperm Capacitation Process. Andrology 4: 128. doi:10.4172/2167-0250.1000128

Copyright: © 2015 López-Úbeda R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



associated with capacitation require a longer period of time. During slow capacitation, sperm acquire the ability to fertilize, which is preceded by the preparation of the sperm to undergo the acrosome reaction and change the pattern of motility called hyperactivation. Components in oviductal fluid such as high weight molecular proteins and high density lipoproteins promote cholesterol efflux resulting in an increased capacitation and tyrosine Phosphorylation (PY) using the cAMP signalling pathway/PKA [32]. Additionally, these slow processes also are achievable *in vitro* by incubation of spermatozoa in defined media, which contain a protein source (usually bovine serum albumin (BSA)), and different ions, including HCO_3^- and Ca_2^+ .

Molecules and Mechanisms Involved in the Process of Capacitation

As we mentioned above, bicarbonate, calcium and cholesterol acceptor are essential during capacitation process entirely (Figure 2). These substances induce modifications lipid membrane, loss of cholesterol, activation of cAMP/PKA pathway, increase Ca_2^+ uptake and pH (pHi), hyperpolarisation of membrane potential, and PY [33]. However, there are other pathways in relation to capacitation as NO/sGC/cGMP or protein nitrosylation are being studied.

Bicarbonate and sperm capacitation


Several studies have shown that bicarbonate plays a key role in sperm capacitation and therefore achieve fertilization under both *in vivo* and *in vitro* [34-38]. Epididymal spermatozoa are exposed to low bicarbonate concentrations (3-4mM). However, when they arrive before the capacitation, takes place (oviduct). They are found in much higher level (> 20mM) [39]. Movement of HCO_3^- through the membrane has been associated with increased intracellular pH during capacitation [40]. Moreover, another likely target for the action of bicarbonate on sperm metabolism is the regulation of cAMP [41] by stimulation of sAC [23]. This in turn stimulates PKA to phosphorylate substrates, thereby allowing PY [42,43]. Furthermore, activation of

the PKA results in activation of phospholipase D (PLD), which in turn stimulates the polymerization of F-actin [44], which is an event associated with the process of acrosome reaction.

Bicarbonate and lipid membrane structure: Bicarbonate also modifies the lipid structure of the sperm plasma membrane during capacitation and it is a reversible phenomenon (Figure 2) [36]. Gadella and Harrison [45] showed that influx bicarbonate during capacitation produces change in the lipid membrane structure using path sAC/cAMP/PKA and so is augmented by inhibitors of phosphatases [46]. These changes lead to a reordering of membrane phospholipids phosphatidylethanolamine, phosphatidylserine, sphingomyelin and phosphatidylcholine. Lipid reordering allows relocating the cholesterol in the apical part of the sperm head. Apparently, this relocation has the function of removal of cholesterol [47]. Albumin, High-Density Lipoprotein (HDL), and β -cyclodextrins promote sperm capacitation acting as acceptors of cholesterol by removing it from the plasma membrane [48]. As a result of this process, decrease ratio of cholesterol/phospholipid consequently contributes to an increased membrane fluidity promoting increase of ion permeability [32,49-51].

Bicarbonate and sterol depletion: Albumin acts in synergy with bicarbonate by mediating efflux of sterols from the sperm surface [52,53]. Flesch et al. [47] observed that the addition of albumin causes cholesterol efflux (Figure 2), but only in bicarbonate-responding cells that exhibited virtually no filipin labelling in the sperm head area. In the absence of bicarbonate, albumin had no effect on other lipid components and no affinity to cholesterol. Bicarbonate also induces sperm surface oxysterol formation by activation of signalling pathway of the ROS, which can be inhibited or blocked by addition of antioxidants as vitamin E or vitamin A [38]. These sterols oxidation products (oxysterols), which are more hydrophilic, can be extracted using albumin [53] or can facilitate an oxysterol dependent scavenger-sensitive transport of free sterols to albumin [54].

Bicarbonate and sperm plasma membrane potential: Under normal conditions, spermatozoa maintain intracellular ion concentration markedly different from extracellular environment and these differences provide the resting membrane potential [55].



When spermatozoa are exposed to different environments during transport through the male and female genital tracts, they find different extracellular ion concentration. For example, the epididymal fluid contains high K^+ , low Na^+ , and even lower concentrations of HCO_3^- . After ejaculation, there will be a drastic change in the concentrations of these ions in the seminal fluid and finally into the female tract, where the concentrations of low potassium and high HCO_3^- are present [56,57]. As a result of changes in extracellular ion concentrations, there will be changes in intracellular concentrations of these ions leaving alterations in membrane potential [58,59] which consequently occurs in the hyperpolarisation of sperm plasma membrane [60]. It has been shown in mouse sperm that changes in membrane potential do not occur in BSA or HCO_3^- absence [59]. These results suggest that HCO_3^- present in capacitation media as well as cholesterol efflux may have a direct or indirect function of events allowing hyperpolarisation of the sperm plasma membrane [55]. Arnoult et al. [61] showed that only hyperpolarized sperm populations are capable of undergoing the acrosome reaction in presence of solubilised zona pellucida material.

Calcium and sperm capacitation

In 1915, Loeb [62] was the first to demonstrate that Ca^{2+} is required in the extracellular medium for fertilization to occur in invertebrates. Of all intracellular signalling mechanisms, perhaps the most studied and best characterized one is the mobilization of Ca^{2+} . This pathway involves transitory increase of intracellular calcium concentrations produced by multitude intercellular messengers.

One of the most important consequences of cholesterol efflux from membranes is a massive influx of extracellular Ca^{2+} , which is considered a prerequisite for the acrosome reaction process [63]. This Ca^{2+} influx may be due to changes occurring in the membrane fluidity. The intracellular Ca^{2+} increase in sperm can activate one or more enzymatic pathways (Figure 2). For example, the Adenylate Cyclase (AC) increases during capacitation in response to Ca^{2+} , this enzyme will catalyse the conversion of ATP to cAMP (revised by [48]).

In 1998, Visconti and Kopf [8] suggested a cooperative effect of Ca^{2+} and HCO_3^- in modulating sperm capacitation requiring the presence of both as well as increase in cAMP levels and the subsequent phosphorylation of different proteins. In swine, both Ca^{2+} and HCO_3^- appear to be required for capacitation and their roles are synergistic, since it has been shown that the HCO_3^- will stimulate the entry of Ca^{2+} in this species [64]. However, in mouse spermatozoa, Tateno et al. [65] showed that Ca^{2+} ionophore A23187 can make spermatozoa capable of fertilizing *in vitro* without activation of cAMP-dependent phosphorylation pathways in media bicarbonate free.

Ca^{2+} is important to sperm hyperactivation during capacitation. CatSperm 1 and 2 are voltage dependent calcium channels that are located in the tail of the sperm. Sperm from mice deficient in these calcium channels are infertile and do not exhibit hyperactivation during capacitation despite having PY [66].

Another aspect that influences capacitation related to calcium is intracellular pH. Sperm not capacitated maintain an acidified intracellular pH [67]. This fact acts as a regulator of calcium influx [68] preventing capacitation and acrosome reaction. Intracellular pH becomes more alkaline during capacitation [69]. Today it is believed that increasing intracellular calcium, bicarbonate and the pH during sperm capacitation produce sAC activation and consequently cAMP [9,51,70,71].

In addition, calmodulin, which is a protein binding Ca^{2+} considered to be an important transducer of calcium signals, appears to be diminished during capacitation. This mechanism could be based on inhibition of Ca^{2+} -ATPase plasma membrane by increasing cAMP levels through PDE1 inhibition (reviewed by [72]).

Tyrosine phosphorylation of sperm proteins

Protein phosphorylation or de-phosphorylation is controlled by activity of protein kinases and protein phosphatases, which provide cells a “switch” through which they can activate function of various proteins [73]. Phosphorylation occurs in serine, threonine, and tyrosine. PY is related to capacitation process and sperm hyperactivation in many mammal species (human [74], bovine [75], murine [8] or porcine [76]). In opposite, it has been shown that protein phosphatases decrease their activity during capacitation [73].

Increasing PY during capacitation is regulated by a cAMP-dependent pathway which involves PKA [77]. cGMP-PKG pathway is also involved in this process [78]. In 2002, Visconti et al. [79] described the possible mechanisms, which could regulate the PY dependent signalling pathway cAMP/PKA: a) the direct or indirect stimulation of a tyrosine kinase by PKA, b) the direct or indirect inhibition of a tyrosine phosphatase, and c) direct or indirect phosphorylation of proteins by PKA on serine or threonine residues to prepare these proteins for subsequent phosphorylation on tyrosine residues.


PY is specific for each species. For example, in man PY during sperm capacitation requires the presence of BSA, and HCO_3^- but no calcium [80]. In the case of stallion PY during capacitation requires HCO_3^- but neither BSA nor calcium [81]. Another factor to consider in PY is time. PY in boar sperm occurs close to 1 hour after the addition of bicarbonate [82], whereas in bull sperm it occurs 4 hours after addition of heparin [75].

Although PY is an important key in capacitation, it is not yet entirely clear how the phosphorylation of these proteins is involved in sperm-zona recognition, gamete interaction, or exocytosis of acrosomal content [83]. The level of PY in human sperm correlates strongly with the sperm-zona-binding capacity [84] and alterations in PY have been found in subfertile subjects [85] indicating its physiological role in fertilization. In pigs ejaculated spermatozoa selected in the oviduct adhere to the epithelial cells and suppress PY of sperm proteins. This modulation by the oviductal epithelium on PY and, therefore capacitation could help synchronize sperm functions to the time of ovulation [86].

Nitric oxide (NO) and sperm capacitation

NO, a highly ROS, has been found in several physiological systems and regulated manifold functions in male and female reproductive systems [87].

NO is a very small lipophilic molecule that can rapidly diffuse through biological membrane barriers and acts as an intracellular and extracellular biological messenger in a variety of physiological processes. NO is synthesized *in vivo* from L-arginine by the action of NOS (Figure 3), an enzyme existing in three isoforms: neuronal NOS (nNOS or NOS1), endothelial NOS (eNOS or NOS3), also referred to as constitutive NOS, responsible for the continuous basal release of NO, and both require calcium/calmodulin for activation [88,89]. A third isoform is an inducible calcium-independent form (iNOS or NOS2). NOS activity is dependent on substrate availability and the



co-factors NADPH, Flavin Mononucleotide (FMN), Flavin Adenine Dinucleotide (FAD) and tetrahydrobiopterin (BH4). The availability of these factors determines the cellular rates of NO synthesis [90].

Different NOS isoforms were detected in mammalian spermatozoa such as mouse [91], bull [92], human [93,94] and boar spermatozoa [95] activating the biosynthesis of NO. NO was able to affect sperm motility [94,96,97], acrosomal reaction [98,99], acts on PY of sperm proteins [100,101] and enhancement of sperm-zona pellucida binding ability [102].

NO has different functions in the spermatozoa, acting on different pathways that result in sequential and parallel processes (Figure 3). The main actions of NO are:

A. Activation of soluble guanylate cyclase (sGC)

The most important intracellular signalling role for NO in the spermatozoa is its capacity to activate the soluble isoform of Guanylate Cyclase (sGC) [103]. Activation of sGC (Figure 4) leads to increase in intracellular levels of cGMP, which has been implicated in several sperm signalling pathway functions, such as capacitation, acrosome reaction, chemotaxis, and sperm-egg interaction [20,27,99,104].

There are at least three targets of cGMP: Cyclic Nucleotide-Gated (CNG) channels, cGMP-dependent Protein Kinase (PKG) and Phosphodiesterase (PDE), involved in several physiological events in mammalian spermatozoa. All of these targets result in increased levels of intracellular calcium and the phosphorylation of different proteins causing sperm hyperactivation and acrosome reaction. Calcium influx together with increased protein phosphorylation brings about the capacitation response (Figure 4).

A1: Cyclic Nucleotide-Gated Channels (CNG): CNG channels have been expressed in mammalian sperm [105] mainly along the length of the flagellum [105,106]. They are activated by cGMP and have been proposed to mediate the influx of Ca^{2+} to the cytoplasm during capacitation in mammalian spermatozoa controlling sperm motility [106]. This signalling pathway involving CNG channel activation using cGMP is one of the first events that occurred during capacitation in the mouse sperm (Figure 4A1) [78].

A2: cGMP-dependent Protein Kinase (PKG): PGK is a major cellular receptor of cGMP and plays important roles in cGMP-

dependent signal transduction pathways. Previous studies have identified in mammals two forms of PGK (I and II) [107,108], that are encoded by distinct genes and two different isoforms of PGK-I (designated Ia and Ib) that are produced by alternative splicing [109]. PGK-I seem to play an important role in mediating the acrosome reaction [20,99], modulating several sperm motion patterns and sperm chemotaxis.

The increase in cGMP in the cytoplasm and the subsequent activation of PKG [107,108] results in protein serine/threonine phosphorylation (Figure 4A2) [110], which might also indirectly, mediate a new calcium entry [78]. This promotes sperm capacitation and acrosome reaction [111].

A3: Phosphodiesterase (PDE): Evidence was provided for the involvement of PDE in sperm motility and capacitation [112]. cGMP and cAMP compete for catalytic sites of PDEs that hydrolyse both cyclic nucleotides [113,114]. A rise in intracellular levels of cGMP could inhibit cAMP degradation via cyclic nucleotide phosphodiesterase type 3 [115], which increase intracellular cAMP levels and, consequently, cause an activation of PKA [116] and indirectly increase protein (PY) (Figure 4A3).

B. Protein tyrosine phosphorylation



NO appears to be involved in PY through different mechanisms, acting on two essential pathways for sperm capacitation: on cAMP/PKA or Extracellular Signal Regulated Kinase (ERK) pathway (Figure 5). These mechanisms for the control of PY are not mutually exclusive neither excluding, both pathways act in parallel [28].

NO can influence the cAMP/PKA pathway by activation of sGC (see above) it could also modulate directly sperm Adenylate Cyclase Activity (AC). The activation of PKA represents the point of convergence for these two pathways. Low concentration of NO could stimulate AC with a subsequent increase in cAMP levels [117] to increase PY by activation of PKA. However, high concentrations of NO can inhibit AC [118]. McVey *et al* [118] also demonstrated that the effects of NO on AC activity are reversible, suggesting S-nitrosylation of AC as a possible mechanism of action of the NO (Figure 5).

ERK pathway is a chain of many proteins (Shc, Grb2, Sos, Ras and ERK1/2 module, which includes three kinases sequentially activated: Raf, MEK, and ERK1/2). NO intervenes in the middle of the pathway, modify the Ras structure [119], reacting with cysteine residues [28] and inducing its activation. Ras interact with Raf by activating it [119], leading to MEK activation. MEK phosphorylates Thr and Tyr are residues within the Thr-Glu-Tyr motif, which are located at the active site of ERKs 1 and 2 (Figure 5). All this process is necessary for the subsequent PY [28] and is involved in the acquisition of sperm motility [120].

C. Direct modulation of protein function by S-nitrosylation of exposed cysteine residues

Mature sperm lack the necessary machinery for the transcription or protein modification and thereby require post-translational modifications to control the activity of proteins. NO participates in protein regulation, which acts directly on protein targets (exposed cysteine residues) via S-nitrosylation [121,122]. S-nitrosylation is a regulated post-translational protein modification (Figure 6), analogous to phosphorylation and acetylation [123,124], which



involves the covalent incorporation of a NO into thiol groups (-SH), to form S-nitrosothiol (S-NO). This modification is selective, reversible and stabilizes NO in a uniquely bioactive form. Lefèvre et al. [125] described numerous sperm proteins that can be nitrosylated in human sperm. Some of the S-nitrosylated proteins are involved in processes related to sperm capacitation as energy generation, sperm motility [125] or hyperactivation [126,127]. However, how some of nitrosylated proteins perform their function remains unknown.

For sperm, hyperactivation is necessary mobilisation of stored Ca^{2+} in the sperm neck/midpiece [127]. The calcium store in the neck of the sperm takes place in the Redundant Nuclear Envelope (RNE) [128]. To mobilize calcium from these stores is necessary to enable ryanodine receptors (RyRs); intracellular calcium release channels involved in regulation of cytosolic calcium levels [129]. These proteins contain a large number of thiol groups and are thus subject to S-nitrosylation by NO [130,131]. S-nitrosylation can potentiate opening of RyR [132-137], probably by generations of the membrane permeant product cys-NO [138]. A NO-induced Ca^{2+} elevation was accompanied by an increase in S-nitrosylation levels of endogenous RyR [139,140] and also inhibition of these Ca^{2+} channels can occur under strongly nitrosylating conditions or at high doses of NO (Figure 6) [134,137,140].


Progesterone acts synergistically with NO (by S-nitrosylation) to mobilise Ca^{2+} at the sperm neck/midpiece (by activation of RyRs) [126] contributing to the hyperactivation that is vital for penetration of the egg vestments.

Summary and Perspective

The sperm from being deposited in the female genital until it reaches the place of fecundation undergoes a series of changes known classically as capacitation. This process involves modifications membrane lipids, loss of cholesterol, activation of cAMP/PKA pathway, increase Ca_i^{2+} uptake and pH (pHi), hyperpolarisation of membrane potential and tyrosine phosphorylation. Among the most studied molecules associated with this process include BSA as cholesterol acceptor, bicarbonate as an activator of the cAMP-PKA-tyrosine phosphorylation, and calcium as an activator of channels voltage-dependent and hyperactivity motility process. However, there are other pathways in relation to capacitation as NO/sGC/cGMP, which have some common steps to cAMP-PKA-tyrosine phosphorylation pathways. Another pathway recently described relating to capacitation process is the protein nitrosylation. Nevertheless, this new way of signalling involves numerous proteins whose functions are yet to be determined and that can be important to understand the complex process of sperm capacitation.

Acknowledgements

Supported by Spanish Ministry of Economy and Competitiveness and FEDER, Grant AGL2012-40180-C03-01



References

- Rodríguez-Martínez H, Tienthai P, Suzuki K, Funahashi H, Ekwall H, et al. (2001) Involvement of oviduct in sperm capacitation and oocyte development in pigs. *Reprod Suppl* 58: 129-145.
- Suarez S, Redfern K, Raynor P, Martin F, Phillips DM (1991) Attachment of boar sperm to mucosal explants of oviduct in vitro: possible role in formation of a sperm reservoir. *Biol Reprod* 44: 998-1004.
- Töpfer-Petersen E, Wagner A, Friedrich J, Petrunkina A, Ekhlasi-Hundrieser M, et al. (2002) Function of the mammalian oviductal sperm reservoir. *J Exp Zool* 292: 210-215.
- Rodríguez-Martínez H, Saravia F, Wallgren M, Roca J, Peña FJ (2008) Influence of seminal plasma on the kinematics of boar spermatozoa during freezing. *Theriogenology* 70: 1242-1250.
- AUSTIN CR (1952) The capacitation of the mammalian sperm. *Nature* 170: 326.
- de Lamirande E, Leclerc P, Gagnon C (1997) Capacitation as a regulatory event that primes spermatozoa for the acrosome reaction and fertilization. *Mol Hum Reprod* 3: 175-194.
- Fraser LR (2010) The "switching on" of mammalian spermatozoa: molecular events involved in promotion and regulation of capacitation. *Mol Reprod Dev* 77: 197-208.
- Visconti PE, Kopf GS (1998) Regulation of protein phosphorylation during sperm capacitation. *Biol Reprod* 59: 1-6.
- Breitbart H (2003) Signaling pathways in sperm capacitation and acrosome reaction. *Cell Mol Biol (Noisy-le-grand)* 49: 321-327.
- Zhang H, Zhou Q, Li X, Zhao W, Wang Y, et al. (2007) Ginsenoside Re promotes human sperm capacitation through nitric oxide-dependent pathway. *Mol Reprod Dev* 74: 497-501.
- de Lamirande E, Lamothe G (2009) Reactive oxygen-induced reactive oxygen formation during human sperm capacitation. *Free Radic Biol Med* 46: 502-510.
- Rosselli M, Dubey RK, Rosselli MA, Macas E, Fink D, et al. (1996) Identification of nitric oxide synthase in human and bovine oviduct. *Mol Hum Reprod* 2: 607-612.
- Ekerhovd E, Brännström M, Weijdegård B, Norström A (1999) Localization of nitric oxide synthase and effects of nitric oxide donors on the human Fallopian tube. *Mol Hum Reprod* 5: 1040-1047.
- Lapointe J, Roy M, St-Pierre I, Kimmins S, Gauvreau D, et al. (2006) Hormonal and spatial regulation of nitric oxide synthases (NOS) (neuronal NOS, inducible NOS, and endothelial NOS) in the oviducts. *Endocrinology* 147: 5600-5610.
- Reyes R, Vázquez ML, Delgado NM (2004) Detection and bioimaging of nitric oxide in bovine oocytes and sperm cells. *Arch Androl* 50: 303-309.
- Tao Y, Fu Z, Zhang M, Xia G, Yang J, Xie H (2004) Immunohistochemical localization of inducible and endothelial nitric oxide synthase in porcine ovaries and effects of NO on antrum formation and oocyte meiotic maturation. *Molecular and cellular endocrinology* 222: 93-103.
- Ekerhovd E, Brännström M, Alexandersson M, Norström A (1997) Evidence for nitric oxide mediation of contractile activity in isolated strips of the human Fallopian tube. *Hum Reprod* 12: 301-305.
- Bryant CE, Tomlinson A, Mitchell JA, Thiemermann C, Willoughby DA (1995) Nitric oxide synthase in the rat fallopian tube is regulated during the oestrous cycle. *J Endocrinol* 146: 149-157.
- Rosselli M, Imthurn B, Macas E, Keller P, Dubey R (1994) Endogenous nitric oxide modulates endothelin-1 induced contraction of bovine oviduct. *Biochemical and biophysical research communications* 201: 143-148.
- Miraglia E, Rullo ML, Bosia A, Massobrio M, Revelli A, et al. (2007) Stimulation of the nitric oxide/cyclic guanosine monophosphate signaling pathway elicits human sperm chemotaxis in vitro. *Fertil Steril* 87: 1059-1063.
- Jin M, Fujiwara E, Kakiuchi Y, Okabe M, Satouh Y, et al. (2011) Most fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona pellucida during in vitro fertilization. *Proceedings of the National Academy of Sciences* 108: 4892-4896.
- Suarez SS (2008) Control of hyperactivation in sperm. *Hum Reprod Update* 14: 647-657.
- Visconti PE (2009) Understanding the molecular basis of sperm capacitation through kinase design. *Proc Natl Acad Sci U S A* 106: 667-668.
- Finkel T (2001) Reactive oxygen species and signal transduction. *IUBMB Life* 52: 3-6.
- Aitken J, Fisher H (1994) Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk. *Bioessays* 16: 259-267.
- Aitken RJ (1997) Molecular mechanisms regulating human sperm function. *Mol Hum Reprod* 3: 169-173.
- Herrero MB, de Lamirande E, Gagnon C (2003) Nitric oxide is a signaling molecule in spermatozoa. *Curr Pharm Des* 9: 419-425.
- O'Flaherty C, de Lamirande E, Gagnon C (2006) Reactive oxygen species modulate independent protein phosphorylation pathways during human sperm capacitation. *Free Radical Biology and Medicine* 40: 1045-1055.
- Funahashi H (2002) Induction of capacitation and the acrosome reaction of boar spermatozoa by L-arginine and nitric oxide synthesis associated with the anion transport system. *Reproduction* 124: 857-864.
- Aquila S, Giordano F, Guido C, Rago V, Carpino A (2011) Nitric oxide involvement in the acrosome reaction triggered by leptin in pig sperm. *Reprod Biol Endocrinol* 9: 133.
- Wennemuth G, Carlson AE, Harper AJ, Babcock DF (2003) Bicarbonate actions on flagellar and Ca²⁺-channel responses: initial events in sperm activation. *Development* 130: 1317-1326.
- Visconti PE, Ning X, Fornés MW, Alvarez JG, Stein P, et al. (1999) Cholesterol efflux-mediated signal transduction in mammalian sperm: cholesterol release signals an increase in protein tyrosine phosphorylation during mouse sperm capacitation. *Developmental biology* 214: 429-443.
- Aitken RJ, Nixon B (2013) Sperm capacitation: a distant landscape glimpsed but unexplored. *Mol Hum Reprod* 19: 785-793.
- Okamura N, Tajima Y, Soejima A, Masuda H, Sugita Y (1985) Sodium bicarbonate in seminal plasma stimulates the motility of mammalian spermatozoa through direct activation of adenylate cyclase. *Journal of Biological Chemistry* 260: 9699-9705.
- Visconti PE, Bailey JL, Moore GD, Pan D, Olds-Clarke P, et al. (1995) Capacitation of mouse spermatozoa. I. Correlation between the capacitation state and protein tyrosine phosphorylation. *Development* 121: 1129-1137.
- Harrison RA, Ashworth PJ, Miller NG (1996) Bicarbonate/CO₂, an effector of capacitation, induces a rapid and reversible change in the lipid architecture of boar sperm plasma membranes. *Mol Reprod Dev* 45: 378-391.
- Harrison RA (2004) Rapid PKA-catalysed phosphorylation of boar sperm proteins induced by the capacitating agent bicarbonate. *Mol Reprod Dev* 67: 337-352.
- Boerke A, Brouwers JF, Olkkonen VM, van de Lest CH, Sostaric E, et al. (2013) Involvement of bicarbonate-induced radical signaling in oxysterol formation and sterol depletion of capacitating mammalian sperm during in vitro fertilization. *Biology of reproduction* 88: 21.
- Rodríguez-Martínez H, Ekstedt E, Einarsson S (1990) Acidification of epididymal fluid in the boar. *Int J Androl* 13: 238-243.
- Zeng Y, Oberdorf JA, Florman HM (1996) pH Regulation in Mouse Sperm: Identification of Na⁺, Cl⁻, and [formula] Dependent and Arylamino benzoate-Dependent Regulatory Mechanisms and Characterization of Their Roles in Sperm Capacitation. *Developmental biology* 173: 510-520.
- Garbers DL, Tubb DJ, Hyne RV (1982) A requirement of bicarbonate for Ca²⁺-induced elevations of cyclic AMP in guinea pig spermatozoa. *J Biol Chem* 257: 8980-8984.
- Gadella B, Harrison R (2002) Capacitation induces cyclic adenosine 3',5'-cyclic monophosphate-dependent, but apoptosis-unrelated, exposure of aminophospholipids at the apical head plasma membrane of boar sperm cells. *Biology of reproduction* 67: 340-350.
- Visconti PE, Galantino-Homer H, Moore GD, Bailey JL, Ning X, et al. (1998) The molecular basis of sperm capacitation. *J Androl* 19: 242-248.
- Cohen G, Rubinstein S, Gur Y, Breitbart H (2004) Crosstalk between protein kinase A and C regulates phospholipase D and F-actin formation during sperm capacitation. *Developmental biology* 267: 230-241.

45. Gadella B, Harrison R (2000) The capacitating agent bicarbonate induces protein kinase A-dependent changes in phospholipid transbilayer behavior in the sperm plasma membrane. *Development* 127: 2407-2420.
46. Harrison RA, Miller NG (2000) cAMP-dependent protein kinase control of plasma membrane lipid architecture in boar sperm. *Mol Reprod Dev* 55: 220-228.
47. Flesch FM, Brouwers JF, Nievelstein PF, Verkleij AJ, van Golde LM, et al. (2001) Bicarbonate stimulated phospholipid scrambling induces cholesterol redistribution and enables cholesterol depletion in the sperm plasma membrane. *Journal of cell science* 114: 3543-3555.
48. Vadnais ML, Galantino-Homer HL, Althouse GC (2007) Current concepts of molecular events during bovine and porcine spermatozoa capacitation. *Systems Biology in Reproductive Medicine* 53: 109-123.
49. Davis B (1976) Inhibitory effect of synthetic phospholipid vesicles containing cholesterol on the fertilizing ability of rabbit spermatozoa. *Experimental Biology and Medicine* 152: 257-261.
50. Cross NL (1998) Role of cholesterol in sperm capacitation. *Biol Reprod* 59: 7-11.
51. Travis AJ, Kopf GS (2002) The role of cholesterol efflux in regulating the fertilization potential of mammalian spermatozoa. *J Clin Invest* 110: 731-736.
52. Boerke A, Tsai P, Garcia-Gil N, Brewis I, Gadella B (2008) Capacitation-dependent reorganization of microdomains in the apical sperm head plasma membrane: functional relationship with zona binding and the zona-induced acrosome reaction. *Theriogenology* 70: 1188-1196.
53. Brouwers JF, Boerke A, Silva PF, Garcia-Gil N, van Gestel RA, et al. (2011) Mass spectrometric detection of cholesterol oxidation in bovine sperm. *Biol Reprod* 85: 128-136.
54. Jessup W, Gelissen IC, Gaus K, Kritharides L (2006) Roles of ATP binding cassette transporters A1 and G, scavenger receptor BI and membrane lipid domains in cholesterol export from macrophages. *Curr Opin Lipidol* 17: 247-257.
55. Salicioni AM, Platt MD, Wertheimer EV, Arcelay E, Allaire A, et al. (2007) Signalling pathways involved in sperm capacitation. *Soc Reprod Fertil Suppl* 65: 245-259.
56. Brooks DE (1983) Epididymal functions and their hormonal regulation. *Aust J Biol Sci* 36: 205-221.
57. Setchell B, Maddocks S, Brooks D (1994) Anatomy, vasculature, innervation, and fluids of the male reproductive tract. *The physiology of reproduction* 1: 1063-1175.
58. Muñoz-Garay C, De la JL, Delgado R, Labarca P, Felix R, et al. (2001) Inwardly Rectifying K⁺ Channels in Spermatogenic Cells: Functional Expression and Implication in Sperm Capacitation. *Developmental biology* 234: 261-274.
59. Demarco IA, Espinosa F, Edwards J, Sosnik J, De La Vega-Beltran JL, et al. (2003) Involvement of a Na⁺/HCO₃⁻ cotransporter in mouse sperm capacitation. *J Biol Chem* 278: 7001-7009.
60. Zeng Y, Clark EN, Florman HM (1995) Sperm membrane potential: hyperpolarization during capacitation regulates zona pellucida-dependent acrosomal secretion. *Dev Biol* 171: 554-563.
61. Arnoult C, Kazam IG, Visconti PE, Kopf GS, Villaz M, et al. (1999) Control of the low voltage-activated calcium channel of mouse sperm by egg ZP3 and by membrane hyperpolarization during capacitation. *Proceedings of the National Academy of Sciences* 96: 6757-6762.
62. Loeb J (1915) On the Nature of the Conditions Which Determine or Prevent the Entrance of the Spermatozoon Into the Egg. *The American Naturalist* 49: 257-285.
63. Flesch FM, Gadella BM (2000) Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochim Biophys Acta* 1469: 197-235.
64. Harrison RA, Mairat B, Miller NG (1993) Flow cytometric studies of bicarbonate-mediated Ca²⁺ influx in boar sperm populations. *Mol Reprod Dev* 35: 197-208.
65. Tateno H, Krapf D, Hino T, Sánchez-Cárdenas C, Darszon A, et al. (2013) Ca²⁺ ionophore A23187 can make mouse spermatozoa capable of fertilizing in vitro without activation of cAMP-dependent phosphorylation pathways. *Proceedings of the National Academy of Sciences*.
66. Carlson AE, Westenbroek RE, Quill T, Ren D, Clapham DE, et al. (2003) CatSper1 required for evoked Ca²⁺ entry and control of flagellar function in sperm. *Proc Natl Acad Sci U S A* 100: 14864-14868.
67. Parrish JJ, Susko-Parrish JL, First NL (1989) Capacitation of bovine sperm by heparin: inhibitory effect of glucose and role of intracellular pH. *Biol Reprod* 41: 683-699.
68. Florman HM, Corron ME, Kim TD-H, Babcock DF (1992) Activation of voltage-dependent calcium channels of mammalian sperm is required for zona pellucida-induced acrosomal exocytosis. *Developmental biology* 152: 304-314.
69. Vredenburg-Wilberg WL, Parrish JJ (1995) Intracellular pH of bovine sperm increases during capacitation. *Mol Reprod Dev* 40: 490-502.
70. Harrison RA, Gadella BM (2005) Bicarbonate-induced membrane processing in sperm capacitation. *Theriogenology* 63: 342-351.
71. Hess KC, Jones BH, Marquez B, Chen Y, Ord TS, et al. (2005) The α -soluble adenylyl cyclase in sperm mediates multiple signaling events required for fertilization. *Developmental cell* 9: 249-259.
72. Bailey JL (2010) Factors regulating sperm capacitation. *Syst Biol Reprod Med* 56: 334-348.
73. Signorelli JR, Díaz ES, Fara K, Barón L, Morales P (2013) Protein phosphatases decrease their activity during capacitation: a new requirement for this event. *PLoS One* 8: e81286.
74. Baldi E, Luconi M, Bonaccorsi L, Forti G (2002) Signal transduction pathways in human spermatozoa. *J Reprod Immunol* 53: 121-131.
75. Galantino-Homer HL, Visconti PE, Kopf GS (1997) Regulation of protein tyrosine phosphorylation during bovine sperm capacitation by a cyclic adenosine 3'5'-monophosphate-dependent pathway. *Biology of reproduction* 56: 707-719.
76. Tardif S, Dubé C, Chevalier S, Bailey JL (2001) Capacitation is associated with tyrosine phosphorylation and tyrosine kinase-like activity of pig sperm proteins. *Biol Reprod* 65: 784-792.
77. Kalab P, Peknicová J, Geussová G, Moos J (1998) Regulation of protein tyrosine phosphorylation in boar sperm through a cAMP-dependent pathway. *Mol Reprod Dev* 51: 304-314.
78. Cisneros-Mejorado A, Hernández-Soberanis L, Islas-Carbajal MC, Sánchez D (2014) Capacitation and Ca²⁺ influx in spermatozoa: role of CNG channels and protein kinase G. *Andrology* 2: 145-154.
79. Visconti PE, Westbrook VA, Chertihin O, Demarco I, Sleight S, et al. (2002) Novel signaling pathways involved in sperm acquisition of fertilizing capacity. *J Reprod Immunol* 53: 133-150.
80. Muratori M, Marchiani S, Tamburrino L, Forti G, Luconi M, et al. (2011) Markers of human sperm functions in the ICSI era. *Front Biosci (Landmark Ed)* 16: 1344-1363.
81. González-Fernández L, Macías-García B, Velez IC, Varner DD, Hinrichs K (2012) Calcium-calmodulin and pH regulate protein tyrosine phosphorylation in stallion sperm. *Reproduction* 144: 411-422.
82. Gadella BM, Van Gestel RA (2004) Bicarbonate and its role in mammalian sperm function. *Anim Reprod Sci* 82-83: 307-19.
83. Flesch F, Wijnand E, Van de Lest C, Colenbrander B, Van Golde L, et al. (2001) Capacitation dependent activation of tyrosine phosphorylation generates two sperm head plasma membrane proteins with high primary binding affinity for the zona pellucida. *Molecular reproduction and development* 60: 107-115.
84. Liu D, Clarke G, Baker H (2006) Tyrosine phosphorylation on capacitated human sperm tail detected by immunofluorescence correlates strongly with sperm-zona pellucida (ZP) binding but not with the ZP-induced acrosome reaction. *Human reproduction* 21: 1002-1008.
85. Buffone MG, Verstraeten SV, Calamera JC, Doncel GF (2009) High cholesterol content and decreased membrane fluidity in human spermatozoa are associated with protein tyrosine phosphorylation and functional deficiencies. *Journal of andrology* 30: 552-558.
86. Victoria Luño, Rebeca López-Úbeda, Francisco Alberto García-Vázquez, Lydia Gil, Carmen Matás (2013) Boar sperm tyrosine phosphorylation patterns in the presence of oviductal epithelial cells: in vitro, ex vivo, and in vivo models. *Reproduction* 146: 315-324.

87. O'Flaherty C, Rodriguez P, Srivastava S (2004) L-arginine promotes capacitation and acrosome reaction in cryopreserved bovine spermatozoa. *Biochim Biophys Acta* 1674: 215-221.
88. Griffith OW, Stuehr DJ (1995) Nitric oxide synthases: properties and catalytic mechanism. *Annu Rev Physiol* 57: 707-736.
89. Snyder SH (1995) Nitric oxide. No endothelial NO. *Nature* 377: 196-197.
90. Rosselli M, Keller PJ, Dubey RK (1998) Role of nitric oxide in the biology, physiology and pathophysiology of reproduction. *Hum Reprod Update* 4: 3-24.
91. Herrero MB, Goin JC, Boquet M, Canteros MG, Franchi AM, et al. (1997) The nitric oxide synthase of mouse spermatozoa. *FEBS Lett* 411: 39-42.
92. Meiser H, Schulz R (2003) Detection and localization of two constitutive NOS isoforms in bull spermatozoa. *Anat Histol Embryol* 32: 321-325.
93. Herrero MB, Pérez Martínez S, Viggiano JM, Polak JM, de Gimeno MF (1996) Localization by indirect immunofluorescence of nitric oxide synthase in mouse and human spermatozoa. *Reprod Fertil Dev* 8: 931-934.
94. O'Bryan MK, Zini A, Cheng CY, Schlegel PN (1998) Human sperm endothelial nitric oxide synthase expression: correlation with sperm motility. *Fertil Steril* 70: 1143-1147.
95. Hou ML, Huang SY, Lai YK, Lee WC (2008) Geldanamycin augments nitric oxide production and promotes capacitation in boar spermatozoa. *Anim Reprod Sci* 104: 56-68.
96. Lewis S, Donnelly E, Sterling E, Kennedy M, Thompson W, et al. (1996) Nitric oxide synthase and nitrite production in human spermatozoa: evidence that endogenous nitric oxide is beneficial to sperm motility. *Molecular human reproduction* 2: 873-878.
97. Donnelly ET, Lewis SE, Thompson W, Chakravarthy U (1997) Sperm nitric oxide and motility: the effects of nitric oxide synthase stimulation and inhibition. *Mol Hum Reprod* 3: 755-762.
98. Revelli A, Soldati G, Costamagna C, Pellerey O, Aldieri E, et al. (1999) Follicular fluid proteins stimulate nitric oxide (NO) synthesis in human sperm: a possible role for NO in acrosomal reaction. *Journal of cellular physiology* 178: 85-92.
99. Revelli A, Costamagna C, Moffa F, Aldieri E, Ochetti S, et al. (2001) Signaling pathway of nitric oxide-induced acrosome reaction in human spermatozoa. *Biol Reprod* 64: 1708-1712.
100. Herrero MB, de Lamirande E, Gagnon C (1999) Nitric oxide regulates human sperm capacitation and protein-tyrosine phosphorylation in vitro. *Biol Reprod* 61: 575-581.
101. Thundathil J, de Lamirande E, Gagnon C (2003) Nitric oxide regulates the phosphorylation of the threonine-glutamine-tyrosine motif in proteins of human spermatozoa during capacitation. *Biol Reprod* 68: 1291-1298.
102. Sengoku K, Tamate K, Yoshida T, Takaoka Y, Miyamoto T, et al. (1998) Effects of low concentrations of nitric oxide on the zona pellucida binding ability of human spermatozoa. *Fertil Steril* 69: 522-527.
103. Murad F (1994) The nitric oxide-cyclic GMP signal transduction system for intracellular and intercellular communication. *Recent Prog Horm Res* 49: 239-248.
104. Revelli A, Ghigo D, Moffa F, Massobrio M, Tur-Kaspa I (2002) Guanylate cyclase activity and sperm function. *Endocr Rev* 23: 484-494.
105. Weyand I, Godde M, Frings S, Weiner J, Müller F, et al. (1994) Cloning and functional expression of a cyclic-nucleotide-gated channel from mammalian sperm. *Nature* 368: 859-863.
106. Wiesner B, Weiner J, Middendorff R, Hagen V, Kaupp UB, et al. (1998) Cyclic nucleotide-gated channels on the flagellum control Ca²⁺ entry into sperm. *J Cell Biol* 142: 473-484.
107. Lohmann SM, Vaandrager AB, Smolenski A, Walter U, De Jonge HR (1997) Distinct and specific functions of cGMP-dependent protein kinases. *Trends Biochem Sci* 22: 307-312.
108. Pfeifer A, Ruth P, Dostmann W, Sausbier M, Klatt P, et al. (1999) Structure and function of cGMP-dependent protein kinases. *Rev Physiol Biochem Pharmacol* 135: 105-149.
109. Wernet W, Flockerzi V, Hofmann F (1989) The cDNA of the two isoforms of bovine cGMP-dependent protein kinase. *FEBS Lett* 251: 191-196.
110. Rahman MS, Kwon WS, Pang MG (2014) Calcium influx and male fertility in the context of the sperm proteome: an update. *Biomed Res Int* 2014: 841615.
111. Miraglia E, De Angelis F, Gazzano E, Hassanpour H, Bertagna A, et al. (2011) Nitric oxide stimulates human sperm motility via activation of the cyclic GMP/protein kinase G signaling pathway. *Reproduction* 141: 47-54.
112. Lefièvre L, Lamirande E, Gagnon C (2000) The Cyclic GMP-Dependent Phosphodiesterase Inhibitor, Sildenafil, Stimulates Human Sperm Motility and Capacitation but Not Acrosome Reaction. *Journal of andrology* 21: 929-937.
113. Bender AT, Beavo JA (2006) Cyclic nucleotide phosphodiesterases: molecular regulation to clinical use. *Pharmacol Rev* 58: 488-520.
114. Conti M, Beavo J (2007) Biochemistry and physiology of cyclic nucleotide phosphodiesterases: essential components in cyclic nucleotide signaling. *Annu Rev Biochem* 76: 481-511.
115. Kurtz A, Götz KH, Hamann M, Wagner C (1998) Stimulation of renin secretion by nitric oxide is mediated by phosphodiesterase 3. *Proc Natl Acad Sci U S A* 95: 4743-4747.
116. Beavo JA (1995) Cyclic nucleotide phosphodiesterases: functional implications of multiple isoforms. *Physiol Rev* 75: 725-748.
117. Belén Herrero M, Chatterjee S, Lefièvre L, de Lamirande E, Gagnon C (2000) Nitric oxide interacts with the cAMP pathway to modulate capacitation of human spermatozoa. *Free Radic Biol Med* 29: 522-536.
118. McVey M, Hill J, Howlett A, Klein C (1999) Adenylyl cyclase, a coincidence detector for nitric oxide. *J Biol Chem* 274: 18887-18892.
119. de Lamirande E, Gagnon C (2002) The extracellular signal-regulated kinase (ERK) pathway is involved in human sperm function and modulated by the superoxide anion. *Mol Hum Reprod* 8: 124-135.
120. Lu Q, Sun QY, Breitbart H, Chen DY (1999) Expression and phosphorylation of mitogen-activated protein kinases during spermatogenesis and epididymal sperm maturation in mice. *Arch Androl* 43: 55-66.
121. Davis KL, Martin E, Turko IV, Murad F (2001) Novel effects of nitric oxide. *Annu Rev Pharmacol Toxicol* 41: 203-236.
122. Ahern GP, Klyachko VA, Jackson MB (2002) cGMP and S-nitrosylation: two routes for modulation of neuronal excitability by NO. *Trends Neurosci* 25: 510-517.
123. Foster MW, Stamler JS (2004) New insights into protein S-nitrosylation. Mitochondria as a model system. *J Biol Chem* 279: 25891-25897.
124. Hess DT, Matsumoto A, Kim SO, Marshall HE, Stamler JS (2005) Protein S-nitrosylation: purview and parameters. *Nat Rev Mol Cell Biol* 6: 150-166.
125. Lefièvre L, Chen Y, Conner SJ, Scott JL, Publicover SJ, et al. (2007) Human spermatozoa contain multiple targets for protein S-nitrosylation: an alternative mechanism of the modulation of sperm function by nitric oxide? *Proteomics* 7: 3066-3084.
126. Machado-Oliveira G1, Lefièvre L, Ford C, Herrero MB, Barratt C, et al. (2008) Mobilisation of Ca²⁺ stores and flagellar regulation in human sperm by S-nitrosylation: a role for NO synthesised in the female reproductive tract. *Development* 135: 3677-3686.
127. Bedu-Addo K, Costello S, Harper C, Machado-Oliveira G, Lefièvre L, et al. (2008) Mobilisation of stored calcium in the neck region of human sperm—a mechanism for regulation of flagellar activity. *International Journal of Developmental Biology* 52: 615.
128. Costello S, Michelangeli F, Nash K, Lefièvre L, Morris J, et al. (2009) Ca²⁺-stores in sperm: their identities and functions. *Reproduction* 138: 425-437.
129. Kakizawa S, Yamazawa T, Iino M (2013) Nitric oxide-induced calcium release: activation of type 1 ryanodine receptor by endogenous nitric oxide. *Channels (Austin)* 7: 1-5.
130. Takeshima H, Nishimura S, Matsumoto T, Ishida H, Kangawa K, et al. (1989) Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature* 339: 439-445.
131. Otsu K, Willard H, Khanna V, Zorzato F, Green N, et al. (1990) Molecular cloning of cDNA encoding the Ca²⁺ release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic reticulum. *Journal of Biological Chemistry* 265: 13472-13483.

132. Stoyanovsky D, Murphy T, Anno PR, Kim YM, Salama G (1997) Nitric oxide activates skeletal and cardiac ryanodine receptors. *Cell Calcium* 21: 19-29.
133. Xu L, Eu JP, Meissner G, Stamler JS (1998) Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science* 279: 234-237.
134. Hart JD, Dulhunty AF (2000) Nitric oxide activates or inhibits skeletal muscle ryanodine receptors depending on its concentration, membrane potential and ligand binding. *J Membr Biol* 173: 227-236.
135. Li N, Zou AP, Ge ZD, Campbell WB, Li PL (2000) Effect of nitric oxide on calcium-induced calcium release in coronary arterial smooth muscle. *Gen Pharmacol* 35: 37-45.
136. Heunks LM, Machiels HA, Dekhuijzen PN, Prakash YS, Sieck GC (2001) Nitric oxide affects sarcoplasmic calcium release in skeletal myotubes. *J Appl Physiol* (1985) 91: 2117-2124.
137. Zima AV, Blatter LA (2006) Redox regulation of cardiac calcium channels and transporters. *Cardiovasc Res* 71: 310-321.
138. Zhang Y, Hogg N (2004) The mechanism of transmembrane S-nitrosothiol transport. *Proc Natl Acad Sci U S A* 101: 7891-7896.
139. Kakizawa S (2013) Nitric oxide-induced calcium release: activation of type 1 ryanodine receptor, a calcium release channel, through non-enzymatic post-translational modification by nitric oxide. *Frontiers in endocrinology* 4.
140. Zahradníková A, Minarovic I, Venema RC, Mészáros LG (1997) Inactivation of the cardiac ryanodine receptor calcium release channel by nitric oxide. *Cell Calcium* 22: 447-454.

Citation: López-Úbeda R, Matás C (2015) An Approach to the Factors Related to Sperm Capacitation Process. *Andrology* 4: 128. doi: [10.4172/2167-0250.1000128](https://doi.org/10.4172/2167-0250.1000128)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission

